

Virtual Consultations: Considerations for the Role of Technology During the Covid-19 Pandemic for People With Type Two Diabetes-UK Perspective.

Lauren Marie Quinn, Melanie J Davies, Michelle Hadjiconstantinou

Submitted to: Journal of Medical Internet Research
on: June 22, 2020

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 4

Supplementary Files..... 14

 Figures 15

 Figure 1..... 16

 Figure 2..... 17

 Figure 3..... 18

Virtual Consultations: Considerations for the Role of Technology During the Covid-19 Pandemic for People With Type Two Diabetes-UK Perspective.

Lauren Marie QuinnMBChB, BMedSci, ; Melanie J DaviesMBChB, MD, FRCP, FRCGP, FMedSci, ; Michelle HadjiconstantinouPhD, CPsychol,

Corresponding Author:

Lauren Marie QuinnMBChB, BMedSci,

Phone: +440116 258 4322

Email: lauren.quinn3@nhs.net

Abstract

The covid-19 pandemic has presented unique challenges for people with diabetes, in addition to their high risk stratification for infection. Supporting people with diabetes to self-care has been critical to reduce their risk of severe infection. This global pandemic has presented an opportunity to digitalise diabetes care and rapidly implement virtual diabetes clinics, with the aim of optimising diabetes management and wellbeing, whilst keeping patients safe. We performed a rapid review of the literature to evaluate the feasibility and effectiveness of virtual clinics in diabetes care before and during the covid-19 pandemic, and have combined these findings with our own reflections in practice. We identified examples demonstrating safety and feasibility of virtual diabetes clinics, which aligns with our own clinical experience during the pandemic. The advantages of virtual clinics include reduced treatment burden, improved therapeutic alliances, societal and psychological benefits, and in our experience, innovative solutions to overcome the challenges presented by the transition from in-person to virtual care. We have provided three infographics to illustrate lessons learnt and key recommendations, including steps to establish a virtual diabetes clinic, a checklist guide for health care professionals conducting virtual clinics and a patient guide for making the most of the virtual clinic. It is important to continue adapting to this pandemic and make technology a sustainable option for the future of diabetes care.

(JMIR Preprints 22/06/2020:21609)

DOI: <https://doi.org/10.2196/preprints.21609>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

✓ **No, I do not wish to publish my submitted manuscript as a preprint.**

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

✓ **Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in a full publication, my title and abstract will remain visible to all users.**

Original Manuscript

Viewpoint

Title:

Virtual Consultations: Considerations for the Role of Technology During the Covid-19 Pandemic for People With Type Two Diabetes-UK Perspective.

Abstract [222 words]

The covid-19 pandemic has presented unique challenges for people with diabetes, in addition to their high risk stratification for infection. Supporting people with diabetes to self-care has been critical to reduce their risk of severe infection. This global pandemic has presented an opportunity to digitalise diabetes care and rapidly implement virtual diabetes clinics, with the aim of optimising diabetes management and well-being, whilst keeping patients safe. We performed a rapid review of the literature to evaluate the feasibility and effectiveness of virtual clinics in diabetes care before and during the covid-19 pandemic, and have combined these findings with our own reflections in practice. We identified examples demonstrating safety and feasibility of virtual diabetes clinics, which aligns with our own clinical experience during the pandemic. The advantages of virtual clinics include reduced treatment burden, improved therapeutic alliances, societal and psychological benefits, and in our experience, innovative solutions to overcome the challenges presented by the transition from in-person to virtual care. We have provided three infographics to illustrate lessons learnt and key recommendations, including steps to establish a virtual diabetes clinic, a checklist guide for health care professionals conducting virtual clinics and a patient guide for making the most of the virtual clinic. It is important to continue adapting to this pandemic and make technology a sustainable option for the future of diabetes care.

Keywords:

Diabetes, virtual clinic, technology, covid-19

Introduction

We are in the midst of the covid-19 pandemic, a novel Severe Acute Respiratory Distress Syndrome coronavirus, which has resulted in thousands of deaths worldwide [1,2]. Increased age and underlying health conditions, including diabetes, cardiovascular disease, obesity and hypertension, significantly increase the risk of developing covid-19 infection [3, 4]. Similarly, disease severity may be worsened and deaths are overrepresented in people with diabetes people with diabetes[4, 5]. Evidence shows poor glycaemic control is both associated with and a consequence of covid-19 infection, the latter demonstrated in older persons with type two diabetes [3,5,6]. More recently, evidence has shown that Black, Asian and Minority Ethnic groups are more severely affected, with higher death rates observed from covid-19 infection in this population; high prevalence of diabetes and comorbidities in this sub-group likely contributes to this increased risk [7]. Despite the challenges for people with diabetes, the covid-19 pandemic has presented a valuable opportunity to digitalise diabetes care. Given the importance of maintaining and improving well-being and glycaemic control during this time, evaluation of novel methods to support self-management remotely is critically important. The aim of this paper is one, to explore the evidence for the role of telemedicine to support people with diabetes, during the covid-19 pandemic and beyond; two, to outline the benefits and challenges presented by virtual diabetes care; three, to present our experience of virtual consultations in clinical settings during the covid-19 pandemic; and (4) to share lessons learnt to assist researchers, clinicians and people with diabetes when integrating technology in diabetes care.

Impact of Social Distancing/Shielding on Diabetes Care and Well-being

With diabetes being classed as a high risk group by the government, it is important that people with diabetes take care of their health now more than ever. people with diabetes are advised to practice social distancing (such as working from home or self-isolating) and adhere to national recommendations around frequent handwashing, to avoid infectious contacts and to abstain from non-essential travel by working from home [8]. In some cases, extremely vulnerable people with diabetes are advised to undertake 'shielding'. Additional guidance, specifically for people with diabetes, focusses on self-management strategies, which help to boost innate immunity for primary prevention [8,9]. The American Association of Diabetes Educators recommend the following seven self-care activities: keeping physically active, healthy diet, following medication regimen, blood glucose monitoring and problem solving, reducing risk of complications and self-empathy; all of which are endorsed for people with diabetes to reduce their risk and severity of covid-19 [10].

Interestingly, for people with diabetes during the pandemic, their engagement with these self-care activities has been highly variable, with some being able to focus more time on their diabetes management and undertake increasing amounts of self-care. For example, Bonora *et al.* [11] showed in a small study that glycaemic control improved for people with type one diabetes during lockdown who self-isolated compared to those who continued working [11]. However, for others, the significant change in lifestyle presented by lockdown has been detrimental to their health and well-being. Barriers specifically for people with diabetes have included difficulty accessing healthy foods because of restricted shopping, and bulk buying; inability to access medications; restricting physical activity to once per day in the local area; and being unable to attend face-to-face appointments with their diabetes care providers [8,9]. With this in mind, diabetes health care professionals have valid concerns that glycaemic control, quality of life, self-management and well-being can be significantly jeopardised during social distancing and shielding, posing considerable risks to people with diabetes, both in the short and longer-term. Lockdown, social distancing, shielding, and the abundance of misinformation in the media, also present additional stressors, which may further exacerbate underlying depression and anxiety [12], conditions that are already highly prevalent in the diabetes

population [13].

Digitalisation of Diabetes Care: Benefits and Challenges

Digital Consultations

Technology has been increasingly integrated into diabetes education and care in modern time, for example through applications (apps), computer-based or web-based education, and telemedicine [14]. The NHS (National Health Service) long-term plan set out to increase digitalisation within the NHS [15, 16]. This included roll-out of virtual or non-face-to-face clinics, with the aim of reducing face-to-face appointments by one third over the next five years [16]. This non-face-to-face activity may be synchronous or asynchronous, meaning a direct or indirect line of communication with a health care professional respectively [17,18]. Synchronous activity would be a video or telephone consultation, whereas asynchronous may be monitoring an email or tracking system and responding to patients' questions through these platforms. The benefits of non-face-to-face appointments are multifactorial and include the opportunity for better care and more joined up patient care pathways, as well as cost savings, and reduced impact on the environment.

Studies evaluating virtual clinics prior to the covid-19 pandemic have demonstrated feasibility, accessibility, comparable effectiveness to in-person consultations and safety [19]. A multicenter mixed methods study evaluating video consultations in diabetes care showed that video consultations were shorter in duration and the people with diabetes did relatively more talking than the health care professional [20,21]. Although from a management perspective video consultations were favoured, there were significant barriers to uptake from the teams implementing them, because of the significant changes introduced to their usual way of working and the care processes, systems and pathways [21]. A Cochrane review analysed twenty one low to high quality studies comparing telemedicine to usual care in people with diabetes and found that improvement in glycaemic control was variable, but LDL and blood pressure were more effectively lowered by telemedicine approaches compared to usual care [22]. Further studies have evaluated virtual clinics for diabetes care in type one and type two diabetes and have reported improved biochemical parameters, including HbA_{1c} (glycosylated haemoglobin) [23]. However many of these studies combined interventions, such as synchronous and asynchronous programmes, rendering it difficult to delineate efficacy of the individual interventions [23]. Studies focussing on synchronous video consultations have been of short duration with a limited sample size [24]. Also, how these findings relate to virtual clinics being implemented during and because of a global pandemic must be considered.

During the covid-19 pandemic, telemedicine has been widely adopted globally in order to reduce exposure and need for people with diabetes to come into hospital, whilst maintaining care standards for people with chronic conditions. For example, virtual consultations have been implemented to triage patients suspected of covid-19 in primary care and to establish need to come into hospital [25-27]. There have been other examples of successful adoption of virtual care for people with newly diagnosed type one diabetes [28], and for people with long standing type one diabetes. A small Italian study showed that for people with type one diabetes who were not working during the pandemic and using continuous glucose monitoring, their time in range significantly improved from 54% to 65%, and this was attributed to decreased hyperglycaemia [11]. Blood glucose variability and average glucose readings from continuous glucose monitoring also significantly improved in this study. In contrast to patients who continued working, there was no difference in glycaemic control [11]. Early reports suggest that virtual clinics are feasible, with some centres increasing virtual clinic consultations from 1% of all consultations prior to covid-19 to 70% afterwards and it is technology that has made this rapid transformation possible [29-31].

A recent linguistic ethnographic study has shown that video consultations among people with a long-term condition and their clinician was found to be effective [26]. When patients experienced technical or operational issues with their video equipment or internet connection, they generally

found a solution to resolve the problem. Familiarity and experience with technology helped in situations like these. Technical interruptions and delays of connectivity either on patient or clinician device were clearly evident in the study, nevertheless, remote physical examinations were able to be carried out, allowing the patient and/or carer to take an active role in the visual/audio adjustments to their device [26].

Benefits of Virtual Clinics

Patient Safety

In line with social distancing advice and minimising risk for people with diabetes, virtual clinics bring specialist care to their home. The virtual clinic is the ideal solution to enable patients to access specialist care but without unnecessarily exposing them to a high risk environment, or close contact with other staff or patients. This is particularly pertinent given the new data supporting the number of covid-19 infections developed whilst being in hospital [32]. Reduced burden of treatment: Improved accessibility and overall experience

Virtual clinics increase accessibility to specialist diabetes care. For example, those shielding can still attend a virtual appointment, those working can more easily take the time out from their day (for example over a coffee break) and those who normally are unable to travel, no longer need to. In our experience, we have found that patients who regularly 'Did Not Attend' appointments, are now attending their virtual clinic appointments. The virtual clinic may also be an improved patient experience because people with diabetes are not required to take additional travel to the hospital for regular appointments, which can significantly reduce the burden of treatment that some may experience with their diabetes care. They are also not brought into a clinical environment, which for some may be a daunting place. The virtual clinics, may provide a solution – video or phone calls may provide a relaxed environment for some people with diabetes, as they are within their own familiar surroundings, which may create a more relaxed atmosphere for their virtual appointment. From our clinic team experience of using virtual consultations with people with diabetes, it is evident that some feel comfortable to show the Health Care Professional (HCP) their house environment, exercise regimen and introduce them to their family members, within professional boundary. This offers a unique insight into environmental or interpersonal factors, which may influence people with diabetes's self-care activities. Long-term, these factors may be considered when discussing the people with diabetes care plan and treatment goals. The opportunities presented in virtual clinics to deliver on increasingly person-centred and individualised care cannot be underestimated.

Improved Therapeutic Alliance and Consultation Dynamics

Virtual consultations could significantly shift the locus of control of the people with diabetes, which would complement the philosophy around the importance of self-management and self-care in diabetes. Virtual consultations could provide an opportunity for shared care between the people with diabetes and the HCP. Multiple members of the diabetes multidisciplinary team (MDT) may come together for the video consultation, to make collaborative decisions with the people with diabetes. From our experience, this results in a more effective consultation, which can be more time-effective. The people with diabetes is benefiting from MDT input, with reduced time commitment for patients and professionals alike. Physical examination, for example conducting a diabetes foot examination, can also be performed. This is particularly important in the context of diabetes foot disease, where people with diabetes can be triaged in virtual clinic to determine need for admission to hospital based on the video examination findings. Technical barriers would be expected when setting up virtual consultations, thus, we have shared a checklist that we developed for setting up virtual clinic (Figure 1).

Societal and Psychosocial Benefits

Virtual clinics reduce the need for travel to and from hospitals, which is a significant benefit given 20% of United Kingdom (UK) traffic is attributed to be health care related [15,16]. Also, virtual clinics mean less time is missed from work, which can be a recurrent issue for people with diabetes, so this reduction in treatment burden is hugely advantageous.

With social isolation becoming more prevalent than ever during the current climate, it is imperative to acknowledge the positive impact that remote consultations may have on individuals who require practicing social distancing and shielding. Virtual consultations, whether these are conducted by telephone or video, provide people with diabetes the opportunity to virtually connect with their HCP, mitigating the psychological effects of social isolation.

Overcoming the Challenges With Virtual Care

Elements of the routine diabetes clinic, for example checking blood pressure, sampling the urine, HbA_{1c}, spot checks and physical examination are not possible in the same way in a virtual consultation. However, given the importance of these methods to screen for diabetes complications, alternative strategies can be developed to make them possible. For example, people with diabetes can get their own blood pressure monitor. This may likely lead to more accurate readings, with the absence of white coat hypertension. Also, urine samples can be delivered locally to general practices and examination can still be performed via video consultation. For example, diabetes foot disease can be screened for and people with diabetes can be triaged to determine need for further assessment.

Patient Preference

The people with diabetes can choose whether to have a telephone or video consultation, to reflect variability in access to technology and systems for video recording and to ensure access to care services. However, virtual care is a significant change from the usual hospital attendance for their diabetes care, thus, some people with diabetes would still prefer a face-to-face consultation. However, in light of the covid-19 pandemic, people with diabetes have readily transitioned to virtual methods and it is hoped that by familiarisation with these new consultation methods, this may increase uptake for virtual care in the longer term. There are also concerns that people with diabetes may find it difficult to build rapport with their HCP in a virtual clinic, but in our experience, the opposite is true. people with diabetes have been far more relaxed in the virtual clinic and able to speak more openly with their diabetes team, which is essential to improve their diabetes management. But this is dependent on high quality video-image and sound, highlighting the essential role of technology to optimise communication between the people with diabetes and HCP.

Engaging the Multidisciplinary team

Engaging professionals and the wider team with virtual care can be a challenge in itself. This is a significant change for the diabetes team as well as the patient, and whole new processes and pathways have needed to develop rapidly to manage the people with diabetes's care remotely (Figure 2). For example, ways in which clinic outcomes are actioned have changed and the wider clinical and non-clinical teams have needed to collaborate to generate these new systems. It is essential that the whole team are committed to the virtual care approach, because this ensures the delivery of safe, high quality care. It is anticipated that with increased familiarity and established systems, HCP will be more open to adopt virtual care into their routine practice and overcome the natural aversion to change how it has always been.

The Future of Virtual Clinics

Anecdotally, in our experience the virtual clinics have been feasible and accessible, with high patient satisfaction. Virtual clinical consultations offer a different kind of benefit compared to conventional

face-to-face appointments, particularly around convenience, logistics, cost-effectiveness and clinician-patient dynamic/relationship [26]. However, we must acknowledge the pitfalls of these new modes of communication and the challenges that may lie ahead with the clinical quality and safety of appointments.

Building on these developments, we are looking to make virtual clinics sustainable for the long-term. This is in line with the NHS long-term plan, to make better use of data and digital technology, in the next five years [15, 16]. In a time of a pandemic, individualised care is more important than ever and virtual clinics provide a readily accessible solution to facilitate this. Having applied virtual clinics in our setting for the last two months, lessons we have learnt include: (i) the importance of integrating multiple members of the MDT into the one virtual consultation; (ii) avoiding the checklist approach and instead focusing on an individualised, person-centred consultation; (iii) and acknowledging that video consultation may be preferred to telephone because of the additional benefits of human contact, body language and the opportunity to gain better insight into the people with diabetes's lifestyle and livelihood to tailor medical support accordingly. In our experience, virtual clinics may be better suited to individuals with longstanding diabetes and where possible should be performed by a professional with whom they have already built a strong rapport with.

Virtual clinics could be an alternative option, however, there are key elements that must be considered to make the consultation as efficient as possible: firstly, even though virtual consultations would require less resources compared to face-to-face, organisational factors prior to virtual consultation would still be required in order to book and record a clinic appointment; secondly, logistical and administrative factors must be integrated within the NHS system; a system which for so many years has been based on delivering face-to-face patient services. We are programmed to deliver our outpatient services in a 'traditional' way, and therefore would anticipate a colossal challenge adapting this existing pathway to a more digital-focused platform.

By acknowledging the complexity of integrating virtual consultations, we also acknowledge the challenges that may come with technology, in terms of the security and safety of every patient and HCP. Data protection and privacy is of critical importance; the technology, software and programmes used in virtual clinics must be encrypted and meet General Data Protection Regulation and information governance standards to maintain patient confidentiality at all times. We acknowledge that virtual consultations are not for everyone, however, providing options to people, enables them to choose an approach that is tailored around their diabetes needs, and also their lifestyle demands; with the aim of reducing the burden of treatment that so many people with diabetes may be experiencing. We aim to prepare our patients to attend virtual consultations, by creating a safe environment and respecting their safety (Figure 3). But to also create a digital platform that would integrate within our current NHS system.

Conclusion

Virtual consultations may become a necessity following this pandemic. The current system pressures with covid-19 have led to numerous challenges to the delivery of routine diabetes care and education. Despite the relative lack of data to support virtual care, in the face of adversity, these virtual measures have been imperative to maintain a line of communication with people with diabetes and to support self-management and self-care remotely. With the right infrastructure and systems in place, technology is the key to evolution in diabetes care and virtual consultations could effectively be embedded into the UK national routine diabetes care. In the time that we find ourselves, virtual clinics may be an ideal platform to reduce social isolation, encourage self-management remotely, and reduce burden of treatment in a less intrusive manner. The covid-19 outbreak will shift the culture of health care across the world and the way we interact within clinical settings will gradually change, to ensure that care can be delivered within social distancing rules

Acknowledgments and author contributions

LQ, MJD and MH had substantial contributions to conception and design, drafting the article and revising it critically for important intellectual content; and final approval of the version to be published.

Conflicts of interest

None declared.

Abbreviations

NHS	National Health Service
HbA _{1c}	glycosylated haemoglobin
HCP	health care professional
MDT	multidisciplinary team
UK	United Kingdom

References

1. World Health Organization. 2020. Novel Coronavirus – China. [online] Available at: <https://www.who.int/csr/don/12-january-2020-novel-coronavirus-china/en/> [Accessed 19 July 2020]
2. GOV.UK. 2020. Staying At Home And Away From Others (Social Distancing). [online] Available at: <https://www.gov.uk/government/publications/full-guidance-on-staying-at-home-and-away-from-others> [Accessed 19 July 2020]
3. Shenoy A, Ismail M, Bajaj M. Diabetes and covid-19 : a global health challenge. *BMJ Open Diabetes Res Care* 2020;8(1):e001450. PMID:32345580
4. Bornstein SR, Rubino F, Khunti K, Mingrone G, Hopkins D, Birkenfeld AL, et al. Practical recommendations for the management of diabetes in patients with COVID-19. *Lancet Diabetes Endocrinol* 2020;8(6):546-550. PMID:32334646
5. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323(13):1239-1242. PMID:32091533
6. Chen G, Li L, Chen G, Li L. Blood glucose levels in elderly subjects with type 2 diabetes during COVID-19 outbreak: a retrospective study in a single center. *medRxiv* 2020. doi: <https://doi.org/10.1101/2020.03.31.20048579>
7. The King's Fund. 2020. Ethnic Minority Deaths And Covid-19. [online] Available at: <https://www.kingsfund.org.uk/blog/2020/04/ethnic-minority-deaths-covid-19> [Accessed 19 July 2020]
8. Gupta R, Ghosh A, Singh AK, Misra A. Clinical considerations for patients with diabetes in times of COVID-19 epidemic. *Diabetes Metab Syndr Clin Res Rev* 2020;14(3):211-212. PMID:32172175
9. Banerjee M, Chakraborty S, Pal R. Diabetes self-management amid COVID-19 pandemic. *Diabetes Metab Syndr Clin Res Rev* 2020;14(4):351-354. PMID:32311652
10. Diabeteseducator.org. 2020. [online] Available at: <https://www.diabeteseducator.org/docs/default-source/practice/practice-resources/position-statements/aade7-self-care-behaviors-position-statement.pdf?sfvrsn=6> [Accessed 19 July 2020]
11. Bonora BM, Boscari F, Avogaro A, Bruttomesso D, Fadini GP. Glycaemic control among people with type 1 diabetes during lockdown for the SARS-CoV-2 outbreak in Italy. *Diabetes Ther* 2020;11(6):1-11. PMID:32395187
12. Kar P. Partha Kar: Covid-19 - We must keep faith in our experts. *BMJ* 2020;368. PMID:32193174
13. Semenkovich K, Brown ME, Svrakic DM, Lustman PJ. Depression in type 2 diabetes mellitus: Prevalence, impact, and treatment. *Drugs* 2015;75(6):577–87. PMID:25851098
14. Cahn A, Akirov A, Raz I. Digital health technology and diabetes management. *J Diabetes* 2018;10(1):10–17. PMID:28872765
15. Plan, N., 2020. Chapter 5: Digitally-Enabled Care Will Go Mainstream Across The NHS. [online] NHS Long Term Plan. Available at: <https://www.longtermplan.nhs.uk/online-version/chapter-5-digitally-enabled-care-will-go-mainstream-across-the-nhs/> [Accessed 19 July 2020].
16. Alderwick H, Dixon J. The NHS Long Term Plan. *BMJ* 2019;364:l84. PMID:30617185
17. UCLPartners. 2020. How-To Guide: Non Face-To-Face Clinics | Uclpartners. [online] Available at: <https://uclpartners.com/non-face-to-face-clinics-resource/> [Accessed 19 July 2020].
18. England.nhs.uk. 2020. [online] Available at: <https://www.england.nhs.uk/wp-content/uploads/2020/01/online-consultations-implementation-toolkit-v1.1-updated.pdf> [Accessed 19 July 2020]

2020]

19. Armstrong N, Hearnshaw H, Powell J, Dale J. Stakeholder perspectives on the development of a virtual clinic for diabetes care: Qualitative study. *J Med Internet Res* 2007;9(3):e23. PMID:17942385
20. Greenhalgh T, Vijayaraghavan S, Wherton J, Shaw S, Byrne E, Campbell-Richards D, et al. Virtual online consultations: Advantages and limitations (VOCAL) study. *BMJ Open* 2016;6(1):1–13. PMID:26826147
21. Shaw S, Wherton J, Vijayaraghavan S, Morris J, Bhattacharya S, Hanson P, et al. Advantages and limitations of virtual online consultations in a NHS acute trust: the VOCAL mixed-methods study. *Heal Serv Deliv Res* 2018;6(21):1–136. PMID:29939519
22. Flodgren G, Rachas A, Farmer AJ, Inzitari M, Shepperd S. Interactive telemedicine: Effects on professional practice and health care outcomes. *Cochrane Database Syst Rev* 2015 (9):CD002098. PMID:26343551
23. Dixon RF, Zisser H, Layne JE, Barleen NA, Miller DP, Moloney DP, et al. A virtual type 2 diabetes clinic using continuous glucose monitoring and endocrinology visits. *J Diabetes Sci Technol* 2019;1932296819888662. PMID:31762302
24. Jennings A, Powell J, Armstrong N, Sturt J, Dale J. A virtual clinic for diabetes self-management: pilot study. *J Med Internet Res* 2009;11(1):1–10. PMID:21821504
25. Greenhalgh T, Koh GCH, Car J. Covid-19: A remote assessment in primary care. *BMJ* 2020;368:1–5. PMID:32213507
26. Shaw SE, Seuren LM, Wherton J, Cameron D, A’Court C, Vijayaraghavan S, et al. Video consultations between patients and clinicians in diabetes, cancer, and heart failure services: linguistic ethnographic study of video-mediated interaction. *J Med Internet Res* 2020;22(5):e18378. PMID:32391799
27. The Health Foundation. 2020. Using Virtual Consultations In The Fight Against COVID-19 | The Health Foundation. [online] Available at: <https://www.health.org.uk/news-and-comment/newsletter-features/using-virtual-consultations-in-the-fight-against-covid-19> [Accessed 19 July 2020]
28. Garg SK, Rodbard D, Hirsch IB, Forlenza GP. Managing new-onset type 1 diabetes during the COVID-19 pandemic: challenges and opportunities. *Diabetes Technol Ther* 2020;22(6):1–9. PMID:32302499
29. Jones MS, Goley AL, Alexander BE, Keller SB, Caldwell MM, Buse JB. Inpatient transition to virtual care during COVID-19 pandemic. *Diabetes Technol Ther* 2020;22(6):6–10. PMID:32396395
30. Wosik J, Fudim M, Cameron B, Gellad ZF, Cho A, Phinney D, et al. Telehealth transformation: COVID-19 and the rise of virtual care. *J Am Med Informatics Assoc* 2020;27(6):957-962. PMID:32311034
31. Javaid M, Haleem A, Vaishya R, Bahl S, Suman R. Industry 4.0 technologies and their applications in fighting COVID-19 pandemic. *Diabetes Metab Syndr Clin Res Rev* 2020;14(4):419-422. PMID:32344370
32. GOV.UK. 2020. Reducing The Risk Of Transmission Of COVID-19 In The Hospital Setting. [online] Available at: <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/reducing-the-risk-of-transmission-of-covid-19-in-the-hospital-setting> [Accessed 19 July 2020]


Supplementary Files

Figures

Checklist for setting up a virtual clinic.



How to conduct a virtual clinic: a health care professional's guide.



Making the most of your virtual diabetes clinic: a patient's guide.

