

# **Application of mobile health to support the elderly during the COVID-19 outbreak: A systematic literature review**

Saeedeh Fehrest, Elham Monaghesh

Submitted to: JMIR mHealth and uHealth  
on: January 04, 2021

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

## ***Table of Contents***

---

Original Manuscript.....	4
Supplementary Files.....	20

Preprint  
JMIR Publications

# Application of mobile health to support the elderly during the COVID-19 outbreak: A systematic literature review

Saeedeh Fehrest<sup>1</sup>; Elham Monaghesh<sup>2</sup> MSc

<sup>1</sup>Department of Health Economics and Management, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran Tehran IR

<sup>2</sup>Department of Health Information Technology, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran. Tabriz IR

## Corresponding Author:

Saeedeh Fehrest

Department of Health Economics and Management, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran  
Tehran University of Medical Sciences  
Tehran  
IR

## Abstract

**Background:** The COVID-19 outbreak has affected the elderly's physical and mental health. The application of information and communication technology, such as mobile health (m-health), can play a significant role in combating this pandemic by changing the behavior and lifestyle of the elderly during this time of crisis.

**Objective:** This systematic review aimed to synthesize m-health capabilities in providing health services to the elderly during the COVID-19 pandemic, and to identify the factors associated with the success of these tools.

**Methods:** To find the relevant studies, a search was conducted in PubMed, Web of Science, Scopus, ProQuest, and Google Scholar. The inclusion criteria were: studies in English that used m-health intervention in all aspects of elderly healthcare during the COVID-19 outbreak, published in peer-reviewed journals from 31 December 2019, and had any research design and methodology. Two authors independently took all the steps of this review, and finally performed narrative synthesis to report the findings.

**Results:** Our initial search identified 421 studies, of which 10 met the inclusion criteria. The data analysis showed that all the m-health interventions had positive effects on the health of the elderly. The m-health services for the elderly during the current pandemic were used for therapy, information provision, self-help, monitoring, and mental health consultation purposes. The results also indicated that various factors affected the elderly's use of m-health tools.

**Conclusions:** The application of m-health keeps the elderly and healthcare providers safe, accelerates health service provision, reduces the costs of service delivery, and decreases the risk of morbidity and mortality during the COVID-19 outbreak. The successful use of m-health tools for the elderly in health programs during the current crisis greatly depends on supporting the elderly and overcoming the barriers.

(JMIR Preprints 04/01/2021:26929)

DOI: <https://doi.org/10.2196/preprints.26929>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in a peer-reviewed journal, I will be able to remove the preprint label and make the full manuscript available to all users.

## Original Manuscript

## Application of mobile health to support the elderly during the COVID-19 outbreak: A systematic literature review

Sara Abbaspur-Behbahani<sup>1</sup>, Elham Monaghesh<sup>2</sup>, Saeedeh Fehrest<sup>3\*</sup>

### Abstract

**Background:** The COVID-19 outbreak has affected the elderly's physical and mental health. The application of information and communication technology, such as mobile health (m-health), can play a significant role in combating this pandemic by changing the behavior and lifestyle of the elderly during this time of crisis.

**Objectives:** This systematic review aimed to synthesize m-health capabilities in providing health services to the elderly during the COVID-19 pandemic, and to identify the factors associated with the success of these tools.

**Methods:** To find the relevant studies, a search was conducted in PubMed, Web of Science, Scopus, ProQuest, and Google Scholar. The inclusion criteria were: studies in English that used m-health intervention in all aspects of elderly healthcare during the COVID-19 outbreak, published in peer-reviewed journals from 31 December 2019, and had any research design and methodology. Two authors independently took all the steps of this review, and finally performed narrative synthesis to report the findings.

**Results:** Our initial search identified 421 studies, of which 10 met the inclusion criteria. The data analysis showed that all the m-health interventions had positive effects on the health of the elderly. The m-health services for the elderly during the current pandemic were used for therapy, information provision, self-help, monitoring, and mental health consultation purposes. The results also indicated that various factors affected the elderly's use of m-health tools.

**Conclusion:** The application of m-health keeps the elderly and healthcare providers safe, accelerates health service provision, reduces the costs of service delivery, and decreases the risk of morbidity and mortality during the COVID-19 outbreak. The successful use of m-health tools for the elderly in health programs during the current crisis greatly depends on supporting the elderly and overcoming the barriers.

**Keywords:** M-health, Coronavirus, COVID-19, Outbreak, Elderly

### Introduction

On December 31, 2019, a cluster of a genus of the Coronaviridae family was officially reported from Wuhan, Hubei Province, China, which caused the novel Coronavirus disease (COVID-19) (1, 2). This infection widely spread to other regions of the world, and primary evidence revealed that elderly people and those with underlying medical problems (e.g. cardiac diseases, diabetes, hypertension, and cancers) are more susceptible to the disease in its most intensive form (3).

As the older population has a much higher risk of mortality compared to younger people during the COVID-19 pandemic, it is emphasized that the elderly should be protected with social distancing or,

<sup>1</sup> Department of Health Economics and Management, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Department of Health Information Technology, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran.

<sup>3</sup> Department of Health Economics and Management, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

\*Corresponding author: Saeedeh Fehrest. Telephone number: +933758700. E-mail address: [sa.fehrest666@yahoo.com](mailto:sa.fehrest666@yahoo.com). ORCID: 0000-0002-9919-8875

if necessary, social isolation (4, 5). Loneliness and social isolation may lead to many problems such as depression, cognitive dysfunction, disability, cardiovascular diseases, and increased mortality among older adults (6). The ongoing COVID-19 outbreak in the world has highlighted the need for mobile technology solutions to diminish the risk of cross-contamination caused by close contact (4, 7, 8). Smart technology (e.g. mobile-based technology) has been important for exploring meaning in life, self-care, consultation, and creating a positive feeling for confined older adults (9, 10).

The term m-health refers to smart phone applications and is increasingly used in the healthcare sector (11). In another definition, m-health is proposed as a technology which may be either short-distance or long-distance, or be system-driven (12). More specifically, m-health is defined as the provision of health intervention using mobile-based tools, including phone calls, video consultation, wearable devices, personal digital assistants, tablets, and personal computers (PCs) (13). The main concept of m-health is that it can provide health services, anytime and anywhere, to anyone, by removing geographical, temporal, and other barriers (14). It is also reported that mobile-based tele-visits can improve access for an estimated 6.3 million older people who are inexperienced with technology tools or have visual problems (15). Remote interventions are potentially effective and utilized by older people, but healthcare providers should ensure that digital services are suitable for the older demographic and provide both the technology and the technical support needed for interventions in the current pandemic (16).

Previous studies have reported the potential role of m-health tools in providing long-term support and health promotion for the elderly (17-19). However, there is no review study on the role of m-health in improving the health of the elderly during the COVID-19 pandemic. Thus, this study aimed to synthesize the literature on m-health capabilities in providing health services for the elderly during the COVID-19 outbreak, and to identify the factors associated with the success of these tools. The research questions were:

- What are the capabilities of available m-health services during the COVID-19 pandemic for the elderly?
- What factors are associated with the success of m-health tools for the elderly during this global emergency?

## **Materials and methods**

### **Study design**

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. All the steps of this review, namely the search in databases, study selection, quality evaluation, data extraction, and data analysis, were independently performed by two authors (S.AB. and S.F.).

### **Bibliography search strategy**

To identify studies on m-health tools used to help the elderly which were published during the COVID-19 pandemic, a comprehensive search was conducted in PubMed, ProQuest, Web of Science, Scopus, and Google Scholar, since the appearance of COVID-19 in Wuhan, Hubei Province, China, until December 31, 2019. Given the aim for this review, the search was conducted on titles and abstracts via a combination of relevant keywords: (1) COVID19, COVID-19, Coronavirus, Novel Coronavirus, 2019-nCoV, Wuhan Coronavirus, SARS-CoV-2, SARS2, (2) Aging, senior, aged, elderly, old\* (3) M-health, m-health, handheld computer, mobile phone, smartphone, mobile application, mobile app, cellular phone, mobile device, mobile technology, SMS, or text message. Hand searching in web-based resources was also accomplished, and finally, the selected articles' references were reviewed to identify additional studies (reference by reference). EndNote x7 was employed for the management of the sources. The search strategy was adapted for each bibliography database, and the search was conducted on December 1, 2020. As an example, the full search strategy on PubMed is given in Table 1.

Table 1. Search Terms and Strategy Used in PubMed

**Eligibility criteria**

All the included studies had to meet the following inclusion criteria:

- Study types: interventional, observational, modeling, or case studies
- Date of publication: December 31, 2019, to December 1, 2020
- Language: English
- Place of publication: peer-reviewed journals
- Population: elderly people (aged >60 years)
- Analysis criteria: studies on the use of m-health to support the elderly living at home or at a healthcare setting during the COVID-19 outbreak

Studies about other technologies (e.g. Internet of medical things or IoMT), studies which had not stated the primary objective of the intervention or reported incomplete information, conference abstracts, review articles, letters to the editor, and opinion papers were excluded.

**Study selection**

In the process of study selection, first, a screening was performed to remove duplications from the datasets retrieved from the mentioned databases, and the papers were reviewed on the basis of their titles. Subsequently, relevant papers were further extracted in the screening process by screening based on abstracts. After initial screening, the remaining papers were screened in the third phase by reviewing their full text according to predefined inclusion and exclusion criteria.

**Data extraction and quality assessment**

The authors used the Microsoft Word Application (version 2016) to design a standard data extraction form. The following information on the eligible studies was extracted: study identification (authors and publication year), study design, country, population (N), the device or technology used, intervention, and key findings.

We assessed the methodology of each included study in accordance with the Critical Appraisal Skills Program (CASP) checklists. The included studies were also divided into three categories of weak, medium, and good on the basis of their quality assessment scoring.

**Evidence analysis**

Owing to the heterogeneity of the designs and objectives of the included studies, a narrative synthesis was performed to summarize the overall evidence by comparing and contrasting the data. The authors reached a consensus on the findings' categorized data according to the details, and qualitatively described and presented the results.

**Results****Selection of articles**

All the identified studies were imported into the EndNote X7 bibliography management software, which retrieved 312 studies after the removal of duplicates. After screening the title and abstracts of the studies, 47 relevant studies with a full text were appraised according to the inclusion and exclusion criteria. After reading the full text of the articles, the authors included nine papers for evidence synthesis. One study was also added after reference screening (reference by reference) and, finally, 10 articles were included for final analyses. The details of the screening procedure are

depicted in the PRISMA flow diagram in Figure 1.

Figure 1: Search results and publication selection procedures

### Characteristics of the included articles

Table 2 presents details of the included studies. This corpus comprised studies that had the following distribution: six studies on the capability of m-health applications in elderly life during this emergency condition, and four studies reporting factors associated with the success of m-health to ensure the application of m-health tools. The 10 studies that met the eligibility criteria had been conducted in various settings, including the Netherlands, China, Canada, United Kingdom, France, Sweden, Italy, Mexico, Portugal, and Spain. Eight studies (80%) possessed good quality, two (20%) had medium quality, and no study was excluded on the basis of the level of quality.

Table 2: General characteristics and key findings of the Included Studies

### Principal findings

Our review identified precious data on how m-health technologies have been used as a means of supporting the elderly during the COVID-19 outbreak. The data analysis showed that health organizations, researchers, and companies suggest mobile-based tools as a method of providing information, facilitating communication, remotely monitoring vital signs, and limiting the number of times healthcare professionals and patients need to come into close contact during the COVID-19 pandemic. Furthermore, the m-health technology (text, apps, calls, etc.) is used for another variety of purposes, e.g. connecting older people with their family and friends during social distancing, purchasing masks, showing the green code for traveling, storing health information, managing chronic diseases, making appointments, and creating a upbeat feeling in the elderly. According to these studies, systems supporting the elderly range from simple tablet- and telephone-based systems to mostly app-based platforms. Advantages provided by m-health tools include the ability to massively implement them in all healthcare settings, practicing social distancing, decreasing the transmission of diseases to the elderly as high-risk individuals, and reducing the mortality and morbidity rate among the elderly population during this pandemic. Even if some challenges exist, the relevance and impact that m-health solutions can offer during COVID-19 are apparent, as they allow health service providers to effectively manage the elderly's well-being to prevent pressures on the healthcare system. Generally, m-health tools for the elderly during the current pandemic are used for therapy, information provision, self-help, monitoring, and mental health consultation purposes.

The reviewed studies reported factors associated with the use of m-health tools by older adults during the COVID-19 pandemic. Most of the reported factors, e.g. the availability of m-health tools, internet access, and the elderly's willingness and capability, are important determinants of the use of m-health tools. Meanwhile, some of the largest facilitators in the adoption of m-health technologies include support from the government and family, empowerment of the elderly, addressing the digital problems in deprived areas, and increasing accessibility to simple mobiles or tablets, which can be directly related to the use of m-health tools by the elderly to improve their health and combat COVID-19 outbreak.

### Discussion

This systematic review exhibited evidence on the scope of m-health capabilities in providing health services for the elderly at the time of COVID-19 outbreak, as well as the factors associated with the success of these tools. The majority of the included studies (60%) aimed to design and develop a mobile-based app to address the health and wellbeing of the elderly by addressing specific COVID-19 issues. These included apps, web-portals, video consultations, and telephone outreach, with the study population mainly consisting of older people with social isolation and residents of independent elderly, most often in need of communication and mental health promotion. This is not surprising, as many psychological problems during quarantine can be prevented through positive communication



or by improving the lives of older adults (23, 26).

The global confinement during the COVID-19 pandemic has highlighted the effect of social isolation and loneliness on the older population (30-32). In this global isolation, European countries (United Kingdom, France, Sweden, etc.) have designed and deployed health mobile apps to assist families and peer groups in maintaining contact with older people. These mobile apps can tackle new and urgent problems in managing COVID-19, and can be applied as a valuable tool for connecting vast population geographies, especially the elderly (20). An important issue to consider when using these online services during the pandemic is the willingness and ability of older people to use m-health tools. For example, older adults hospitalized in healthcare settings were able to independently complete phone calls rather than video calls, and tended to make phone calls more often than video calls (33).

Our results showed that m-health tools have been potential solutions for managing the problems of the elderly during the COVID-19 pandemic. For example, according to the study conducted in the USA, the COVID mobile app is an innovative tool for the management of long-term care facility and can help provide a new channel for real-time communication, promote the early detection of suspected COVID-19 cases, and decrease the number of deaths and confirmed cases of COVID-19 in long-term care settings (28).

COVID-19 has been a crisis of huge proportions for health delivery services associated with chronic diseases, especially among the elderly, and online services have been recognized as a potential solution for fighting these diseases (25, 29). Van Dijk et al. in the Netherlands has designed an online Psychotherapy Day-treatment Program which provides online psychotherapy sessions, with activities and movements, and mindfulness exercises (25). This online mobile-based program can help avoid direct physical contact, reduce the risk of exposure to infections, and prevent the potential transmission of infections to the elderly and healthcare providers during the current emergency conditions. Generally, mobile-based online tools can help healthcare institutional staff and clinicians address the COVID-19 pandemic by providing a channel for real-time communication (28).

In this review, we also presented evidence on the factors associated with the use of m-health tools for health promotion among older adults to overcome COVID-19. These factors are important determinants of the use of m-health tools by the elderly. According to Cenzer et al., there are some lingering issues with the capability and willingness to participate in video visits, particularly among older adults residing in rural areas. Factors affecting the use of VA Video Connect (VVC) by the elderly include the Internet access, email access, access to devices with a camera, and the capability/willingness to participate in VVC appointments (22). To incentivize the elderly and increase their usage of m-health tools, the studies suggest replacing computers with mobile internet, providing government and family support, addressing the digital divide, and making educational and policy efforts to combat the COVID-19 pandemic are essential (22, 24). Despite the huge potential of m-health tools in the promotion of elderly health, it is necessary that the government and health policy-makers solve the mentioned challenges. This can help the management of various problems that have emerged, and improve the control of COVID-19 among the elderly population.

### Future research

The majority of the databases that we reviewed published studies about the role of m-health tools in the older population, but only a few reports focused on a specific older group during the COVID-19 pandemic. Accordingly, health organizations and researchers should pay more attention to this domain so that the elderly can be taken care of in emergency situations such as COVID-19 with the help of m-health tools. More research has to be conducted to develop and evaluate new methods, and social worker training programs should be provided to ensure that older people can access these services. During this time of crisis, more evidence on the efficacy and cost-effectiveness of m-health tools in promoting health and well-being of older adults is required. It is recommended that future studies determine how to set up m-health in healthcare settings (e.g. home nursing and elderly care

centers), and assess the elderly and healthcare professionals' satisfaction with mobile-based services.

### **Limitations**

The use of multiple and varied outcome measures during the COVID-19 pandemic limits the reliability of judgments about the efficacy of m-health interventions in support of the elderly. Moreover, we did not have access to some other databases such as CINAHL and PsycINFO, and could not access articles published in languages other than English.

### **Conclusion**

The results of this systematic review highlight the potentials of m-health tools in the promotion of health and well-being among the older population, and show that there is a variety of usages of such tools during the current pandemic. The use of m-health keeps the elderly and healthcare providers safe, accelerates health service provision, reduces the costs of service delivery, and decreases the risk of morbidity and mortality during the time of COVID-19. Due to the high risk of COVID-19 infection in the elderly, the application of m-health tools for the health promotion of this group would be a feasible approach to address many key challenges and eventually diminish morbidity and mortality. Despite the advantages of using m-health tools for the elderly people, the results of this review showed that there are some challenges in the implementation of m-health to support the elderly who need attention from the government, healthcare providers, and society. During the COVID-19 pandemic, older people need the highest degree of support, but there appear to be barriers to accessing this support through mobile apps, which need to be resolved.

### **Acknowledgements**

None.

### **Conflicts of Interest**

None declared

### **Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### **Ethics approval and consent to participate**

Not applicable.

### **References**

1. WHO. Q&A on coronaviruses (COVID-19). Available from: <https://www.who.int/news-room/q-a-detail/q-a-coronaviruses>. 2020.
2. Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LF. The trinity of COVID-19: immunity,

inflammation and intervention. *Nature Reviews Immunology*. 2020;1-12.

3. Lipsitch M, Swerdlow DL, Finelli L. Defining the epidemiology of Covid-19—studies needed. *New England journal of medicine*. 2020;382(13):1194-6.
4. Verma J, Mishra AS. COVID-19 infection: Disease detection and mobile technology. *PeerJ*. 2020;8:e10345.
5. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet infectious diseases*. 2020.
6. Morley JE, Vellas B. COVID-19 and older adult. *J Nutr Health Aging*. 2020;24(4):364-5.
7. Smith AC, Thomas E, Snoswell CL, Haydon H, Mehrotra A, Clemensen J, et al. Telehealth for global emergencies: Implications for coronavirus disease 2019 (COVID-19). *Journal of telemedicine and telecare*. 2020;1357633X20916567.
8. Kouroubali A, Kondylakis H, Kavlentakis G, Logothetides F, Stathiakis N, Petrakis Y, et al. An eHealth platform for the holistic management of COVID-19. *Stud Health Technol Inform*. 2020;273:182-8.
9. Martel D, Lauzé M, Agnoux A, de Lacroix LF, Daoust R, Émond M, et al. Comparing the effects of a home-based exercise program using a gerontechnology to a community-based group exercise program on functional capacities in older adults after a minor injury. *Experimental gerontology*. 2018;108:41-7.
10. Trzebiński J, Cabański M, Czarnecka JZ. Reaction to the COVID-19 pandemic: the influence of meaning in life, life satisfaction, and assumptions on world orderliness and positivity. *Journal of Loss and Trauma*. 2020;1-14.
11. Lindeman D. mHealth technologies: applications to benefit older adults. Oakland, CA: Center for Technology and Aging; 2011. Ed; 2015.
12. Mirza F, Norris T. Opportunities and barriers for mobile health in New Zealand. *Studies in health technology and informatics*. 2007;129(1):102.
13. Kay M, Santos J, Takane M. mHealth: New horizons for health through mobile technologies. World Health Organization. 2011;64(7):66-71.
14. Zhang H, Cocosila M, Archer N. Factors of adoption of mobile information technology by homecare nurses: a technology acceptance model 2 approach. *CIN: Computers, Informatics, Nursing*. 2010;28(1):49-56.
15. Donaghy E, Atherton H, Hammersley V, McNeilly H, Bikker A, Robbins L, et al. Acceptability, benefits, and challenges of video consulting: a qualitative study in primary care. *British Journal of General Practice*. 2019;69(686):e586-e94.
16. Frost R, Nimmons D, Davies N. Using Remote Interventions in Promoting the Health of Frail Older Persons Following the COVID-19 Lockdown: Challenges and Solutions. *Journal of the American Medical Directors Association*. 2020;21(7):992-3.

17. Czaja SJ. Long-term care services and support systems for older adults: The role of technology. *American Psychologist*. 2016;71(4):294.
18. McMurtrey ME, Zeltmann SM, Downey JP, McGaughey RE. Seniors and technology: Results from a field study. *Journal of Computer Information Systems*. 2011;51(4):22-30.
19. Karim H, Choobineh H, Kheradbin N, Ravandi MH, Naserpor A, Safdari R. Mobile health applications for improving the sexual health outcomes among adults with chronic diseases: A systematic review. *Digital health*. 2020;6:2055207620906956.
20. Meinert E, Milne-Ives M, Surodina S, Lam C. Agile requirements engineering and software planning for a digital health platform to engage the effects of isolation caused by social distancing: case study. *JMIR Public Health and Surveillance*. 2020;6(2):e19297.
21. Jaana M, Paré G. Mobile Health Technologies Use for Self-tracking: A Survey Comparing Seniors to the General Adult Population in Canada. *JMIR Mhealth and Uhealth*. 2020.
22. Padala KP, Wilson KB, Gauss CH, Stovall JD, Padala PR. VA Video Connect for Clinical Care in Older Adults in a Rural State During the COVID-19 Pandemic: Cross-Sectional Study. *Journal of medical Internet research*. 2020;22(9):e21561.
23. van Dyck LI, Wilkins KM, Ouellet J, Ouellet GM, Conroy ML. Combating Heightened Social Isolation of Nursing Home Elders: The Telephone Outreach in the COVID-19 Outbreak Program. *The American Journal of Geriatric Psychiatry*. 2020.
24. Wang J, Katz I, Li J, Wu Q, Dai C. Mobile digital divide and older people's access to 'Internet plus social work': implications from the COVID-19 help-seeking cases. *Asia Pacific Journal of Social Work and Development*. 2020:1-9.
25. van Dijk S, Bouman R, Folmer E, den Held R, Warringa J, Marijnissen R, et al. (Vi)-rushed into online group schema-therapy based day-treatment for older adults by the COVID-19 outbreak in the Netherlands. *The American Journal of Geriatric Psychiatry*. 2020.
26. Rodenstein MS, Merchant TS, Pendergrast TR, Lindquist LA. Reducing social isolation of seniors during COVID-19 through medical student telephone contact. *Journal of the American Medical Directors Association*. 2020;21(7):948-50.
27. von Humboldt S, Mendoza-Ruvalcaba NM, Arias-Merino ED, Costa A, Cabras E, Low G, et al. Smart technology and the meaning in life of older adults during the Covid-19 public health emergency period: a cross-cultural qualitative study. *International Review of Psychiatry*. 2020:1-10.
28. Echeverría P, Bergas MAM, Puig J, Isnard M, Massot M, Vedia C, et al. COVIDApp as an innovative strategy for the management and follow-up of COVID-19 cases in long-term care facilities in Catalonia: implementation study. *JMIR public health and surveillance*. 2020;6(3):e21163.
29. McLellan LJ, Morelli M, Simeone E, Khazova M, Ibbotson SH, Eadie E. SmartPDT®: Smartphone enabled real-time dosimetry via satellite observation for daylight photodynamic therapy. *Photodiagnosis and photodynamic therapy*. 2020;31:101914.
30. Armitage R, Nellums LB. COVID-19 and the consequences of isolating the elderly. *The*

Lancet Public Health. 2020;5(5):e256.

31. Cudjoe TK, Kotwal AA. "Social Distancing" Amid a Crisis in Social Isolation and Loneliness. Journal of the American Geriatrics Society. 2020.

32. Santini ZI, Jose PE, Cornwell EY, Koyanagi A, Nielsen L, Hinrichsen C, et al. Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal mediation analysis. The Lancet Public Health. 2020;5(1):e62-e70.

33. Sacco G, Léonart S, Simon R, Noublanche F, Annweiler C, Group TS. Communication technology preferences of hospitalized and institutionalized frail older adults during COVID-19 confinement: cross-sectional survey study. JMIR mHealth and uHealth. 2020;8(9):e21845.

Figure 1: Search results and publication selection procedures




Table 1.

Table 1. Search Terms and Strategy Used in PubMed
(COVID-19[title/abstract] OR COVID19[title/abstract] OR Coronavirus [title/abstract] OR Novel coronavirus [title/abstract] OR 2019-nCoV [title/abstract] OR Wuhan coronavirus [title/abstract] OR SARS-CoV-2[title/abstract] OR SARS2[title/abstract])  AND

(Elderly [title/abstract] OR Old\* [title/abstract] OR Aged [title/abstract] OR Senior [title/abstract] OR Aging [title/abstract])

AND

(mhealth [title/abstract] OR m-health [title/abstract] OR Handheld computer [title/abstract] OR Mobile phone [title/abstract] OR Smartphone [title/abstract] OR Mobile application [title/abstract] OR Mobile technology [title/abstract] OR Mobile device [title/abstract] OR Cellular phone [title/abstract] OR Mobile app [title/abstract] OR Text message [title/abstract] OR SMS [title/abstract]).

Table 2: General characteristics and key findings of the Included Studies

Year	Study design	Country	Population of study (N)	Device or technology used	Intervention	Key finding
2020 (Paré et al.)	Case study	United Kingdom, France, and Sweden	Older users and their family members (N=6000 user)	Digital health mobile app	Design, develop, and deploy a digital health mobile app to provide a means of assisting families and peer groups in maintaining contact with older people	<ul style="list-style-type: none"> <li>• Solution with the capacity of connecting vast geographies of individuals</li> <li>• Creates a capability for virtual interconnectedness</li> <li>• Tackle a new and urgent issue arising to combat the COVID-19 worldwide pandemic</li> <li>• Connect the older adults with their family and friends during social distancing</li> <li>• To encourage positive nutrition, enhance physical activity, and promote mental</li> </ul>
2020 (Paré et al.)	Cross-sectional survey	Canada	Canadian residents/65 years and older (N=4109)	Not Applicable	To compare older adults' use of m-health technologies to that of the general	<ul style="list-style-type: none"> <li>• Considerable number of older adults reported using internet and smart technology</li> <li>• Support the m-health technologies of</li> </ul>

Year	Study design	Country	Population of study (N)	Device or technology used	Intervention	Key finding
					adult population in Canada and to investigate the factors that affect their use	<p>older people health and well-being needs in the community</p> <ul style="list-style-type: none"> <li>• Used mobile apps by users to store and monitor health-related data</li> <li>• The potential of m-health technologies for self-tracking purposes in the context of older adults</li> <li>• Necessary to culture shift in the provision of care to older adults is deemed keep up with the development of m-health technologies and the changing demographics and expectations of patients and their caregivers</li> </ul>
er et (2020)	Cross-sectional survey	United States	older veterans (N=118)	Not Applicable	To examine the capability and willingness of older veterans to participate in a VA Video Connect (VVC) visit during the COVID-19 pandemic	<ul style="list-style-type: none"> <li>• The 77% successful in the scheduled VVC visits</li> <li>• There are some lingering issues with capability and willingness to participate in video telehealth visits, particularly among older adults residing in rural areas</li> <li>• Education have a stronger influence on internet access</li> </ul> <p>- Factor affected the use VA Video Connect (VVC) by older:</p> <ul style="list-style-type: none"> <li>• Availability of internet access (77%)</li> <li>• Availability of email access (70%)</li> <li>• Availability of a device with a camera (56%)</li> <li>• Capable of participating in VVC appointments (56%)</li> <li>• Willingness to participate in VVC appointments (58%)</li> <li>• Capable and willing to participate in VVC appointments (53%)</li> </ul> <p>- Suggestions for more access to VVC appointments:</p> <ul style="list-style-type: none"> <li>• Necessity of policy efforts to bring broadband internet access to rural residents</li> <li>• Addressing the digital divide in rural area</li> <li>• Needed of targeted educational efforts to train those residing in rural areas</li> <li>• Need to distribution of video-capable devices be augmented by the provision of high-speed internet and age-friendly</li> </ul>



Year	Study design	Country	Population of study (N)	Device or technology used	Intervention	Key finding
						technical assistance
et (0)	Observational study	United States	Nursing home residents (N=30)	Telephone Outreach in the COVID-19 Outbreak (TOCO) Program	To address social isolation in nursing home residents, the Student Interest Group initiated a Telephone Outreach in the COVID-19 Outbreak (TOCO) Program that implements weekly phone calls with student volunteers.	<ul style="list-style-type: none"> <li>• Positive effect of TOCO Program among seniors</li> <li>• Benefited the seniors from meaningful conversations with their volunteer companions</li> <li>• The nursing home residents shared stories of their lives and their families with student volunteers</li> <li>• Identify the unique and actionable needs of the senior companions by volunteers</li> <li>• Recognized common challenges experienced by many of these older adults (e.g. feelings of restlessness and anxiety as isolation continues, and fearfulness as COVID-19 enters nursing home facilities)</li> <li>• Exacerbated challenges for the elderly's due to lack of access to technology (e.g. limited computer and internet access to communicate with loved ones)</li> <li>• Alleviate social isolation experienced by nursing home residents during the COVID-19 pandemic</li> </ul>
g et (0)	Qualitative study	China	Elderly people (N=8)	Not Applicable	How has the COVID-19 pandemic highlighted the age-based mobile digital divide in China, and what are the implications for gerontological internet-based social work?	<ul style="list-style-type: none"> <li>• Replacing the mobile internet instead the PC to become the dominating online platform in China</li> <li>• Need of younger (e.g. their granddaughter) to support older people to seek help through internet-based</li> <li>• Used the mobile internet to seek help, purchase masks, or show the Green Code for travelling by older people</li> <li>• Need to simple apps for many older people</li> <li>• Need to used mobile phone or tablet with a big screen for elderly</li> <li>• Expensively of tablet with the call function for bought by elderly</li> <li>• Internet-based ICT used by social workers should be better designed and accessible to older people in terms of visibility, price, and operation</li> <li>• Urgent need to make more efforts to provide internet-based social work services</li> <li>• Need to be better trained for social workers to deliver services tailored to the specific needs of elderly population'</li> </ul>

Author year	Study design	Country	Population of study (N)	Device or technology used	Intervention	Key finding
						<ul style="list-style-type: none"> <li>Assess the capabilities of each elderly client by social workers to use different forms of internet based methods before they provide services to that client</li> </ul>
Dijk al. (2020)	Case study	Netherlands	Elderly patient aged between 64 and 70 years (n=4)	Online psychotherapy day-treatment program	To present first impressions on feasibility of transforming a psychotherapy day-treatment program for older persons into an online program	<ul style="list-style-type: none"> <li>Comfort in setting up the individual treatment plan by video conferencing</li> <li>Increasing the motivation of therapists for positive change</li> <li>Existence of time for therapists to prepare the psychotherapy sessions</li> <li>Online and therapy of older ran smoothly</li> <li>Providing active movements (gymnastics), relaxation, and mindful exercises (e.g., Tai Chi) to the older patients</li> </ul>
Re et al. (2020)	Observational study	United States	Older adults at risk for increased social isolation during COVID-19 (N=-)	Phone call outreach program (Seniors Overcoming Social Isolation (SOS))	Created a phone call outreach program in which medical and health professions student volunteers called older adults, living in long-term-care facilities (LTCFs) and the community, at risk of social isolation during COVID-19	<ul style="list-style-type: none"> <li>Creating happy and enjoyment among older patients to have someone to talk</li> <li>Interrupting social isolation of older adults and residents of independent</li> <li>Improving the lives of older adults during the COVID-19 pandemic</li> <li>Understanding the needs of older adults in the community</li> </ul>
Abold al. (2020)	Qualitative study	Italian, Mexican, Portuguese and Spanish	Older participants aged 65-87 years (N= 351)	Not Applicable	To fill a knowledge gap about older adult perceptions of how smart technology influenced their meaning in life during the Covid-19	<ul style="list-style-type: none"> <li>Used of smart technology to connect with family, colleagues and friends and create meaningful relations)</li> <li>Important instrument for carrying out older individual activities</li> <li>Helped to develop a sense of spirituality</li> <li>Health and safety-related support of older</li> <li>Allowed to self-growth and reflection about how to achieve self-growth during the Covid-19</li> <li>To continue physical activity</li> </ul>
Over et al.	Observational	Spain	Older residents and individuals	COVID App	To report the implementation of	<ul style="list-style-type: none"> <li>Helped to institutional staff from long-term care facilities and primary care</li> </ul>

Year	Study design	Country	Population of study (N)	Device or technology used	Intervention	Key finding
2020	Study		with physical and mental disabilities in long-term care facilities (N=)	mobile tool	COVIDApp innovative tool for the management of long-term care facility residents as a high-risk population, specifically for early identification and self-isolation of suspected cases, remote monitoring of mild cases, and real-time monitoring of the progression of the infection.	<p>clinicians address the COVID-19 pandemic</p> <ul style="list-style-type: none"> <li>• Providing a new channel for real-time communication</li> <li>• Early detection of suspected COVID-19 cases</li> <li>• Rapid development of epidemiological actions (e.g. self-isolation of suspected cases and contacts and relocation after positive or negative results)</li> <li>• Remote management of mild COVID-19 cases treated in institutions</li> <li>• Knowledge of progression of the infection in long-term care facilities</li> <li>• Advise and support older people in nursing homes through a website that enables direct contact between a senior geriatrician and centers for older people</li> <li>• Monitor the progression of the infection</li> <li>• Decrease in the number of deaths and confirmed cases of COVID-19</li> <li>• Enabled to intervene proactively by isolating residents with suspected infection early and by monitoring contacts</li> <li>• Reduce transmission in a vulnerable population</li> </ul>
ella et al. 2020	Observational study	United Kingdom	Elderly patient (N=)	Smartphone-based application and web-portal (smartPDT®)	Determine the accuracy of the algorithms used in the commercial SmartPDT® application	<ul style="list-style-type: none"> <li>• Provide treatment for the common pre-malignant condition of chronic photo damaged skin actinic keratosis</li> <li>• Creating a more convenient and reliable therapy</li> <li>• Reducing risk in an elderly patient population during the Covid-19 pandemic</li> </ul>

## Supplementary Files