

A Cross-Country Comparative Study on the Role of Information and Communication Technology Policy and Infrastructure to Curb the Spread of Novel Coronavirus

Nam Ji Eum, Seung Hyun Kim

Submitted to: JMIR Public Health and Surveillance on: June 08, 2021

Disclaimer: © **The authors. All rights reserved.** This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on it's website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressively prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript	. 5
Supplementary Files	
Multimedia Appendixes	26
Multimedia Appendix 1	

A Cross-Country Comparative Study on the Role of Information and Communication Technology Policy and Infrastructure to Curb the Spread of Novel Coronavirus

Nam Ji Eum¹ BA; Seung Hyun Kim¹ BA, MS, PhD

¹Yonsei University Seoul KR

Corresponding Author:

Seung Hyun Kim BA, MS, PhD Yonsei University 50 Yonsei-ro, Sinchon-dong, Seodaemun-gu Seoul KR

Abstract

Background: Despite a worldwide effort, control of COVID-19 transmission and of its aftereffect is lagging. As can be seen from the case of SARS-CoV-2 and influenza, such worldwide crisis and its side effects are likely to recur in the future because of extensive international interactions. Consequently, there is an urgent need to find the determinants that can mitigate the disease diffusion. Meanwhile, we observed that the pace of diffusion and its consequences varied substantially across countries, signaling the need for country-level investigation.

Objective: We conducted research on how (1) distancing-enabling ICT infrastructure and (2) medical ICT infrastructures and related policies have affected the cumulative confirmed cases, fatality rates, and initial diffusion speed across different countries. We analyze the determinants of COVID-19 diffusion during the relatively early days of the pandemic.

Methods: Based on our data for country-level characteristics including demographics, culture, ICT infrastructure, policies, economic status, and diffusion of COVID-19, we conducted regression to analyze the dataset. To gain further insight, we conducted a subsample analysis for countries with low population density.

Results: Contrary to expectations, our analysis indicated that the ICT infrastructure is mostly ineffective in reducing the COVID-19 transmission and fatality rate. Instead, our full sample analysis showed that implied telehealth policy is associated with lower fatality rates when controlled for cultural characteristics (p-value =.004). In particular, the fatality rate for countries with an implied telehealth policy was lower than that for others by 2.7 percentage point. Interestingly, explicit telehealth policy was found to be not effective in curbing the fatality rates (p-value =.299). Furthermore, countries with government health website had 36% fewer confirmed cases than those without it, when controlled for cultural characteristics (p-value=.03). To investigate their heterogeneous effects across countries, 1) interaction effect of both implied and explicit telehealth policy with other medical ICT infrastructure variables and 2) subsample regression for countries with lower population density were conducted. Our analysis further revealed that the interaction between implied telehealth policy and training ICT health was significant (p<0.01), suggesting that implied telehealth policy may be more effective when in-service training on ICT is provided to health professionals. In addition, a rate of credit card ownership, as an enabler of convenient e-commerce transactions and distancing, showed a negative associate with fatality rates only in full sample analysis (p-value =.04), but not in subsample analysis (p-value =.76), highlighting that distancing-enabling ICT is more useful in densely populated countries.

Conclusions: Our findings demonstrate important relationships between national traits and COVID-19 infections, suggesting guidelines for policymakers to minimize the negative consequences of pandemics. Especially, the findings suggest (1) physician's autonomous use of medical ICT and (2) strategic allocation of distancing-enabling ICT infrastructure in high population density countries to maximize efficiency. This study also urges further research to investigate the role of health policies in combatting COVID-19 and other pandemics.

(JMIR Preprints 08/06/2021:31066)

DOI: https://doi.org/10.2196/preprints.31066

Preprint Settings

- 1) Would you like to publish your submitted manuscript as preprint?
- **✓** Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users. Only make the preprint title and abstract visible.

- No, I do not wish to publish my submitted manuscript as a preprint.
- 2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?
- ✓ Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain ves, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <a href="https://example.com/above

Original Manuscript

Original Paper

A Cross-Country Comparative Study on the Role of Information and Communication Technology Policy and Infrastructure to Curb the Spread of Novel Coronavirus

Abstract

Background: Despite worldwide efforts, control of COVID-19 transmission and of its aftereffect is lagging. As seen from the case of SARS-CoV-2 and influenza, such worldwide crisis and its side effects are likely to recur in the future because of extensive international interactions. Consequently, there is an urgent need to find the determinants that can mitigate the disease spread. Meanwhile, we observed that the transmission speed and severity of consequences varied substantially across countries, signaling the need for country-level investigation.

Objective: We aimed to investigate how distancing-enabling information and communication technology (ICT) infrastructure and medical ICT infrastructures and related policies have affected the cumulative confirmed cases, fatality rates, and initial speed of transmission across different countries.

Methods: We analyzed the determinants of COVID-19 transmission during the relatively early days of the pandemic by conducting regression analysis based on our data for country-level characteristics including their demographics, culture, ICT infrastructure, policies, economic status, and transmission of COVID-19. To gain further insights, we conducted a subsample analysis for countries with low population density.

Results: Our full sample analysis showed that *implied telehealth policy*, which refers to the lack of specific telehealth related policy but is referred to in general eHealth policy, was associated with lower fatality rates when controlled for cultural characteristics (*p*-value =.004). Particularly, the fatality rate for countries with an implied telehealth policy was lower than that for others by 2.7 percent. Interestingly, *stated telehealth policy*, which refers to existence of specified telehealth policy, was found not to be associated with lower fatality rates (*p*-value =.299). Furthermore, countries with government-run health website had 36% fewer confirmed cases than those without it, when controlled for cultural characteristics (*p*-value=.03). Our analysis further revealed that the interaction between implied telehealth policy and training ICT health was significant (*p*-value<0.01), suggesting that implied telehealth policy may be more effective when in-service training on ICT is provided to health professionals. In addition, a rate of credit card ownership, as an enabler of convenient e-commerce transactions and distancing, showed a negative association with fatality rates in full sample analysis (*p*-value =.04), but not in subsample analysis (*p*-value =.76), highlighting that distancing-enabling ICT is more useful in densely populated countries.

Conclusions: Our findings demonstrate important relationships between national traits and COVID-19 infections, suggesting guidelines for policymakers to minimize the negative consequences of pandemics. The findings suggest physicians' autonomous use of medical ICT and strategic allocation of distancing-enabling ICT infrastructure in countries with high population density to maximize efficiency. This study also encourages further research to investigate the role of health policies in combatting COVID-19 and other pandemics.

Keywords: health policy; telehealth; physical distancing; disease transmission; COVID-19

Introduction

First identified in December 2019, the novel coronavirus disease (COVID-19) outbreak has rapidly spread worldwide. As of September 23, 2021, the World Health Organization reported that over 229.8 million people were infected worldwide and with over 4.7 million deaths caused by COVID-19 [1]. Furthermore, on December 19, 2020, a mutant of COVID-19, labeled B.1.1.7, was found, causing further havoc specifically in European countries. Consequently, U.K. Prime Minister Boris Johnson imposed another lockdown, this one even stricter than the previous lockdowns [2]. This kind of catastrophic pandemic is not unprecedented as from 1918 to 1919; the H1N1 influenza A virus was first manifested, and went on to infect approximately 40% of the global population with a mortality rate of more than 2% [3]. The H1N1 flu persists to this date, over 100 years from its first appearance, and has undergone significant genetic mutations. Virologists expect COVID-19 to follow a similar pattern [4,5]. As such, the prevalence of pandemics and genetic mutations is not a one-off phenomenon. Other outbreaks with disruptive social and economic consequences are probable [6], demanding research on how to control the spread of a virus in its early stages.

With this urgent need in mind, it is noteworthy that the infection and fatality rates as well as speed of transmission varied widely across countries. Countries such as Israel, Singapore, South Korea, and Taiwan are regarded as relatively successful in curbing the transmission [7], whereas European countries and the United States experienced an explosive increase in the number of confirmed cases [1]. It has been known that minimizing physical contact between individuals without disturbing their daily lives and improved medical practices are crucial in managing the spread of infectious diseases in general [8]. First, as a means of reducing physical contacts and enabling social distancing, national information and communication technology (ICT) infrastructure such as e-commerce and high speed Internet connection have played a key role in many countries [25-26]. Second, the possible importance of medical ICT policies and infrastructures has also been recognized. For example, effective use of telehealth practices has been credited with the successful management of other infectious diseases such as severe acute respiratory syndrome (SARS) or Middle East respiratory syndrome (MERS), pointing to a possible role for telehealth in controlling pandemics [12]. In addition, other medical ICTs, such as computerized physician order entry (CPOE) and e-prescribing, have been acknowledged as drivers of healthcare improvements in quality and efficiency [13]. Third, countries differed significantly across other dimensions, such as implementation of early lock-down and conformity to government policies because of their cultural differences, that could have affected their success in social distancing.

Therefore, to gain a broader insight, there is a need for country-level analysis of the national-level characteristics that mitigate the spread of COVID-19 through successful distancing and improved medical practices. Nevertheless, previous studies on the spread of COVID-19 tended to have narrower scopes such as individuals, several cities, a single country, or a single continent rather than a global focus [16-18]. Although there are some exceptions, those studies were still limited in number of countries, possibly due to difficulty of data collection, for investigations [19,20]. For instance, the latest cross-country study on the effect of threat or coping appraisal on distancing compliance was conducted by comparing three countries only [19]. In addition, prior research also focused on how demographic, cultural, or political factors of a country affected COVID-19 infection [21-24], but did not include the ICT related factors.

To address such research gap, the primary objective of this study is to identify what national characteristics have a major impact on curtailing contagious diseases during the relatively early days of the pandemic. In particular, we focus on the role of distancing-enabling ICT infrastructure (DistancingICT, hereafter) and medical ICT infrastructure and policy (MedicalICT, hereafter) in

containing COVID-19 infections, fatality rates and transmission speed.

Methods

Data Collection

The main sources of country-level data used in this study include the UN, the World Bank, WHO, Worldometers, OWID, Ookla, Hofstede Insights, and Wikipedia. Among these, Worldometers is a reference site that provides real-time statistics on diverse topics. As a widely used source of research, media and teaching, OWID is an online scientific publication institute that focuses on global issues such as poverty and disease. Ookla provides analyses of internet access performance metrics. Hofstede Insights provides culture scores of each country based on Hofstede's cultural dimension theory [25], and is widely used in academic research. For example, to predict growth of COVID-19 confirmed cases across several countries, Hofstede dimensions are used to account for cultural factor [26]. Wikipedia is used only to determine which countries enforced national lockdowns in the early days of COVID-19. The accuracy of enforcement and the dates of enforcement are further confirmed through research in the media. All the data were collected between July and August 2020. This sampling strategy allows us to analyze the determinants of COVID-19 transmission during the early days of the pandemic.




Figure 1. Map chart of countries used in analysis

We limit our analysis to the countries that reported statistics related to the spread of COVID-19. For example, countries without accurate statistics on deaths as of July 28, 2020, are excluded. China, the country at the epicenter of COVID-19, is excluded because the patterns of disease spread and governmental control differ greatly from other countries. By matching the data for the social, economic, and demographic statuses of countries as well as their physical distancing and healthcare-related ICT infrastructure, our final dataset consists of 98 countries. The countries included in our analysis are shown in (Figure 1).

We focus on three dependent variables that represented the early state of spread of COVID-19 in each country: (a) the cumulative number of infections, (b) fatality rate, (c) number of days from initial infection to 1,000th infection. The first dependent variable represents the cumulative number of COVID-19 infections per country as of July 28, 2020. The second dependent variable is the fatality rate, which is the death toll over the cumulative number of infections. The third dependent variable is the transmission speed of COVID-19 in its initial stage. This is represented for each country by the number of days from the date of the first confirmed case to the date of the 1,000th confirmed case. For the third dependent variable, countries with fewer than 1,000 cases as of July 28, 2020, are excluded due to the difficulty of calculating the speed of transmission.

The two main categories of our main independent variables are DistancingICT, and MedicalICT. For DistancingICT, we chose two variables based on findings of previous research: rate of credit card ownership and broadband internet speed. Research and Markets reported that during the COVID-19 pandemic, North America's online sales surged, and credit cards were the top payment method for these online sales [23]. Previous studies further support the relationship between e-commerce and credit card usage. Meyll and Walter (2019) surveyed more than 25,000 U.S households, confirming that individuals using mobile payments are likely to use credit cards [24]. Given this direct relationship between e-commerce and credit card usage, we identified credit cards as a major enabler of distancing through their serving as alternatives to offline shopping. Broadband internet speed also assists individuals to comply with stay-at-home orders during COVID-19 [25-26]. Next, the MedicalICT variables involve the availability of a national telehealth policy or strategy; availability of government-supported multilingual health internet websites that provide information; institutions with healthcare ICT training; and national electronic health records (EHR). These variables represent MedicalICT because the variables relate to either reliable online sharing of medical information (government health internet sites and national EHR) or the effective use of existing medical ICT technology (national telehealth policy and healthcare ICT training).

Particularly, national telehealth policy is distinguished into two types: implied and stated. The *implied telehealth policy* means that a country does not have a specific national telehealth policy or strategy; such a policy is referred to in the overall national eHealth policy. Australia, Finland, and the U.S. are examples of such countries. The U.K. and Norway, on the other hand, have separate telehealth policies. For instance, the Norwegian Ministry of Health commissioned the Norwegian Centre for Telemedicine to foster telehealth services, while assuring "the necessary actions to secure a successful dissemination of the services" [27]. As such, countries with a specific national telehealth policy or strategy — apart from national eHealth strategy — are accounted for as "stated" telehealth policy in our model. Although the term telehealth and eHealth are at times used interchangeably, they differ in purpose of use. While telehealth indicates usage of ICT to promote long-distance care, eHealth indicates usage of ICT for health in general [28]. For example, in the U.S. before COVID-19, telehealth was only for people who need long-distance care due to limited access to nearby

hospitals [29,30], while eHealth was widely applied to patients regardless of hospital accessibility. Accordingly, reimbursement on telehealth has not been prioritized nor sufficiently instituted, as compared to that on general eHealth [30]. As such, telehealth, comparably, lacked clearly stated guidance before the pandemic.

Our control variables were selected based on the results of prior research that primarily focused on how demographic, cultural, or political factors of a country affected COVID-19 infections [21-24]. The larger the scale of these countries' economies, the greater their potential for economic activities such as job hunting and international exchanges that increase the opportunities for infections. Thus, we included gross domestic products at purchasing power parity (GDP PPP) and unemployment rates as controls for the economic statuses of countries. Moreover, because several studies have indicated that high temperatures and humidity may influence the infection rate of COVID-19, we included annual rainfall and temperature as controls [31]. Similarly, we added other controls such as the proportion of senior citizens, early implementation of a national lockdown, and population density to our model, along with two culture-related variables, individualism and uncertainty avoidance. Overall, we selected additional control variables that are identified as important determinants of the spread of contagious viruses in the literature. Detailed explanations of our main variables and of additional control variables are summarized in (Table 1).

Table 1. Variable Description

Variable name	Description (by Country)	Source	Year Measured
Dependent Variables			
Total Cases ^a	Number of individuals infected by COVID-19	Worldometers	2020
Fatality Rate ^a	(0-100)	Worldometers	2020
Number of Days ^b	Number of days elapsed from the first confirmed case to the 1000th confirmed case	OurWorldinData	2020
Control Variable			
GDP PPP	GDP by purchasing power parity in billions	World Bank	2019
Unemployment Rate	Unemployment rate (0-100)	World Bank	2019
Population Density	People per sq. km of land area	World Bank	2019
Percent 60 or Over	Percentage of people ages 60 or older	United Nations	2019
Annual Rainfall	The average annual rainfall in mm	World Bank	2019
Annual Temperature		World Bank	2019
Early Lockdown	Implementation of a national lockdown within one month of the first confirmed case (dichotomous)	Wikipedia, Press	2020
Individualism	Cultural dimension score for preference for a loosely knit social framework in which individuals are expected to take care of only themselves and their immediate families $(0-100)$	Hofstede	2015
Uncertainty Avoidance	Cultural dimension score for degree to which the members of a society feel uncomfortable with uncertainty and ambiguity $(0-100)$	Hofstede	2015
ICT Infrastructure En	abling Physical Distancing		
Credit Card Ownership	The percentage of respondents who report having a credit card (Ages 15+)	World Bank	2017

Broadband Speed	Broadband internet speed in Mbps	Ookla	2019
Medical ICT Infrastru	cture and Policy		
• • • • • • • • • • • • • • • • • • • •	Country with stated telehealth policy or strategy (1: Yes, 0: Otherwise)	WHO	2015
	Country with no specific telehealth policy or strategy but is referred in overall eHealth policy or strategy (1: Yes, 0: Otherwise)	WHO	2015
Govt Health websites	Government-supported health internet sites provide information (1: Available, 0: Not Available)	WHO	2015
Training ICT Health	Institutions offering in-service training to health professionals on ICT for health (1: Available, 0: Not Available)	WHO	2015
National EHR	Country with national Electronic Health Records (EHR) (1: Available, 0: Not Available)	WHO	2015

Empirical Analysis

For each dependent variable, we specified our models as follows:

$$\log \left(\text{Total Case } \mathbf{s}_{i} \right) = \alpha_{1} + \beta_{11} \text{Contro } \mathbf{l}_{i} + \beta_{12} \text{DistancingICT}_{i} + \beta_{13} \text{MedicalICT}_{i} + \epsilon_{1i}$$
 (1)

Fatality Rate_i =
$$\alpha_2 + \beta_{21} \text{Control}_i + \beta_{22} \text{DistancingICT}_i + \beta_{23} \text{MedicalICT}_i + \epsilon_{2i}$$
 (2)

$$log (Number of Days_i) = \alpha_3 + \beta_{31} Control_i + \beta_{32} Distancing ICT_i + \beta_{33} Medical ICT_i + \epsilon_{3i}$$
 (3)

where i denotes an individual country. For simplicity, an OLS estimator was used to estimate the coefficients. The variables of main interest are DistancingICT and MedicalICT. The positive coefficient values of DistancingICT and MedicalICT in Equations (1) and (2) demonstrate that the variables increase the total number of confirmed cases and the fatality rate. In contrast, the positive coefficients of DistancingICT and MedicalICT in Equation (3) represent a slower transmission speed. In all three models, we used the same set of control variables, including GDP PPP, unemployment rate, population density, elderly population ratio, annual rainfall, annual temperature, and early lockdown. For normality, we log transformed all the variables, including Total Cases and Number of Days, that displayed skewed distributions and were nonnegative.

For each dependent variable, our baseline model includes all the main independent variables and controls—but without the two culture-related variables of individualism and uncertainty avoidance. In the second model, we added these culture-related variables to the baseline model. We performed this separate estimation because the content for the culture-related variables is not available for all 98 countries. Thus, adding them to the model reduces the sample size from 98 to 69. In the third model, we added several interaction terms to check for possible interaction effects between MedicalICT variables. To ensure that independent variables in the analysis were not correlated, we calculated variance inflation factor (VIF). All the independent variables had VIF values less than 10, which indicates no multicollinearity violations [32]. Lastly, robust standard errors were used to address any possible heteroskedastic error.

We also conducted a subsample analysis in which we removed countries with high population

Definitions for Hofstede variables are obtained from https://hi.hofstede-insights.com/national-culture

^a Data collected as of July 28th, 2020

^b Data collected as of August 9th, 2020

density. A prior study showed positive correlation between population density and COVID-19 infection [33]. Residents of densely populated countries inevitably incur interactions and contacts with more people offline, whereas loosely populated countries can reduce such possibilities when the infrastructure is built up. Moreover, less populated countries may not have sufficient localized medical services, thus requiring more medical ICT infrastructure than other countries. The needs and utilization of ICT would significantly vary by population density as well. Thus, to determine if our results might be biased by the inclusion of countries with higher populations densities, we conducted an additional analysis with only countries with lower population density.

Results

Main Analysis

The descriptive statistics for the variables used in this analysis and their correlations are presented in (Table 2) and (Table S1 in Multimedia Appendix 1), respectively.

Table 2. Summary Statistics

Tubic 2. Julilliary Julistics					
Variable Names	No. of countries	Mean	SD	Min	Max
Dependent Variables					
Total Cases	98	113,021.2	464,974.3	20	4,498,343
Fatality Rate	95	3.32	3.177	0.053	15.26
Number of Days	91	51.02	29.3	11	139
Control Variables					
GDP PPP	98	734.76	2,290.02	9.03	21,427.7
Unemployment Rate	98	6.55	4.68	0.09	28.18
Population Density	98	256.50	919.32	2.11	8,737.02
Percent Ages 60 or Older	98	15.32	8.85	3.19	34.02
Annual Rainfall	98	78.85	54.56	1.53	244.87
Annual Temperature	98	16.74	8.57	-4.97	29.29
Early Lockdown	98	0.36	0.48	0	1
Individualism	68	42.85	23.57	6	91
Uncertainty Avoidance	68	66.97	22.67	8	100
Distancing Enabling ICT Infra	structure				
Credit Card Ownership Rate	98	21.23	22.25	0	83
Broadband Speed	98	45.47	36.40	4.18	191.93
Medical ICT Infrastructure					
Telehealth Policy (Stated)	98	0.22	0.42	0	1
Telehealth Policy (Implied)	98	0.37	0.49	0	1
Govt Health Websites	98	0.61	0.49	0	1
Training ICT health	98	0.82	0.39	0	1
National EHR	98	0.47	0.5	0	1

The results for regressions with robust standard error are shown in (Tables 3-5). For each dependent variable, number of confirmed cases (Table 3), fatality rate (Table 4), and transmission speed (Table 5), Model (1) is the baseline model while Model (2) adds the culture-related variables as controls. Model (3) includes all the control variables as well as the interaction effects between distancing ICT and medical ICT variables (DistancingICT $_i$ × MedicalICT $_i$). As mentioned above, in the case of Number of Days $_i$, countries with no reported cases of the 1,000th infection as of July 28, 2020, were excluded from the analysis. As for the goodness of fit, R^2 for Total Cases $_i$, is higher than that of

Fatality Rate $_i$ and of Number of Days $_i$, indicating that the national characteristic variable used in the analysis explains the cumulative number of infected better than the two other dependent variables. For Total Cases $_i$, the R^2 values for the three models are 61.5%, 73.3%, and 75.4%, respectively. Considering that the R^2 averages of Fatality Rate $_i$ and Number of Days $_i$ are 44.5% and 37%, respectively, the overall explanatory power of the models for Total Cases $_i$, exceeds that of the two others. Therefore, national characteristics account for a significant portion of the differences in cumulative number of confirmed cases by country.

Table 3. Main regression results for the full sample: log (Total Cases)

Variables	DV ^a : log (To	tal Cases)				
	(1)		(2)		(3)	
	Baseline	P value	Including	P value	Interaction	P value
	Model		Culture		Effects	
log (GDP PPP)	1.136	<.001	1.124	< 0.001	1.104	<0.001
	(0.116) ^b		(0.122)		(0.133)	
Unemployment Rate	0.033	0.04	0.015	0.40	0.019	0.32
	(0.016)		(0.018)		(0.019)	
og (Population Density)	0.250	0.06	0.278	0.07	0.359	0.03
	(0.129)		(0.15)		(0.162)	
Percent Ages 60 or Older	-0.027	0.06	-0.054	0.006	-0.055	0.007
	(0.014)		(0.019)		(0.02)	
og (Annual Rainfall)	-0.253	0.12	-0.023	0.93	-0.115	0.66
	(0.169)		(0.246)		(0.258)	
Annual Temperature	-0.021	0.07	-0.030	0.06	-0.035	0.03
	(0.012)		(0.015)		(0.016)	
Early Lockdown	-0.018	0.91	0.04	0.80	-0.06	0.74
-	(0.151)		(0.158)		(0.179)	
ndividualism			0.005	0.42	0.005	0.37
			(0.006)		(0.006)	
Uncertainty Avoidance			0.007	0.08	0.006	0.15
			(0.004)		(0.004)	
Distancing Enabling ICT In	frastructure					
Credit Card Ownership Rate	-0.002	0.64	0.0002	0.97	-0.0002	0.97
	(0.005)		(0.005)		(0.005)	
log (Broadband Speed)	0.068	0.83	0.156	0.64	0.104	0.78
3 ((0.316)		(0.335)		(0.364)	
Medical ICT Infrastructure	()		()		()	
Telehealth Policy (Stated)	0.055	0.76	0.192	0.32	0.861	0.19
resemble of (States)	(0.179)		(0.191)	0.52	(0.648)	0,10
Telehealth Policy (Implied)	-0.003	0.98	-0.098	0.56	-0.13	0.77
Teleneurin Foliey (Implieu)	(0.152)	0.50	(0.166)	0.50	(0.433)	0.77
Govt Health Websites	-0.221	0.17	-0.440	0.03	-0.675	0.02
Government vectories	(0.161)	0.17	(0.193)	0.05	(0.278)	0.02
Training ICT Health	0.174	0.35	0.096	0.64	0.276	0.38
Truming 101 Treatm	(0.183)	0.55	(0.205)	0.04	(0.311)	0.50
National EHR	0.179	0.20	0.094	0.52	0.206	0.38
Ivational Effic	(0.137)	0.20	(0.144)	0.52	(0.231)	0.50
Interaction	(0.15/)	<u> </u>	(0.144)		(0.231)	
Telehealth Policy (Stated)					0.040	0.01
*Govt Health Websites					-0.049	0.91
					(0.443)	0.14
Telehealth Policy (Implied)					0.527	0.14
*Govt Health Websites					(0.351)	0.46
Telehealth Policy (Stated)					-0.567	0.46
*Training ICT Health					(0.758)	

Telehealth Policy (Implied)					-0.241	0.57
*Training ICT Health					(0.417)	
Telehealth Policy (Stated)					-0.117	0.79
*National EHR					(0.442)	
Telehealth Policy (Implied)					-0.25	0.52
*National EHR					(0.384)	
Constant	-8.285	< 0.001	-8.577	< 0.001	-8.140	< 0.001
	(1.347)		(1.579)		(1.745)	
Observations, n	98		69		69	
\mathbb{R}^2	0.615		0.733		0.754	
Adjusted R ²	0.	55	0.65		0.636	

Table 4. Main regression results for the full sample: Fatality Rate

Variables	DV ^a : Fatality	Rate				
	(1) Baseline Model	P value	(2) Including Culture	P value	(3) Interaction Effects	P value
log (GDP PPP)	1.635	0.005	1.573	0.02	1.993	0.007
	(0.571) ^b		(0.67)		(0.711)	
Jnemployment Rate	-0.042	0.59	-0.019	0.84	0.043	0.67
	(0.077)		(0.095)		(0.099)	
log (Population Density)	0.036	0.95	1.579	0.06	1.962	0.03
	(0.623)		(0.805)		(0.843)	
Percent Ages 60 or Older	0.143	0.04	0.045	0.66	0.025	0.81
	(0.07)		(0.101)		(0.103)	
log (Annual Rainfall)	0.198	0.81	2.373	0.08	2.224	0.11
	(0.835)		(1.325)		(1.346)	
Annual Temperature	-0.011	0.85	-0.033	0.69	-0.041	0.62
	(0.058)		(0.082)		(0.082)	
Early Lockdown	-0.617	0.41	-1.000	0.25	-0.535	0.57
	(0.737)		(0.86)		(0.931)	
Individualism			0.138	< 0.001	0.141	< 0.001
	/		(0.031)		(0.031)	
Uncertainty Avoidance			0.034	0.13	0.036	0.11
			(0.022)		(0.022)	
Distancing Enabling ICT In	frastructure					
Credit Card Ownership Rate	0.0002	0.99	-0.057	0.04	-0.074	0.01
•	(0.023)		(0.027)		(0.028)	
log (Broadband Speed)	-1.287	0.40	-1.199	0.51	-0.015	0.99
	(1.513)	7	(1.799)		(1.893)	
Medical ICT Infrastructure			(=1.55)		(=:000)	
Telehealth Policy (Stated)	-0.715	0.42	-1.093	0.30	-1.03	0.76
resemble of the control of the contr	(0.888)		(1.042)	0.00	(3.367)	01.0
Telehealth Policy (Implied)	-1.176	0.12	-2.684	0.004	-6.231	0.008
(implied)	(0.743)	V.1 L	(0.903)	2.30 .	(2.242)	2,000
Govt Health Websites	-0.053	0.95	-1.721	0.11	-2.985	0.05
	(0.8)	2.00	(1.058)		(1.473)	3,03
Training ICT Health	-0.37	0.68	-0.091	0.93	0.029	0.99
	(0.904)	2.00	(1.1)	3.55	(1.611)	3,00
National EHR	-0.508	0.44	0.888	0.26	-0.206	0.87
	(0.655)		(0.78)	J.20	(1.208)	3.07
Interaction	(0.000)		(0., 0)		(1,200)	

 ^a DV: Dependent Variable
 ^b Standard errors in parentheses
 ^c N/A: Not applicable

Telehealth Policy (Stated)					2.896	0.22
*Govt Health Websites					(2.315)	
Telehealth Policy (Implied)					1.7	0.36
*Govt Health Websites					(1.832)	
Telehealth Policy (Stated)					-0.595	0.88
*Training ICT Health					(3.934)	
Telehealth Policy (Implied)					1.654	0.45
*Training ICT Health					(2.162)	
Telehealth Policy (Stated)					-1.256	0.59
*National EHR					(2.293)	
Telehealth Policy (Implied)					3.419	0.096
*National EHR					(2.009)	
Constant	-13.999	0.04	-24.255	0.007	-30.459	0.002
	(6.791)		(8.6)		(9.241)	
Observations, n	95	95		68		8
\mathbb{R}^2	0.2	0.271		0.496		669
Adjusted R ²	0.1	43	0.338		0.358	

Table 5. Main regression results for the full sample: log (Number of Days)

Variables	DV ^a : log (Nu	ımber of Day	vs)			
	(1)		(2)	5.0	(3)	
	Baseline	P value	Including	P value	Interaction	P value
	Model		Culture		Effects	
log (GDP PPP)	-0.130	0.005	-0.08	0.15	-0.061	0.29
	$(0.044)^{b}$		(0.055)		(0.058)	
Unemployment Rate	-0.001	0.84	-0.001	0.87	0.004	0.61
	(0.006)		(0.007)		(800.0)	
og (Population Density)	-0.068	0.16	-0.089	0.17	-0.079	0.24
	(0.047)		(0.064)		(0.066)	
Percent Ages 60 or Older	0.005	0.30	0.01	0.23	0.003	0.71
	(0.005)		(800.0)		(0.008)	
log (Annual Rainfall)	0.032	0.62	-0.05	0.63	-0.057	0.59
	(0.063)		(0.104)		(0.105)	
Annual Temperature	0.010	0.03	0.008	0.21	0.006	0.36
	(0.005)		(0.007)		(0.007)	
Early Lockdown	-0.096	0.09	-0.118	0.09	-0.118	0.13
	(0.055)		(0.069)		(0.075)	
Individualism			-0.002	0.46	-0.002	0.46
			(0.002)		(0.002)	
Uncertainty Avoidance			-0.003	0.17	-0.002	0.26
•			(0.002)		(0.002)	
Distancing Enabling ICT In	frastructure					
Credit Card Ownership Rate	-0.0003	0.88	-0.001	0.56	0.0004	0.88
•	(0.002)		(0.002)		(0.002)	
log (Broadband Speed)	-0.125	0.27	-0.208	0.14	-0.260	0.08
	(0.113)		(0.14)		(0.146)	
Medical ICT Infrastructure						
Felehealth Policy (Stated)	0.002	0.98	0.009	0.91	-0.328	0.21
	(0.066)		(0.081)		(0.26)	
Telehealth Policy (Implied)	0.001	0.99	0.042	0.56	-0.357	0.06
	(0.055)		(0.071)		(0.183)	
Govt Health Websites	0.017	0.78	0.047	0.59	0.067	0.57

 ^a DV: Dependent Variable
 ^b Standard errors in parentheses
 ^c N/A: Not applicable

(0.061)		(0.085)		(0.117)	
0.055	0.44	0.042	0.65	-0.203	0.12
(0.07)		(0.091)		(0.128)	
-0.029	0.55	0.018	0.77	0.038	0.70
(0.049)		(0.062)		(0.096)	
				-0.015	0.94
				(0.183)	
				0.02	0.89
				(0.144)	
				0.415	0.18
				(0.304)	
				0.506	0.01
				(0.188)	
				-0.013	0.94
				(0.178)	>
				-0.107	0.50
				(0.159)	
3.131	< 0.001	3.078	< 0.001	3.163	< 0.001
(0.52)		(0.69)		(0.723)	
9	1	65		65	
0.3	41	0.329		0.44	
0.219 0.105		0.146			
	0.055 (0.07) -0.029 (0.049) 3.131 (0.52) 9	0.055	0.055 0.44 0.042 (0.07) (0.091) -0.029 0.55 0.018 (0.049) (0.062) 3.131 <0.001 3.078 (0.52) (0.69) 91 6 0.341 0.3	0.055 0.44 0.042 0.65 (0.07) (0.091) -0.029 0.55 0.018 0.77 (0.049) (0.062) 3.131 < 0.001 3.078 < 0.001 (0.52) (0.69) 91 65 0.341 0.329	0.055 0.44 0.042 0.65 -0.203 (0.07) (0.091) (0.128) -0.029 0.55 0.018 0.77 0.038 (0.049) (0.062) (0.096) -0.015 (0.183) 0.02 (0.144) 0.415 (0.304) 0.506 (0.188) -0.013 (0.178) -0.107 (0.178) 3.131 <0.001

^a DV: Dependent Variable

Our results suggest that medical ICT policy, rather than the ICT infrastructure itself, is negatively associated with the fatality rate. For example, the coefficient for implied telehealth policy in (Table 4), Column (2) is -2.68 and significant (p<0.01). That is, the fatality rate for countries with an implied telehealth policy was lower than that for others by 2.7 percentage point. Moreover, the coefficient for the rate of credit card ownership in (Table 4), Column (2) is -0.06 and statistically significant (p<0.05), suggesting that credit card usage could have lessened the fatality rate. However, broadband internet speed was not associated with any of the three measures of transmission of COVID-19. Lastly, the presence of a government-run health website showed a negative and significant relationship with the total number of confirmed cases in (Table 3), Column (2) (β =-0.44, p<0.05). This implies that countries with government-run health website had 36% fewer confirmed cases than those without it.

The effects of DistancingICT and MedicalICT on transmission speed of COVID-19 were not statistically significant. For the interaction terms, although most coefficients were insignificant, the interaction between implied telehealth policy and training ICT health was significant in (Table 5), Column (3) (β = 0.51, p<0.01). Therefore, an implied telehealth policy may be more effective when in-service training on ICT is provided to health professionals (β =-0.357+0.51=0.153).

Despite this not being the main focus of the study, it would be meaningful to examine the effects of other control variables in light of the lack of country-level empirical studies on the spread of COVID-19. Interestingly, early lockdowns, contrary to expectations, were statistically uncorrelated with the total number of infections and fatality rates. Moreover, the coefficients for GDP with the total number of confirmed cases and the fatality rate in (Table 4), Column (1) and (Table 4), Column

^b Standard errors in parentheses

^c N/A: Not applicable

(1) are 1.136 (p<0.01) and 1.635 (p<0.01), respectively. Population density is positively associated with the total number of cases and fatality rates, but temperature is negatively associated with the total number of infections. Lastly, the ratio of the elderly population and the total number of infections shows a negative relationship (β =-0.03; p<0.10) in (Table 3), Column (1).

As for the cultural dimensions of COVID-19, our results suggest no significant relationship between fatality rate and uncertainty avoidance, one of the two cultural dimensions from Hofstede Insights. However, individuals' tendency to care only for themselves and the immediate family, as represented by individualism, displayed a positive relationship with the fatality rate. The coefficient in (Table 4), Column (2) is 0.138 (p<0.01).

Additional Analysis

Although it has not been long since the COVID-19 outbreak and the transmission mechanism of the virus has not yet been clarified, it is apparent that human-to-human interaction increases the risk of infection [33,37]. Moreover, prior research has found a positive association between dense populations and infection rates [33], possibly because high density enhances the probability of an individual's exposure to the virus. However, people in such areas could be more aware of the risk, consequently taking precautions or complying with government regulations to avoid an epidemic. Moreover, highly concentrated urbanization is more likely to offer entrenched contact-free systems (e.g., delivery, retail kiosks) than less populated regions, thus stagnating or reducing the spread of virus. As such, the effect of DistancingICT or MedicalICT may vary substantially between countries with high and low population densities. Therefore, we conducted a further regression analysis on countries outside of the top 30% in population density in our sample. The results of further analysis are shown in (Tables S2-4 in Multimedia Appendix 1).

The key results are not remarkably different. Telehealth policy, rather than technology itself, may promote efficient management of COVID-19's aftermath regardless of population density. Unlike other national traits, the presence of telehealth policy —stated and implied— showed a negative association with the fatality rate when we omitted countries in the top 30% of population density from our sample. These consistent results highlight the importance of telehealth policy development in infection containment. However, it is noteworthy that credit card ownership was no longer significant. It is conceivable that countries with lower population density also have lesser rates of offline physical interaction, thus lowering the need for credit cards and online shopping.

Discussion

Principal Findings

In this study, we conducted exploratory research on the role of national characteristics, especially in regard to ICT and medical ICT infrastructures and concomitant policies that enable physical distancing, on the cumulative number of confirmed cases, fatality rates, and initial transmission speed of COVID-19. Findings suggested that medical ICT policies, especially when in-service training on ICT is provided, could potentially curb fatality rate. Government health websites were negatively associated with total number of confirmed cases. Moreover, possession of credit card was observed to decrease a fatality rate.

Discussion of Findings

The analysis results countered general intuition that ICT infrastructure should play a crucial role in slowing COVID-19 transmission. Overall, we found that the relationship between ICT infrastructure and COVID-19 infection or its consequences is less than expected. Nevertheless, there are some

important findings to highlight.

First, the presence of a telehealth policy manifests a negative correlation with the fatality rate. Surprisingly, only implied telehealth policy, but not stated telehealth policy shows statistically significant correlation with fatality rate. This raises the possibility that an implied telehealth policy may be more effective than a stated one because setting specific guidelines as in a stated policy sets boundaries that hinder clinicians' flexible decision making or system utilization amidst crisis [34]. The result remains consistent even when countries with high population concentrations are excluded. Moreover, it is important to note that the telehealth policy becomes more relevant when in-service training on ICT is provided. Such interaction result is aligned with WHO Digital Health Guidelines that state "Extensive training on the technology and operating the device should be done before introducing the system for use directly with clients" [38]. It is also notable that government health websites had a negative and significant relationship with the total number of confirmed cases. Consistent with preceding studies [39,40], it advocates governmental online communication in case of disease outbreaks to facilitate efficient interconnection between specialists from various fields. Otherwise, prompt and precise communication attempts by the government could enhance information transparency and built trust amongst public, increasing likelihood of public compliance to suggested guidelines including vaccine acceptance [41].

Second, our findings suggest that possession of a credit card, a widely used payment method for e-commerce [9,10], is related to a lower fatality rate. The results varied when countries with higher population density were omitted; this could happen either because of uneven development of contact-free systems in areas of lower density or because of prevention of crowding through widespread interventions such as contact tracing [42]. Nonetheless, such finding denotes how credit card ownership facilitates distancing compliance via online commerce. Alternatively, credit cards could have advanced financial inclusion and cushioned the financial burden even amongst challenging times. For instance, cardholders, especially those with low-income, have benefited from financial assistance programs including deferred payments, waived late fees, and even skipping payments from credit card issuers [43]. With financial assistance, perceived burden would have been decreased, fostering adherence to suggested guidelines [44].

Third, we had interesting findings from the control variables on the spread of COVID-19 and its aftermath. Early lockdown enforcement displayed no relationship with COVID-19 transmission, although policymakers intuitively assumed that compulsory restriction of contacts would be helpful in diminishing the total number of infections, fatality rate, and speed of transmission. Such intuitions are inferred from their actions to tighten stay-at-home restrictions [45]. Our result is consistent with the outcome of a preceding analysis that stated infection control was apparently effective before the mandate, and thus voluntary social behavior, rather than legal enforcement, is more crucial in combating a pandemic [46,47].

Stronger individualistic culture exhibited higher fatality rate whereas uncertainty avoidance showed no association with fatality rate. Related to individualism, recent study clarified the role of individualism-collectivism on perceived risk of COVID-19 and sense of responsibility [48]. In essence, individuals with strong collectivistic orientation perceived greater fear because of their higher physical and social interconnection with others [48]. Moreover, collectivism-oriented people, due to their strong sense of integrity and responsibility within society, are willing to follow containment guidelines whereas individualism-oriented people are not [49]. This finding is in line with preceding research that identified the relationship between pathogen risk and societal individualism [35]. Societal collectivism, which is more prevalent in Eastern cultures, did "[serve] as a natural guard against disease transmission" [36].

Previous study also revealed that countries with relatively high uncertainty avoidance are less likely to engage in public gathering, potentially decreasing number of infections and fatality rate [23]. However, in this study, such effect is not observed. The discrepancy could stem from different data collection periods. Huynh conducted an analysis at the initial stage of COVID-19, from February 16, 2020 to March 29, 2020 [23], but we used COVID-19 transmission data until July 28, 2020. It is challenging to refrain from public gatherings for several months; the impact of uncertainty avoidance would be weakened eventually.

The analysis result indicates that GDP is positively related to number of total cases and fatality rate. It is plausible that economically active countries, as represented by higher GDP, involve more interactions between individuals, causing an inevitable increase in the number of infections. Alternatively, in larger economies, the number of confirmed cases may reflect better testing because these countries have the economic capacity to conduct more tests. Lastly, higher proportion of elderly population proportion is correlated with fewer number of infections in total. The perceived risk of infection among elders could have possibly affected stay-at-home compliance, thus reducing physical contacts and infections.

Implications for Research and Practice

The results of this study pose several implications. Theoretically, we contribute to the effect of ICT infrastructure and policy in epidemics. Prior studies have examined ICT adoption intention of healthcare workers or how ICT use improves public health or physical wellness in general [50-53]. Moreover, past research on ICT and health have paid attention to how ICT mitigates various health-related challenges by providing access to health-related information and fostering communication between patients and physicians [54-56]. For example, previous research found that telephone usage for healthcare lowered rates of depression [54] and increased immunization rates [55]. However, they rarely showed interest for the ICT use in the context of epidemics, possibly due to its unlikelihood. Similarly, previous research on ICT policy in healthcare mainly focused on the "limitations concerning design and implementation of policies" of public health improvement [57]. Considering that ICT, by its nature, enables faster communication to the public, it is surprising that ICT effect on epidemics, which are widespread and abrupt, was not investigated sufficiently. In this study, we have addressed such void by examining the relationship between ICT policy with total cases, fatality rate and transmission speed under pandemic circumstance. Therefore, by expanding the scope on the role ICT on people's health, this study contributes to the literature on ICT and health-related challenges.

Practically, this study advocates autonomous use of medical ICT, rather than playing it by the book. Contrary to the assumption that detailed and rigorous policy statement limits or prevents a wide range of health threats such as smoking habits and cardiovascular disease [58,59], the findings indicate that stated medical ICT policy is in fact less likely to taper fatality rate than implied medical ICT policy. It is plausible that relatively less restrictions are helpful for better medical service because for jobs with high variety tasks, such as medical practice, increased autonomy boosts job performance of workers [60]. Also relevant in broader context of digital health field, this finding provides significant empirical insight that could improve the outcome of long-distance medical care and guide future clinical decision support system (CDSS) related strategies. This finding is aligned with WHO guidelines which suggest that "health workers may deviate from the recommendations" of CDSS based on physicians' own rationale [38], although algorithm based CDSS is conventionally perceived as competent. As such, policymakers need to consider independent and flexible decision-making of physicians in context of medical ICT usage.

Regarding distancing enabling infrastructure, this study implies that the government should prioritize

in providing ICT infrastructure that enable physical distancing in densely populated areas. As the budget for ICT infrastructure is limited, the government should strategically allocate budget to achieve greatest benefit. Especially, strategic budgeting is vital to developing countries where tax revenue is relatively insufficient. Because our finding shows that difference in population density yields different outcomes of ICT implementation, governors can consider investing in populous areas first, in order to maximize the profit with limited resources.

Conclusions

Despite our findings on the relationship between national characteristics and disease dispersion, our study is not without limitations. Although we included most established countries, we were not able to include all countries in our analysis. Consequently, the sample size for regression was small. In addition, the data for medical ICT infrastructure and the rate of credit card ownership were not up-todate. Therefore, there remains a possibility that their impact during the observation period may not been accurately estimated. However, it is important to note that ICT policy and infrastructure have a delayed "lag" effect on country-level outcomes because people need to adopt, trust and alter their behaviors in line with new technology and policies [61]. For instance, a recent study on the role of ICT on women's health outcomes showed that maternal fatality rate was lowered while modern medical care seeking behavior increased after kiosk was implemented and used for some years [61]. In addition, whereas we included broadband speed in the model, broadband coverage may also play a distinct role. For instance, high broadband speed offers fast communication via online, advocating real-time information sharing in dire situations such as COVID-19 [11]. Instead, broadband coverage enables seamless internet connectivity with personal devices; when individuals get out of the service range at some point, they would not get broadband access. While lack of decent broadband coverage indicates inability to use the Internet, lack of decent broadband speed denotes unattainability of prompt communication with others. Nevertheless, the correlation between broadband coverage and speed was high (Corr=.79). Accordingly, only one of them was used for our analysis to avoid a multicollinearity problem. Moreover, there are other potential confounders such as mask adherence that were not included in our study. However, we believe that our culture-related variables such as individualism and uncertainty avoidance may account for such confounders [62]. Lastly, because the COVID-19 pandemic has not ended, the long-term effect of the ICT infrastructure and ICT policies could not be examined.

By conducting an analysis at country level, we ensured the generalizability of our work and developed tentative guidelines to control the spread of infectious diseases. We have especially emphasized the importance of medical telehealth policies that contribute to reduce the consequences of COVID-19. By collecting updated COVID-19 data, future research can clarify the long-term effects of the aforementioned national traits. We hope that this study can broaden the scope of research on the impact of ICT infrastructure and policies and give guidance to better policy-making in the healthcare domain.

Acknowledgements

This work is (partially) supported by the Barun ICT Research Center at Yonsei University. This research was supported by the Yonsei University Research Fund of 2019 (2019-22-0051). This research was supported by the Yonsei Signature Research Cluster Program of 2021 (2021-22-0006).

Abbreviations

CPOE: computerized physician order entry

GDP PPP: gross domestic product by purchasing power parity

ICT: information and communication technology

OLS: ordinary least squares OWID: Our World in Data

UN: United Nations

WHO: World Health Organization VIF: variance inflation factor

References

- 1. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. 2020. https://covid19.who.int/ [accessed Sep 23, 2021].
- 2. Kupferschmidt K. Mutant coronavirus in the United Kingdom sets off alarms, but its importance remains unclear. Science. December 20, 2020. https://www.sciencemag.org/news/2020/12/mutant-coronavirus-united-kingdom-sets-alarms-its-importance-remains-unclear [accessed Jan 19, 2021].
- 3. Carter RW, Sanford JC. A new look at an old virus: patterns of mutation accumulation in the human H1N1 influenza virus since 1918. Theoretical Biology and Medical Modelling. 2012;9(1):1-19.
- 4. Grubaugh ND, Hanage WP, Rasmussen AL. Making sense of mutation: what D614G means for the COVID-19 pandemic remains unclear. Cell. 2020;182(4):794-5.
- 5. Jin X, Lian J-S, Hu J-H, Gao J, Zheng L, Zhang Y-M, et al. Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. Gut. 2020;69(6):1002-9.
- 6. United Nations Development Programme(UNDP). Socio-economic impact of COVID-19. 2020. https://www.undp.org/content/undp/en/home/coronavirus/socio-economic-impact-of-covid-19.html [accessed Jan 2, 2021].
- 7. Ghose A, Li B, Macha M, Sun C, Foutz NZ. Trading Privacy for the Greater Social Good: How Did America React During COVID-19? Available at SSRN 3624069. 2020.
- 8. Coronavirus: access to the internet can be a matter of life and death during a pandemic. Article 19. March 31, 2020. https://www.article19.org/resources/access-to-the-internet-can-be-a-matter-of-life-and-death-during-the-coronavirus-pandemic/ [accessed Feb 17, 2021].
- 9. B2C e-commerce and payments in North America: an assessment of COVID-19's impact on the industry. GlobeNewswire. Jul 6, 2020. https://www.globenewswire.com/news-release/2020/07/06/2057872/0/en/B2C-e-Commerce-and-Payments-in-North-America-An-Assessment-of-COVID-19-s-Impact-on-the-Industry.html [accessed Feb 17, 2021].
- 10. Meyll T, Walter A. Tapping and waving to debt: Mobile payments and credit card behavior. Finance Research Letters. 2019;28:381-7.
- 11. Chiou L, Tucker C. Social distancing, internet access and inequality. National Bureau of Economic Research, 2020 0898-2937.
- 12. Keshvardoost S, Bahaadinbeigy K, Fatehi F. Role of telehealth in the management of COVID-19: lessons learned from previous SARS, MERS, and Ebola outbreaks. Telemedicine and e-Health. 2020;26(7):850-2.
- 13. Draft O. Guide for Measuring ICTs in the Health Sector. Internal document for the OECD task forces. 2013;20120452.
- 14. Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, et al. Social contacts and mixing patterns relevant to the spread of infectious diseases. PLoS Med. 2008;5(3):e74.
- 15. Leung K, Jit M, Lau EH, Wu JT. Social contact patterns relevant to the spread of respiratory infectious diseases in Hong Kong. Scientific reports. 2017;7(1):1-12.
- 16. Coccia M. Factors determining the diffusion of COVID-19 and suggested strategy to prevent

- future accelerated viral infectivity similar to COVID. Science of the Total Environment. 2020;729:138474.
- 17. Denis E, Telle O, Benkimoun S, Mukhopadhyay P, Nath S. Mapping the lockdown effects in India: how geographers can contribute to tackle Covid-19 diffusion. The Conversation. 2020.
- 18. Bontempi E. First data analysis about possible COVID-19 virus airborne diffusion due to air particulate matter (PM): the case of Lombardy (Italy). Environmental research. 2020;186:109639.
- 19. Al-Hasan A, Khuntia J, Yim D. Threat, Coping, and Social Distance Adherence During COVID-19: Cross-Continental Comparison Using an Online Cross-Sectional Survey. Journal of medical Internet research. 2020;22(11):e23019.
- 20. Altmann S, Milsom L, Zillessen H, Blasone R, Gerdon F, Bach R, et al. Acceptability of appbased contact tracing for COVID-19: Cross-country survey study. JMIR mHealth and uHealth. 2020;8(8):e19857.
- 21. Sebhatu A, Wennberg K, Arora-Jonsson S, Lindberg SI. Explaining the homogeneous diffusion of COVID-19 nonpharmaceutical interventions across heterogeneous countries. Proceedings of the National Academy of Sciences. 2020;117(35):21201-8.
- 22. Jinjarak Y, Ahmed R, Nair-Desai S, Xin W, Aizenman J. Accounting for global COVID-19 diffusion patterns, January–April 2020. Economics of disasters and climate change. 2020;4(3):515-59.
- 23. Huynh TLD. Does culture matter social distancing under the COVID-19 pandemic? Safety Science. 2020;130:104872.
- 24. Atalan A. Is the lockdown important to prevent the COVID-19 pandemic? Effects on psychology, environment and economy-perspective. Annals of medicine and surgery. 2020;56:38-42.
- 25. Hofstede G. Culture's consequences: International differences in work-related values: sage; 1984. ISBN: 0803913060.
- 26. Yeung AY, Roewer-Despres F, Rosella L, Rudzicz F. Machine Learning—Based Prediction of Growth in Confirmed COVID-19 Infection Cases in 114 Countries Using Metrics of Nonpharmaceutical Interventions and Cultural Dimensions: Model Development and Validation. Journal of Medical Internet Research. 2021;23(4):e26628.
- 27. Zanaboni P, Knarvik U, Wootton R. Adoption of routine telemedicine in Norway: the current picture. Global Health Action. 2014;7(1):22801.
- 28. Centers for Disease Control and Prevention(CDC). Telehealth and Telemedicine: A Research Anthology of Law and Policy Resources. https://www.cdc.gov/phlp/publications/topic/anthologies/anthologies-telehealth.html [accessed July 8, 2021].
- 29. Jetty A, Moore MA, Coffman M, Petterson S, Bazemore A. Rural family physicians are twice as likely to use telehealth as urban family physicians. Telemedicine and e-Health. 2018;24(4):268-76.
- 30. Kircher SM, Mulcahy M, Kalyan A, Weldon CB, Trosman JR, Benson AB. Telemedicine in oncology and reimbursement policy during COVID-19 and beyond. Journal of the National Comprehensive Cancer Network. 2020;1(aop):1-7.
- 31. Wang J, Tang K, Feng K, Li X, Lv W, Chen K, et al. High temperature and high humidity reduce the transmission of COVID-19. arXiv preprint arXiv:200305003. 2020.
- 32. Hair JF. Multivariate data analysis. 2009.
- 33. Coccia M. Two mechanisms for accelerated diffusion of COVID-19 outbreaks in regions with high intensity of population and polluting industrialization: the air pollution-to-human and human-to-human transmission dynamics. MedRxiv. 2020.
- 34. Nancarrow SA. Six principles to enhance health workforce flexibility. Human resources for health. 2015;13(1):1-12.

35. Fincher CL, Thornhill R, Murray DR, Schaller M. Pathogen prevalence predicts human cross-cultural variability in individualism/collectivism. Proceedings of the Royal Society B: Biological Sciences. 2008;275(1640):1279-85.

- 36. Travers M. Cultural psychology research suggests the U.S. may be less individualistic after coronavirus pandemic. Forbes. March 23, 2020. https://www.forbes.com/sites/traversmark/2020/03/23/what-will-the-post-covid-19-world-look-like-lessons-from-cultural-psychology/?sh=69f540563f58 [accessed Feb 18, 2021].
- 37. Jarvis CI, Van Zandvoort K, Gimma A, Prem K, Klepac P, Rubin GJ, et al. Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. BMC medicine. 2020;18:1-10.
- 38. World Health Organization. WHO guideline: recommendations on digital interventions for health system strengthening. 2019. ISBN: 9241550503.
- 39. Komenda M, Bulhart V, Karolyi M, Jarkovský J, Mužík J, Májek O, et al. Complex reporting of the COVID-19 epidemic in the Czech Republic: Use of an interactive web-based app in practice. Journal of medical Internet research. 2020;22(5):e19367.
- 40. Ngai CSB, Singh RG, Lu W, Koon AC. Grappling with the COVID-19 health crisis: content analysis of communication strategies and their effects on public engagement on social media. Journal of medical Internet research. 2020;22(8):e21360.
- 41. Marlow LA, Waller J, Wardle J. Trust and experience as predictors of HPV vaccine acceptance. Human vaccines. 2007;3(5):171-5.
- 42. Hsu J. Population density does not doom cities to pandemic dangers. Scientific American. September 16, 2020. https://www.scientificamerican.com/article/population-density-does-not-doom-cities-to-pandemic-dangers/ [accessed Feb 23, 2021].
- 43. White A. Here's our most up-to-date list of credit card financial assistance programs during coronavirus. CNBC. July 8, 2021. https://www.cnbc.com/select/credit-card-issuers-offer-customer-assistance-amid-coronavirus-financial-hardship/ [accessed July 16, 2021].
- 44. Kim E, Gupta S, Bolge S, Chen C-C, Whitehead R, Bates JA. Adherence and outcomes associated with copayment burden in schizophrenia: a cross-sectional survey. Journal of Medical Economics. 2010;13(2):185-92.
- 45. Fancourt D. People started breaking Covid rules when they saw those with privilege ignore them. The Guardian. January 2, 2021. https://www.theguardian.com/commentisfree/2021/jan/02/follow-covid-restrictions-break-rules-compliance [accessed Feb 20, 2021].
- 46. Kepp KP, Bjornskov C. Lockdown Effects on Sars-CoV-2 Transmission—The evidence from Northern Jutland. medRxiv. 2020;2020.12. 28.20248936.
- 47. Gupta S, Nguyen TD, Rojas FL, Raman S, Lee B, Bento A, et al. Tracking public and private responses to the COVID-19 epidemic: evidence from state and local government actions. National Bureau of Economic Research, 2020 0898-2937.
- 48. Germani A, Buratta L, Delvecchio E, Mazzeschi C. Emerging adults and COVID-19: the role of individualism-collectivism on perceived risks and psychological maladjustment. International journal of environmental research and public health. 2020;17(10):3497.
- 49. Carvalho LdF, Pianowski G, Gonçalves AP. Personality differences and COVID-19: are extroversion and conscientiousness personality traits associated with engagement with containment measures? Trends in psychiatry and psychotherapy. 2020;42(2):179-84.
- 50. Melas CD, Zampetakis LA, Dimopoulou A, Moustakis V. Modeling the acceptance of clinical information systems among hospital medical staff: an extended TAM model. Journal of biomedical informatics. 2011;44(4):553-64.
- 51. Jimoh L, Pate MA, Lin L, Schulman KA. A model for the adoption of ICT by health workers in Africa. International journal of medical informatics. 2012;81(11):773-81.
- 52. Ganju KK, Pavlou PA, Banker RD. Does information and communication technology lead to the well-being of nations? A country-level empirical investigation. MIS Quarterly

- (Forthcoming). 2015:15-072.
- 53. Venkatesh V, Rai A, Sykes TA, Aljafari R. Combating Infant Mortality in Rural India: Evidence from a Field Study of eHealth Kiosk Imlementations. MIS Q. 2016;40(2):353-80.
- 54. Simon GE, Ludman EJ, Tutty S, Operskalski B, Von Korff M. Telephone psychotherapy and telephone care management for primary care patients starting antidepressant treatment: a randomized controlled trial. Jama. 2004;292(8):935-42.
- 55. Kaplan WA. Can the ubiquitous power of mobile phones be used to improve health outcomes in developing countries? Globalization and health. 2006;2(1):1-14.
- 56. Tan SS-L, Goonawardene N. Internet health information seeking and the patient-physician relationship: a systematic review. Journal of medical Internet research. 2017;19(1):e9.
- 57. Gelband H, Sankaranarayanan R, Gauvreau CL, Horton S, Anderson BO, Bray F, et al. Costs, affordability, and feasibility of an essential package of cancer control interventions in low-income and middle-income countries: key messages from Disease Control Priorities. The Lancet. 2016;387(10033):2133-44.
- 58. Busse R, Blümel M. Tackling chronic disease in Europe: strategies, interventions and challenges: WHO Regional Office Europe; 2010. ISBN: 9289041927.
- 59. Heidenreich PA, Trogdon JG, Khavjou OA, Butler J, Dracup K, Ezekowitz MD, et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. Circulation. 2011;123(8):933-44.
- 60. Dodd NG, Ganster DC. The interactive effects of variety, autonomy, and feedback on attitudes and performance. Journal of organizational behavior. 1996;17(4):329-47.
- 61. Venkatesh V, Sykes T, Zhang X. ICT for development in rural India: A longitudinal study of women's health outcomes. MIS Quarterly. 2020;44(2):605-29.
- 62. Nørgaard M, Ehrenstein V, Vandenbroucke JP. Confounding in observational studies based on large health care databases: problems and potential solutions—a primer for the clinician. Clinical epidemiology. 2017;9:185.

Supplementary Files

Multimedia Appendixes

Untitled.

URL: http://asset.jmir.pub/assets/582efd020092bac16d4ed9ca5e6632b7.docx