

Mobile Phones and HIV testing: Multi-country Instrumental Variable Analysis from Sub-Saharan Africa

Francesco Iacoella, Nyasha Tirivayi

Submitted to: Journal of Medical Internet Research
on: May 07, 2023

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 4

Supplementary Files..... 42

Figures 43

 Figure 1..... 44

 Figure 2..... 45

 Figure 3..... 46

Multimedia Appendixes 47

 Multimedia Appendix 0..... 48

Mobile Phones and HIV testing: Multi-country Instrumental Variable Analysis from Sub-Saharan Africa

Francesco Iacoella^{1*}; Nyasha Tirivayi^{2*}

¹Maastricht University Maastricht NL

²UNICEF Office of Research Innocenti Florence IT

*these authors contributed equally

Corresponding Author:

Francesco Iacoella
Maastricht University
Minderbroedersberg 4-6
Maastricht
NL

Abstract

Background: Sub-Saharan Africa carries a disproportionate burden of HIV infections and AIDS-related diseases. In 2016, of the 6000 new HIV infections happening globally every day, two out of three occurred in sub-Saharan Africa. The uptake of new technologies could help reduce this burden. SSA has been a powerhouse of mobile phone technology adoption over the past decades, yet the impact that mobile phones had on reproductive health has not been a research priority.

Objective: This study aims to fill this research gap by investigating the role of mobile phone connectivity on HIV testing in sub-Saharan Africa. We make use of the novel and comprehensive OpencellID cell tower database, and DHS geocoded information for over 400,000 women in 28 Sub-Saharan African countries.

Methods: Through the use of OLS and IV regressions, We examine whether women's community distance from the closest cell-tower influences knowledge about HIV testing facilities and the likelihood of ever been tested for HIV.

Results: We find a negative and significant impact of distance on our main outcomes (0.7 and 0.5 percentage point decrease respectively for each kilometer away from a cell tower). We investigate the mechanisms through which such effects might occur to find that proximity to a cell tower increases HIV-related knowledge as well as reproductive health knowledge. Similar results are observed when the analysis is performed at community level.

Conclusions: Results suggest that the effect of mobile phone connectivity is channelled through increased knowledge of HIV, STIs, and modern contraceptive methods. Further analysis shows that cell phone ownership has an even larger impact on HIV testing and knowledge. This paper adds to recent literature on the impact of mobile-based HIV prevention schemes by showing through large-scale analysis that better mobile network access is a powerful tool to spread reproductive health knowledge and increase HIV awareness.

(JMIR Preprints 07/05/2023:48794)

DOI: <https://doi.org/10.2196/preprints.48794>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in a peer-reviewed journal, I will be able to make the full manuscript available to all users.

Original Manuscript

**Mobile Phones and HIV testing:
Multi-country Instrumental Variable Analysis from Sub-Saharan
Africa**



Abstract**Background**

Sub-Saharan Africa has been a technological hothouse when it comes to mobile phone technology adoption. However, evidence on the role played by mobile technology on infectious disease prevention has been mostly limited to experimental studies.

Objective

This observational study investigates the role of mobile phone connectivity on HIV testing in sub-Saharan Africa.

Methods

We make use of the novel and comprehensive OpencellID cell tower database, and DHS geocoded information for over 400,000 women in 28 Sub-Saharan African countries. We examine, through OLS and IV regressions, whether women's community distance from the closest cell-tower influences knowledge about HIV testing facilities and the likelihood of ever been tested for HIV.

Results

After finding a negative and significant impact of distance on our main outcomes, we investigate the mechanisms through which such effects might occur. Our analysis shows that proximity to a cell tower increases HIV-related knowledge as well as reproductive health knowledge. Similar results are observed when the analysis is performed at community level.

Conclusions

Results suggest that the effect of mobile phone connectivity is channelled through increased knowledge of HIV, STIs, and modern contraceptive methods. Further analysis shows that cell phone ownership has an even larger impact on HIV testing and knowledge. This paper adds to recent literature on the impact of mobile-based HIV prevention schemes by showing through large-scale analysis that better mobile network access is a powerful tool to spread reproductive health knowledge and increase HIV awareness.

Introduction

Sub-Saharan Africa carries a disproportionate burden of HIV infections and AIDS related diseases.

In 2016, of the 6000 new HIV infections happening globally every day, two out of three occurred in sub-Saharan Africa [1]. Global AIDS-related deaths have sharply decreased in the past decade, with sub-Saharan African countries leading this downward trend [2], largely due to the increased access to anti-retroviral drugs. AIDS-related mortality has decreased by approximately 40% since 2010, and by 60% since it reached its peak in 2004 [2]. However, the number of new infections has not decreased at the same pace. Although the number of new HIV infections decreased by approximately 23% worldwide since 2010, with southern and eastern Africa showing the largest reduction, in 2019 there were over 1.7 million new infections, of which 1.4 million happened in Sub-Saharan Africa (and 59% of whom are women) [2]. More importantly, not every person living with HIV is aware of her condition [4]. Estimates for people aware of their condition in Sub-Saharan African countries range between 45% [3] and 65% [4], hence, the numbers of infected individuals are potentially being underestimated by the hundreds of thousands. In order to increase the capacity of governments to correctly identify HIV-infected individuals and therefore reduce the spreading of the virus, standard HIV-testing facilities and awareness raising campaigns are being paired with alternative methods such as community-based testing, self-test kits, and point-of-care testing for early infants [4]. Major barriers to testing (e.g. stigma, inaccessibility of services, and low awareness) are being gradually overcome by these new techniques which are currently being rolled-out in numerous Sub-Saharan countries [3–5]. Expanding HIV-related knowledge and introducing optimised HIV-testing techniques such as community-based testing and self-testing can potentially increase the returns from national anti-HIV programs [6].

Sub-Saharan Africa has proven to be a technological hothouse when it comes to the expansion of mobile phone coverage and use. The rapid increase in mobile subscribers' numbers and the expansion of network penetration has resulted in SSA becoming the world's fastest-growing mobile region. Mobile phones, facilitate individuals' access to information from sources that would

otherwise be unreachable or costly and enable near-instant communication through voice and text (or video, in the case of smartphones). Mobile phone penetration had a pivotal role in the region, influencing economic growth human development [7,8]. It has been argued that the “communication” component of mobile phones holds the most potential of supporting economic development, by creating knowledge-sharing networks [9–11]. Recently, researchers have been investigating the role of mobile phones in public health and found them to be powerful awareness raising tools and facilitators of individuals’ communication with health workers and other patients [12–14]. For mobile phone users, new information is, therefore, acquired mostly through communication with other individuals. However, mobile phones can also facilitate access to institutional resources (e.g. by receiving informative text messages from a health clinic, listening to radio, or, for those with access to smartphone, surfing the internet). Evidence from rural Indonesia has shown that midwives’ use of mobile phones increased access to institutional resources and, consequently, fostered better reproductive health knowledge. Further, access to peer resources was associated with higher self-efficacy, which was as well positively associated with health knowledge [15]. Mobile phone ownership has been found to significantly influence contraception use and reproductive health attitudes [16,17]. Phones have also been used in recent years as a new way of reaching HIV-infected and at-risk individuals with awareness raising campaigns (see Project Masiluleke in South Africa (<https://www.innovations.harvard.edu/project-masiluleke>) and Text to Change in Uganda (<http://images.aarogya.com/aids/pdf/unandzain.pdf>)). Finally, recent evidence identified mobile phone ownership as a more relevant determinant of contraceptive use and reproductive healthcare access than TV or radio [18].

Some studies have focused on investigating mobile-based awareness-raising messaging and adherence to HIV/AIDS treatment programmes through randomized control trials [19–23]. Lester et al. [18], in a seminal study on this topic, examined the association between higher adherence to antiretroviral treatment (ART) and SMS-based support in Kenya. More recent studies have

questioned the role of SMS-based support as a “silver bullet” in resource-limited settings, especially after second-line ART failure, although it still remains a valuable tool for promoting higher participation in anti-HIV/AIDS programmes in general [19]. A study on the role of mobile apps for newly diagnosed HIV-positive patients found that young individuals in particular are more responsive to mobile-based treatment adherence campaigns [20], while another is planning to test an innovative self-testing service that uses AI [23]. However, the impact of mobile technology on HIV, and HIV-testing knowledge and uptake might go way beyond the effectiveness of SMS-based or app-based support.

The lack of literature on the topic makes it difficult to formulate hypotheses on the role played by cellular phones, although it has been shown that innovative mobile-based awareness-raising campaigns at community- and individual-level represent an economically sensible alternative to traditional awareness raising campaigns [4]. Moreover, mobile technology-enabled networks facilitate peer-to-peer communication, which has been proven to increase HIV knowledge and treatment adherence, while reducing HIV risk [24–26]. Finally, mobile devices may accord privacy in communication, which could facilitate the discussion of a topic such as HIV prevention which is still stigmatized. We hypothesise that mobile-enabled communication increases HIV- and reproductive-health-related knowledge and, through that, it affects HIV testing knowledge and uptake.

Considering the high number of new infections that happen every year in Sub-Saharan African countries, and the fact that a large portion of people living with HIV are still unaware of their condition, it is crucial to better understand if and how mobile technology can support HIV prevention and testing. Although there is an emerging body of evidence linking mobile phones with health promotion and awareness, to our knowledge, evidence on the role played by mobile phone ownership or connectivity on HIV-testing knowledge and attitudes, and an investigation of impact pathways, is lacking. Our study seeks to address this knowledge gap. It examines the role mobile connectivity can

play in promoting HIV testing through multi-country analysis in Sub-Saharan Africa. We focus on mobile connectivity instead of mobile ownership to account for the fact that phone sharing is very common in Sub-Saharan African countries [9,13]. Our study is, to our knowledge, the first of its kind to establish a connection between widespread access to mobile phone network and knowledge of HIV testing facilities and ultimately HIV testing uptake. We utilize recent data on cell-tower locations and data on women's HIV testing and knowledge in 28 sub-Saharan African countries. In our paper, we also explore the mechanisms through which mobile phone connectivity affects HIV testing knowledge and uptake. We test if mobile connectivity has a direct effect on levels of HIV knowledge, and whether it influences overall knowledge about reproductive health (i.e. knowledge about other sexually transmissible diseases (STIs) and about contraceptive methods). We find that indeed mobile connectivity influences general knowledge about HIV and reproductive health.

With our research, we establish a causal link between mobile network connectivity and HIV testing knowledge and practices using individual-level data from 28 African countries and we shed a light on potential mechanisms for these results. Our work fills a significant gap in HIV-testing literature by providing observational evidence and investigating the role of both mobile connectivity and mobile phone ownership.

Methods

Data

Several data sources have been used to build the dataset used in our analysis.

Cell connectivity information

Information on cell towers' location is derived from OpenCellid dataset. OpenCellid is the largest, daily updated, publicly available database of cell towers. Linear distance between household clusters and the nearest cell tower is calculated in kilometres and serves as a proxy for cell connectivity and signal strength and as our main explanatory variable. OpenCellid datasets reports location, operator, and type of signal for over 40 million towers world-wide (additional information at <https://opencellid.org/>). Database entries for the African continent start in 2008. Unfortunately, the date of entry does not unequivocally represent the date of construction of a tower. Because to this, many countries show total absence of cell towers before year 2013, a clearly misleading representation. This is why we calculate distance from nearest cell tower including all cell towers in each country regardless of the date in which they have been added to OpenCellid dataset, under the assumption that towers have been built before their date of entry in the dataset. Since this is a very strong assumption, we test its robustness by running our analysis again, this time limiting the number of cell towers per country to those added to OpenCellid before the year in which individual survey data was collected. As we will show in a later section, results remain consistent even when the number of cell towers (and, therefore, of countries) is reduced. One more issue with OpenCellid data is that tower location is not verified after it is inserted in the database, which might result in a discrepancy between reported and actual location. However, one study comparing independently collected data on cell towers coordinates and OpenCellid information on the same towers shows that the localization error estimated from OpenCellid data agrees well with the experimental error distribution [27]. As an extra precaution, however, we consider only towers which coordinates' maximum error (reported by OpenCellid) lies within one square kilometre.

[Figure 1 about here]

Individual and community level data

Individual data is obtained from the Demographic and Health Survey (DHS) Program from 28 African countries – Angola, Benin, Burkina Faso, Burundi, Cameroon, Chad, Comoros, Ivory Coast, Democratic Republic of Congo, Ethiopia, Gabon, Ghana, Guinea, Kenya, Lesotho, Liberia, Malawi, Mali, Mozambique, Namibia, Nigeria, Rwanda, Senegal, Sierra Leone, Tanzania, Togo, Uganda, Zambia, and Zimbabwe. DHS provides cross-sectional information on women and household access to health and health status together with demographic and labour statistics. Besides women's demographic and employment information, we take from DHS our main dependent variables: a dummy for knowledge of facilities where women could be tested for HIV and a dummy for ever been tested for HIV. We use additional variables on knowledge of reproductive health to test our hypothesis on possible impact pathways. Individual-level characteristics are added to the analysis to serve as confounders. We include information on woman age, education, employment, marital status, if ever had children, and exposure to media (i.e. newspapers, radio, or television). We also include household-level indicators of gender of the household head, place of living (i.e. urban or rural), and a wealth index build by DHS and based on selected assets, dwelling construction materials, and types of water access and sanitation facilities. Finally, to address the role played by ease of access to health facilities in facilitating HIV testing, we include in our analysis two dummy variables for women having visited a health facility in the past 12 months and for perceiving distance to health facilities as a problem (The authors thought about using distance in kilometres to the nearest health facility as a valid alternative to this variable. However, evidence shows that perception of distance might play a bigger role than actual distance when deciding whether to go to health facility for testing [28]). For each country, DHS surveys included in this study have been conducted between 2010 and 2016. The final sample comprises 407,000 women grouped in 19,603 clusters, of who approximately 350,000 are included in the main analysis. As DHS surveys are conducted to be representative at country

level, our study provides estimates relevant for the entire population of the countries involved in it. Geospatial information of households in DHS datasets is clustered to guarantee respondents' privacy. For these clusters, DHS provides spatial information retrieved from secondary sources within a buffer of 10km for rural communities, and 2km for urban ones. We include in the analysis community-level (i.e. cluster-level) information on average travel time to the nearest settlement (i.e. an urban area with at least 50,000 inhabitants) and population density. We also use household-level information on access to electricity to generate a community-level electrification rate indicator. This indicator is simply the percentage of households that have access to electricity within a cluster. Living a long distance from the nearest cell tower might be one of many indicators of a community's geographical isolation. More isolated communities could have lower level of HIV testing regardless of their use of mobile phones [29]. To better measure community isolation, and to control for it in the analysis, we calculate the linear distance of each households' cluster from the nearest primary or secondary road (trails and non-paved routes have not been considered in the analysis). Information on roads in the African continent is obtained through the Digital Chart of the World (DCW) vector basemap. Although last updated in 1998, DCW remains a very detailed map of Africa's road system. Authors have considered using more recent street maps for the analysis, however they ultimately decided to make use of DCW due to its reliability. NASA gROADS (<http://tiny.cc/5grl8y>) data set, although more recent, does not provide users the chance to adequately distinguish types of roads in the African continent. However, for those countries in which comparison between DCW and gROADS was possible (e.g. Zimbabwe), the authors verified that primary roads reported in the latter were also present in the former (graphical representation in Figure A2 in appendix). Distance to road is used, together with distance to nearest settlement and community-level electrification rate, as a proxy for community isolation. Finally, we include information on land gradient and HIV prevalence for each of the clusters. This information is obtained from DHS elaboration of the United States Geological Survey GTOPO30.

HIV prevalence (as number of HIV positive people over 100,000) within a 10km-radius from the cluster is obtained from Dwyer-Lindgren et al. (2019) mapping exercise, which produced a 5x5 Km grid of HIV prevalence among adults for Sub-Saharan Africa. Grids are provided for every year between 2000 and 2017. Since 2010 represents the less recent years in which surveys included in our dataset were collected, we use HIV prevalence information from that year.

Data is analysed using QGIS [31] and Stata 16 [32] software.

Sample characteristics

Table 1 reports descriptive statistics from the sample. The average community distance from cell tower is 13 km. This is considering both rural and urban areas. Average woman's age is 28 years old, with 5 years of education. About 28% of interviewed women live in a female-headed household, 72% are exposed to at least one type of media (i.e. radio, television, newspaper), 52% are in union, 59% are currently working, and 35% live in an urban area. Half of the women in the sample report to have visited a health facility in the last 12 months, although distance from said facility is considered a problem by 41% of them. Over 70% of sampled women have given birth to at least one child. The average population density in the communities was 1290 people per square kilometre (this is considering a buffer of 2 km² for urban areas and 10 km² for rural areas, 2015 population data). Distance from closest road (trails and non-paved routes have been excluded from the analysis) is 6 km on average and average travel time to closest settlement is about 77 minutes. Finally, the prevalence of households with access to electricity in the community is about 32% on average.

Table 1. Descriptive statistics of the sample

	Values	N
Variables		
Distance from nearest cell tower (Km)	13.06	517,290
Woman age	28.46	517,290
Woman years of education	5.18	517,110
Female-headed households (%)	28.34	517,290
Woman exposed to any type of media (%)	72.05	517,257
Woman currently working (%)	58.6	500,001
Ever had a child (%)	72.66	517,290
Number of lifetime sex partners	2.16	407,542
Woman in union (%)	52.34	517,284
Living in urban area (%)	35.4%	517,290
Population density in community (2015) [†]	1290.64	516,411

<i>Wealth Index quintiles</i>		517,290
Poorest (%)	19.42	
Poorer (%)	18.56	
Middle (%)	19.01	
Richer (%)	19.96	
Richest (%)	23.05	
Average travel time to closest settlement (min) [†]	77.48	516,411
Distance from closest road (km) [†]	6.12	517,290
Average HIV prevalence in 10km radius	5.05	517,022
Visited a health facility in the past 12 months (%)	50.31	499,767
Distance to health facility is considered a problem (%)	41.04	458,976
Electrification (cluster-level) [†] (%)	31.81	517,290

Methodology

Using probit and OLS regressions, we design a naïve model at first.

$$(1) HIV t_i = \beta_0 + \beta_1 D_i + \beta_2 Z_i + \beta_3 C + u_i$$

Where $HIV t_i$ is a dummy variable indicating if either woman knows where HIV testing facility is or has ever been tested for HIV. β_1 is a vector of our main explanatory variable, or distance to nearest cell tower D for woman i . and β_2 is a vector of covariates Z for woman i . A full list of covariates is present in Table 1. Finally, β_3 is a vector of community-level characteristics and country-fixed effects C . Location information, from which distance from cell towers was calculated, could only be obtained at community level. Community clusters contain between 1 and 92 households each. In order to address intra-cluster correlation, we cluster standard errors at community level in each regression. Common practice is to cluster standard errors at the most aggregate level [33]. However, it is important to consider that in our case sampling has been done at cluster level and, more importantly, the treatment (i.e. distance from cell-tower) is measured at community level. Correlation in standard errors at community level is, therefore, very plausible and clustering errors at a higher level might be unnecessarily conservative [34]. The authors decided to present results with standard errors clustered at community level to address such correlation. However, robustness checks have been performed by clustering errors at country and administrative area level. Both models produced similar and significant results (results available upon request). De-normalised individual sample weights are included. Individual women weights have been calculated

by de-normalising DHS individual weights following Ruilin Ren note on DHS weight de-normalisation (see <https://userforum.dhsprogram.com/index.php?t=getfile&id=767&S=Google>). As a robustness check, analysis has been run without weights and results remain consistent (results available upon request).

As Klonner and Nolen (2010) observe, cell towers' distribution is not exogenous over a country's territory. It might follow an established path (i.e. roads) and be affected by power supply and other infrastructural needs. Moreover, network providers might consider certain areas not profitable enough to repay them of the investment of building a cell tower. Figure 2 shows this endogenous distribution in the context of Zimbabwe.

[Figure 2 about here]

In order to deal with endogeneity, we look for an instrument that could influence towers' location but is uncorrelated with our dependent variable error terms. Following Dinkelman et al. (2011), we utilize average community land gradient as an instrumental variable. Terrain slope raises the costs of electric line routing and antennas' building [36–38], acting as a disincentive for mobile network companies to build cell towers in areas with higher land gradient. On the contrary, more competitive markets in Sub-Saharan Africa can lead to a considerable increase in cell-phone area coverage [39]. Proxies for market competition have been used in the past in tandem with terrain roughness to explain cell towers' positioning [40]. We build our instrumental variable interacting land gradient and number of phone service providers in each country following Berman et al. [41]. We also add to the list of covariates information on household electrification to control for effects related only to access to electricity and not to access to cell-phone network. Our second model, a two-stage regression is, therefore:

$$(2) D_i = \pi_0 + \pi_1 \text{Slope} * \text{Network Providers}_i + \pi_2 Z_i + \pi_3 \text{Elec}_i + \pi_4 C + \varepsilon_i$$

$$(3) HIV_{it} = \beta_0 + \beta_1 \hat{D}_i + \beta_2 Z_i + \beta_3 \text{Elec}_i + \beta_4 C + u_i$$

In equation (2), the measure of land gradient *Slope* multiplied by number of *Network Providers* is the

exogenous regressors. Electrification *Elec*, ranges between 0 and 1 and represents the percentage of households that have access to electricity in the community. Equation (3) is similar to equation (1), although here distance *D* has been instrumented by *Slope * Network Providers*. Land gradient is calculated by DHS based on United States Geological Survey Global 30 Arc-Second elevation model (GTOPO30) [42], and we consider it invariant through time (or, at least, within the same geological era). This model is estimated via two stage least squares (2SLS).

The identification assumption is that conditional on individual and community characteristics, our instrumental variable does not directly affect knowledge of facilities or decision to test for HIV, and only does so through the distance from the nearest cell tower. To check our assumption, we perform several placebo tests. Mobile phone signal spreads unevenly in the area surrounding cell towers due to a variety of factors. Given the geographical extent of our analysis, we are not able to calculate network coverage exactly. However, technical limitations of standard GSM towers set their maximum signal range to a 35 Km radius, therefore communities located more than 35 Km away from the closest cell tower (9.2% of the sample) can be considered not covered by mobile network. The 35 km maximum range is the one imposed by technical limitations on GSM towers; however, it does not take into account barriers to signal that might be present within the radius of the cell tower. This is why the dummy build with this cut-off point is used only as an addition to our main explanatory variable (i.e. distance from cell tower). See <https://www.3gpp.org/technologies> for additional information on timing advance and TDMA technology, and the way they influence GSM towers range. For the sake of standardisation, we assume towers do not possess features to extend their range. Using this cut-off point, which theoretically divides the sample into connected and non-connected communities, we examine whether our instrumental variable only affects outcomes when communities are covered by a mobile network. Results (Table A1 in appendix) demonstrate that our instrument is only correlated with outcomes in network-covered communities. Our second test investigates whether the instrumental variable or instrumented distance from cell tower are in any

way correlated to proxies of communities' remoteness, a factor that could in turn affect knowledge of HIV testing place and test uptake. We select three different proxies: distance from nearest road (excluding trails and unpaved roads), exposure to any type of media as proxies for remoteness; and average population density in the community as a proxy for dispersion. Both the instrument and the instrumented distance from nearest cell tower show no correlation with any of the proxies, supporting our initial assumption that longer distance from a cell tower does not necessary results from the remoteness of a community (Table A2 in appendix for full results). Lastly, we test the sensibility of our instrumental variable. Our results might be driven by countries where markets are more accessible for mobile phone providers. Competition might decrease mobile technology consumption costs and facilitate knowledge sharing. We run our models once again, this time dividing the sample based on the number of mobile network providers in country of residence. Results remain consistently negative throughout the models (see Figure 3), confirming that the relationship between distance from cell towers and our outcomes does not change based on mobile network market characteristics.

[Figure 3 about here]

Ethical considerations

The study uses secondary data analysis from DHS. Procedures and questionnaires for standard DHS surveys have been reviewed and approved by ICF Institutional Review Board (IRB). Study data are anonymous.

Data Availability

All data used in this study is available on DHS repository, pending authorization, and OpenCellid repository.

Results

Mobile phone connectivity and knowledge of HIV-testing facilities

Our analysis examines whether better mobile connectivity is correlated with women's knowledge of

which facilities offer HIV testing. Results from are presented in Table 2. Probit regression results show a strong negative correlation ($p < .001$) between distance from cell tower and knowledge of an HIV testing facility. Results from the 2SLS estimation are consistent with our Probit results. Due to the nature of the variables, interpreting the magnitude of the impact is not straightforward. We could infer that for each unit increase in distance to the nearest cell tower, the probability of knowing where one can be tested for HIV is 0.7 percentage points lower (0.2 p.p. for Probit analysis). To facilitate the interpretation of results and to put them in the right order of magnitude, we report (in Figure A1 in Appendix) a graphical illustration of our IV model's predicted share of women knowledgeable about testing facilities by deciles of distance to nearest cell tower. As it can be noticed, 90% or more of women living in the top deciles are knowledgeable about HIV testing facilities locations. The share starts dropping starting from the 4th decile, going below 80% in the bottom two deciles (i.e. with a distance $>15\text{km}$), and reaching 40% in the lowest decile (i.e. distance $>33\text{km}$).

The clustering of households' coordinates allows us to calculate distance from nearest cell-tower at community-level only. To check whether this might affect our results, we perform community-level analysis by averaging women characteristics at community level. The impact of distance to cell towers remains highly significant ($P < .001$). The magnitude of the impact also remains consistent, with a 0.8 p.p. decrease in the probability of knowing testing facility (0.1 p.p. for Probit). Underidentification and weak-instrument (Kleibergen-Paap Wald F statistic) tests for all IV regressions show that our instrument is relevant and reasonably strong.

Table 2. Impact of phone connectivity on knowledge of HIV testing facilities.

	Knowledge of HIV testing facility			
	Individual-level		Community-level	
	Probit	IV (2SLS)	Probit	IV (2SLS)
Distance from nearest cell (km)	-0.002*** [<0.001]	-0.007*** [0.002]	-0.001*** [<0.001]	-0.008*** [0.002]
Individual-level characteristic	Yes	Yes	/	/
Community-level characteristics	Yes	Yes	Yes	Yes

Country-year-fixed effects	Yes	Yes	Yes	Yes
Constant	-0.547*** [0.058]	0.505*** [0.026]	0.304*** [0.021]	0.399*** [0.038]
Observations	351,613	351,488	18,437	18,431
R-squared	0.307	0.183	0.682	0.402
Underidentification (p-value)		<0.001		<0.001
Kleibergen-Paap rk Wald F statistic		32.13		28.74

Note: Cluster-robust standard errors in brackets. *** p<.001. Underidentification and weak-instrument (Kleibergen-Paap rk Wald F statistic) tests reported at bottom of the table for IV regressions.

From knowledge to action – mobile connectivity and HIV testing

We have identified a significantly positive effect of proximity to cell towers on knowledge of HIV testing facilities. It is important to also determine whether knowledge acquired through mobile phones ultimately translates into a behavioral change. We therefore investigate whether women living closer to cell towers not only acquire knowledge on where HIV testing can be conducted, but also get tested.

Newly acquired knowledge affects individuals' health behaviour through what has been called the knowledge-attitude-behaviour continuum [43]. Studies show that when agents act on the way information is processed by individuals, one could expect desired behavioural changes to ultimately happen [43]. Activities aimed at increasing knowledge about HIV transmission among pregnant women have been shown to change attitudes and behaviour towards HIV testing [44,45]. At the same time, access to peer networks can influence individuals' attitudes towards HIV testing. Studies in the past have found that peer-based interventions can increase HIV knowledge and adherence to AIDS treatment [46–48].

We examine whether better mobile connectivity has a direct effect on women's probability of ever been tested for HIV. Our analysis is operationalised in a variation of equations (1) and (3) where having tested for HIV is now the dependent variable and the explanatory variable is distance from nearest cell tower. Table 3 reports the results of our analysis. We find that a unit increase in distance from cell towers significantly reduces women's probability of ever been tested for HIV by 0.5 p.p. in the individual-level model and by 0.4 p.p. in the community-level model. Again, we present in Figure A1 in Annex a graphical illustration of our IV model's predicted share of women tested by

deciles of distance to nearest cell tower. Similarly to knowledge of testing facilities, share of tested women is constant at around 70% in the top four deciles. At the start of the bottom two deciles, share of tested women is already below 60% and it reaches approximately 10% in the lowest decile.

Table 3. Impact of phone connectivity on HIV testing uptake

	Ever tested for HIV			
	Individual-level		Community-level	
	Probit	IV	Probit	IV
Distance from nearest cell (km)	-0.005*** [0.001]	-0.005 (P=0.02) [0.002]	-0.001*** [<0.001]	-0.005*** [0.002]
Individual-level characteristic	Yes	Yes	/	/
Community-level characteristics	Yes	Yes	Yes	Yes
Country-year-fixed effects	Yes	Yes	Yes	Yes
Constant	-1.383*** [0.050]	0.147*** [0.025]	-0.036* [0.019]	0.014 [0.029]
Observations	365,108	364,982	18,480	18,474
R-squared	0.323	0.343	0.793	0.734
Underidentification (p-value)		<0.001		<0.001
Kleibergen-Paap rk Wald F statistic		34.21		29.08

Note: Cluster-robust standard errors in brackets. *** p<.001. Underidentification and weak-instrument (Kleibergen-Paap rk Wald F statistic) tests reported at bottom of the table for IV regressions

Robustness checks

Reduced number of cell towers

As a first robustness check, we examine whether the results remain significant when distance from nearest cell tower is calculated using only the cell towers inserted in the OpenCellid dataset starting from the year in which each DHS survey was collected. Several countries are dropped from this analysis due to the complete absence of data on cell towers from the year of the survey onwards, namely: Cameroon, Comoros, Gabon, Guinea, Mozambique, and Senegal. However, figures presented in Table 4 show that results remain significant and consistent with previous findings.

Table 4. Impact of phone connectivity on HIV testing (restricted cell towers' sample)

	Knowledge of HIV testing facility				Ever tested for HIV			
	Individual-level		Community-level		Individual-level		Community-level	
	OLS	IV	OLS	IV	OLS	IV	OLS	IV
Distance from nearest cell (km)	<0.001 (P=0.84) [<0.001]	-0.003*** [0.001]	<0.001*** [<0.001]	-0.002*** [<0.001]	<0.001*** [<0.001]	-0.002 (P=0.004) [0.001]	<0.001*** [<0.001]	-0.001 (P=0.007) [<0.001]
Individual-level characteristic	Yes	Yes	/	/	Yes	Yes	/	/
Community-level characteristics	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Country-year-fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes

Constant	-0.636*** [0.066]	0.475*** [0.025]	0.335*** [0.025]	0.369*** [0.041]	-1.568*** [0.059]	0.118*** [0.023]	-0.043* [0.022]	-0.019 [0.032]
Observations	247,902	247,777	12,867	12,861	261,802	261,676	12,910	12,904
(pseudo) R-squared	0.315	0.0897	0.676	0.234	0.354	0.312	0.814	0.643
Underidentification (p-value)		<0.001		<0.001		<0.001		<0.001
Kleibergen-Paap rk								
Wald F statistic		41.26		13.48		43.80		13.41

Note: Cluster-robust standard errors in brackets. *** p<.001. Underidentification and weak-instrument (Kleibergen-Paap rk Wald F statistic) tests reported at bottom of the table for IV regressions.

Impact on other health care seeking and utilization

One question that can arise when looking at our results is whether the impact of mobile connectivity on knowledge of HIV testing facilities and testing uptake is driven by a general impact of access to mobile phones on health seeking and utilization. For example, better connectivity may increase demand for health services from the population while at the same time reducing costs to access such services (e.g. increased knowledge and better networking may provide individuals with more options for obtaining health care). To test whether this is the case, we repeat our analysis and replace our outcomes of interest with dummy variables for having health insurance and having delivered one's last child in a safe structure ("Safe structure" is defined here as either a public or a private health centre). The results, presented in Table A4 in appendix, are not statistically significant in any of our preferred models (i.e. 2SLS estimations). These results suggest that increased mobile connectivity does not have a general impact on healthcare seeking and utilization.

Alternative instrumental variable

As we have shown in our methodology section, terrain roughness and market competition are often associated with infrastructural development. However, there are additional characteristics that influence cell tower positioning in developed and developing countries alike. As an additional robustness check, we test the validity of lightning intensity and cost of mobile phone usage as alternative instruments.

Frequent electrostatic discharges damage mobile phone infrastructures and negatively affect connectivity. Areas with high lightning intensity represent a risk for mobile phone providers since

power surge protection for cell towers is costly and reduces profits from their investments. Being Africa the continent with the highest lightning intensity in the world, evidence has showed that 1 standard deviation increase in lightning intensity leads to a lower penetration rate of mobile phone technology of approximately 0.43 percentage points per year [49]. Mobile phone usage depends on a variety of factors, of which cost is surely one of the most important [50]. Higher costs might be driven by low competition and, at the same time, act as an incentive for new companies to provide network access to a larger share of the population.

We interact average intensity of lightning strikes in a 10km radius around each DHS cluster with country-level monthly cost of running a mobile phone as a percentage of per-capita income. Lightning strikes intensity data is publicly available at <https://ghrc.nsstc.nasa.gov/lightning/>. The satellite data is collected by NASA's Global Hydrology Resource Center. We included in our analysis yearly average lightning strikes for countries included in the analysis for 2010. Lightning strike incidence is proved to be stable through years, making the yearly average for 2010 a good proxy for the subsequent decade [49]. Information on monthly cost of running mobile phone by country is retrieved from 2015 ITU report on Measuring the Information Society, available at <https://www.itu.int/en/ITU-D/Statistics/Documents/publications/misr2015/MISR2015-w5.pdf>. A slight modification to equation (2) produces

$$(4) D_i = \pi_0 + \pi_1 \text{Flash intensity} * \text{Mobile cost}_i + \pi_2 Z_i + \pi_3 \text{Elec}_i + \pi_4 C + \varepsilon_i$$

Analysis conducted using the alternative instrumental variable produces results consistent with previous findings which remain statistically significant (results presented in Table A3 in Appendix).

Mobile phone ownership instead of connectivity

We have focused our analysis on the impact of mobile network connectivity on women's HIV and health knowledge and practices regardless of their ownership of a cellular phone. We did so because evidence has shown that mobile phone sharing is a common practice in Sub-Saharan Africa [9]. It is

the intention of the authors to argue for the potential of better phone connectivity regardless of increased ownership, and our results have supported our claims. However, it is logical to think that the easier the access to a mobile phone the more relevant its impact on our outcomes of interest should be, and ownership of a mobile phone grants a much higher exposure to mobile technology than borrowing. DHS surveys have recently started including questions on mobile phone ownership in their questionnaires as part of individuals' and households' assets. This provides us with a reduced sample of women that reported whether or not they own a cellular phone. Information on phone ownership is present for Angola (2015), Burundi (2016), Ethiopia (2016), Malawi (2015), Tanzania (2015), Uganda (2016), and Zimbabwe (2015).

To analyse the role of phone ownership, we make use of Probit models as in equation (1), while we specify a new Probit-2SLS model for the instrumental variable analysis (in this case, we use the same instrument from our main analysis, the interaction between land gradient and number of network providers). Given the binary nature of the phone ownership variable, a binary probit model for the first stage regression is more adequate. We present results in Table 5. According to our IV estimation, phone ownership increases the chance of knowing a facility that conducts HIV tests by 8 p.p., and it increases the chance of ever been tested for HIV by approximately 14 p.p. Considering the percentage of people knowledgeable about HIV testing place and ever tested for HIV who do not own a phone, our percentage points translate into a 9% higher chance of knowing a testing facility and a 22% higher chance of having tested for HIV for mobile phone owners.

Table 5. Phone ownership and HIV testing

	Knowledge of HIV testing facility		Ever tested for HIV	
	Probit	IV	Probit	IV
Mobile phone owner	0.220*** [0.040]	0.078*** [0.016]	0.240*** [0.028]	0.142*** [0.025]
Individual-level characteristic	Yes	Yes	Yes	Yes
Community-level characteristics	Yes	Yes	Yes	Yes
Country-year-fixed effect	Yes	Yes	Yes	Yes
Constant	-0.765*** [0.148]	0.708*** [0.012]	-1.654*** [0.101]	0.183*** [0.015]
Observations	76,241	76,241	81,048	81,048
(pseudo) R-squared	0.300	0.183	0.318	0.320

Note: Cluster-robust standard errors in brackets. *** p<.001

Mechanisms for impact

Cell phone connectivity has been found to positively impact probability of knowing HIV testing facilities and of ever been tested for HIV. Through which mechanisms such impact is achieved is what we investigate in this section. To do so, we need to understand the barriers to HIV testing that mobile phone connectivity might help overcome. Studies have shown that stigma, both self- and social-, remains the most significant barrier to HIV testing in Sub-Saharan Africa [51,52]. Provision of HIV counselling for high-risk subjects has yielded good results among adolescents thanks to increased knowledge and awareness, although cultural change is hard to achieve [53]. Physical barriers to HIV testing are also limiting the effectiveness of awareness raising programs. Distance from health facilities, inconvenient opening hours, and long waiting lines are among the reasons cited for low test uptake [53].

Mobile connectivity has been proven to be a powerful tool to spread information and increase social cohesion in the Global South [54]. Mobile phones also facilitate knowledge diffusion which in turn reduces information asymmetry [7,55]. We argue that, while better mobile phone connectivity might do little to reduce physical barriers to HIV testing, it can increase women's knowledge of HIV related symptoms, can help fight HIV misconception, and foster reproductive health's knowledge.

To test our hypotheses, we perform analysis on the role of distance from nearest cell tower on several intermediate outcomes. We use as dependent variables knowledge of HIV and sexually transmittable infections (STIs), and knowledge of modern contraceptive methods. HIV knowledge is measured, following standard DHS guidelines, as an index ranging from 0 to 1 measuring knowledge of HIV symptoms and of preventive measures [56]. Knowledge of STIs and modern contraceptive methods are dummies assigning value 1 to knowledgeable women. Results of the analysis are presented in Table 6.

Distance from cell towers appears to have a significantly negative impact on HIV- and STI-related

knowledge. In the same way, it also reduces the chance that a woman would be knowledgeable about modern contraceptive methods. These results seem to confirm our hypothesis that better cellular connectivity facilitates acquisition of knowledge on reproductive health and sexually transmittable diseases. These findings are supported by previous evidence on the role played by mobile phones in knowledge-sharing and learning [15,57,58].

Table 6. Analysis of mechanisms of cell connectivity and HIV testing correlation

	Knowledge of HIV		Knowledge of STIs		Knowledge of contraceptive method	
	OLS	IV	Probit	IV	Probit	IV
Distance from nearest cell (km)	<-0.001 (P=0.02) [<0.001]	-0.004*** [0.001]	-0.003*** [0.001]	-0.002*** [0.001]	-0.002*** [0.001]	-0.004*** [0.001]
Individual-level characteristic	Yes	Yes	Yes	Yes	Yes	Yes
Community-level characteristics	Yes	Yes	Yes	Yes	Yes	Yes
Country-year-fixed effect	Yes	Yes	Yes	Yes	Yes	Yes
Constant	0.577*** [0.012]	0.760*** [0.022]	-0.153** [0.077]	0.741*** [0.016]	-0.660*** [0.069]	0.640*** [0.018]
Observations	340,062	339,965	359,431	368,812	369,000	368,874
(pseudo) R-squared	0.222	0.114	0.258	0.0505	0.345	0.0784
Underidentification (p-value)		<0.001		<0.001		<0.001
Kleibergen-Paap rk Wald F statistic		31.52		34.01		34.01

Note: Cluster-robust standard errors in brackets. *** p<.001. Underidentification and weak-instrument (Kleibergen-Paap rk Wald F statistic) tests reported at bottom of the table for IV regressions.

To complete our analysis on mechanisms, we test whether results remain consistent when considering mobile phone ownership instead of mobile connectivity. Findings are reported in Table 7. Our results show that phone ownership significantly increases knowledge about sexual transmittable infections and modern contraceptive methods, while it seems to have no statistically significant impact on general knowledge of HIV.

Recent evidence has shown that mobile phone ownership might have an impact on contraceptive use and knowledge about family planning [59], and facilitate the spread of reproductive, maternal, newborn and child health in vulnerable settings [60]. At the same time, some research has argued that better outcomes for phone owners are mostly driven by their economic status [61]. In our analysis we control for additional factors that could influence women's knowledge, such as media exposure,

education, asset-based wealth, and place of residence, and the effect of mobile ownership remains statistically significant.



Table 7. Phone ownership and mechanisms

	Knowledge of HIV		Knowledge of STIs		Knowledge of contraceptive method	
	Probit	IV	Probit	IV	Probit	IV
Mobile phone owner	0.012*** [0.005]	0.006 (P=0.639) [0.013]	0.243*** [0.050]	0.053*** [0.016]	0.193*** [0.051]	0.089*** [0.015]
Individual-level characteristic	Yes	Yes	Yes	Yes	Yes	Yes
Community-level characteristics	Yes	Yes	Yes	Yes	Yes	Yes
Country-year-fixed effect	Yes	Yes	Yes	Yes	Yes	Yes
Constant	0.651*** [0.019]	0.702*** [0.011]	-0.527*** [0.129]	0.722*** [0.014]	-0.297** [0.137]	0.733*** [0.014]
Observations	66,344	66,344	81,048	81,048	81,048	81,048
Pseudo-R ²			0.325		0.478	
R-squared	0.296	0.239		0.174		0.236

Note: Cluster-robust standard errors in brackets. *** p<.001.

Discussion

This paper analyses the impact of cell phone connectivity on women HIV testing knowledge and behaviour. We used distance from the nearest cell tower as a proxy for phone connectivity strength and quality and we estimated how increasing distance from said cell tower affects the probability of knowing where one can be tested for HIV and the probability of having taken such a test at least once in one's life. Mobile phones enhance individuals' access to information thanks primarily to knowledge-sharing through communication, although they can also put people in contact with institutional resources [15,58]. Due to phone-sharing practices in Sub-Saharan Africa we used cell phone connectivity as the primary indicator for analysis, with mobile phone ownership used for robustness check. We attempt to expand on the work that has been done on a smaller scale on the role played by mobile technology on HIV awareness-raising and testing [19,21]. We make use of a pooled cross-sectional dataset from 28 Sub-Saharan African countries and different methods of analysis to produce robust inference on the role played by phone connectivity.

Results show that mobile connectivity has a statistically significant and positive impact on the knowledge of HIV testing facilities and having been tested for HIV, highlighting its role as an important determinant of these practices. Results are consistent in all models and at both individual and community level. We test the hypothesis and find evidence that the impact of mobile phone connectivity on HIV testing knowledge and behaviour might be driven by improved knowledge of HIV, STIs, and contraceptive methods. Moreover, we find that the positive effects of mobile technology are even greater when we consider phone ownership as the explanatory variable.

Our results are in agreement with previous evidence that shows the role that mobile technology can play in raising awareness about HIV and HIV testing [62]. The study also fits well into the

new body of literature investigating mHealth and the role it could play in developing countries. Promising evidence shows how informal mHealth and access to mobile phones among young Africans can help bridge healthcare gaps in countries like Ghana, Malawi, and South Africa [63,64]. Not only mobile phones, but all digital technology has been found to be beneficial for treating and preventing mental disorders in low-income and middle-income countries [65]. Our results support the idea that stronger mobile connectivity can foster HIV testing practices in Sub-Saharan Africa.


Our study is not exempt from limitations. First, although we are able to establish a link between mobile connectivity and knowledge of HIV testing facilities and mobile connectivity and HIV testing uptake, due to data limitations we cannot dig deeper into the way mobile connectivity shapes women's attitudes toward testing through better knowledge, since no questions were asked to interviewed women about their opinion of HIV testing. Second, OpenCellid dataset is a relatively new dataset that might be lacking detail especially for remote areas. Missing information on the date in which cell towers are built also poses a limitation. We addressed both these issues by only including cell towers with the most reliable GPS coordinates and by performing robustness checks that aligned the building year of the cell towers with the year of the survey. When and if data will be available in the future, additional longitudinal analysis could be conducted to investigate the evolution of HIV testing behaviour through time for women with better mobile connectivity. A third limitation is represented by the clustering of individuals' position under one GPS coordinate representing the community. This clustering allows distance from cell tower to be calculated only at community-level. We show, however, that our results remain significant even if individual figures are averaged at cluster level and community-level analysis is performed.

Our study provides useful insights on the potential that new communication technologies

represent to foster better reproductive health practices in Sub-Saharan Africa, and paves the way for future analysis on how mobile phones enhance knowledge-sharing around HIV/AIDS knowledge, behaviour and practices.

Conflict of interest

The authors have no conflict of interest to declare.



Bibliography

1. Kharsany ABM, Karim QA. HIV Infection and AIDS in Sub-Saharan Africa: Current Status, Challenges and Opportunities. Open AIDS J Bentham Science Publishers; 2016;10:34–48. PMID:27347270
2. UNAIDS. UNAIDS data 2019. 2019. Available from: https://www.unaids.org/sites/default/files/media_asset/2019-UNAIDS-data_en.pdf [accessed Aug 3, 2020]
3. Sweat M, Fonner V. Overcoming resistance to HIV testing in sub-Saharan Africa. Lancet HIV Elsevier; 2016 Mar 1;3(3):e106–e107. doi: 10.1016/S2352-3018(16)00004-7
4. UNAIDS. Knowledge is power — Know your status, know your viral load. 2018. Available from: https://www.unaids.org/sites/default/files/media_asset/jc2940_knowledge-is-power-report_en.pdf [accessed Aug 3, 2020]
5. Jani I V, Meggi B, Loquiha O, Tobaiwa O, Mudenyanga C, Zitha A, Mutsaka D, Mabunda N, Vubil A, Bollinger T, Vojnov L, Peter TF. Effect of point-of-care early infant diagnosis on antiretroviral therapy initiation and retention of patients. AIDS AIDS; 2018;32(11):1453–1463. PMID:29746301
6. Swann M. Economic strengthening for HIV testing and linkage to care: a review of the evidence. AIDS Care AIDS Care; 2018;30(sup3):85–98. PMID:29985055
7. Asongu SA, Boateng A, Akamavi RK. Mobile Phone Innovation and Inclusive Human Development: Evidence from Sub-Saharan Africa. African Gov Dev Inst. 2016 Mar. doi: 10.2139/ssrn.2822103
8. Lee SH, Levendis J, Gutierrez L. Telecommunications and economic growth: an empirical analysis of sub-Saharan Africa. Appl Econ Routledge ; 2012 Feb;44(4):461–469. doi: 10.1080/00036846.2010.508730
9. Aker JC, Mbiti IM. Mobile Phones and Economic Development in Africa. The J Econ

- Perspect 2010;24(3):207–230. Available from:
<https://www.jstor.org/stable/pdf/20799163.pdf?refreqid=excelsior%3A22ec4be2eb589923661e3e5bf632e6e7> [accessed Nov 11, 2019]
10. Donner J. The Use of Mobile Phones by Microentrepreneurs in Kigali, Rwanda: Changes to Social and Business Networks. *Inf Technol Int Dev MIT Press*; 2006 Dec;3(2):3–19. Available from: <http://dev.itidjournal.org/index.php/itid/article/view/221> [accessed Oct 1, 2019]
 11. Nakato GV, Beed F, Bouwmeester H, Ramathani I, Mpiira S, Kubiriba J, Nanavati S. Building agricultural networks of farmers and scientists via mobile phones: case study of banana disease surveillance in Uganda. *Can J Plant Pathol Taylor & Francis*; 2016 Jul 2;38(3):307–316. doi: 10.1080/07060661.2016.1230149
 12. Anstey Watkins JOT, Goudge J, Gómez-Olivé FX, Griffiths F. Mobile phone use among patients and health workers to enhance primary healthcare: A qualitative study in rural South Africa. *Soc Sci Med* 2018 Feb;198:139–147. doi: 10.1016/j.socscimed.2018.01.011
 13. Kaplan WA. Can the ubiquitous power of mobile phones be used to improve health outcomes in developing countries? *Global Health BioMed Central*; 2006 May 23;2(1):9. doi: 10.1186/1744-8603-2-9
 14. Chen S, Fang Y, Chan PS-F, Kawuki J, Mo P, Wang Z. Counseling Supporting HIV Self-Testing and Linkage to Care Among Men Who Have Sex With Men: Systematic Review and Meta-Analysis. *JMIR public Heal Surveill JMIR Public Health and Surveillance*; 2024 Jan 24;10(1):e45647. PMID:38265866
 15. Lee S, Chib A, Kim J-N. Midwives' Cell Phone Use and Health Knowledge in Rural Communities. *J Health Commun Taylor & Francis Group* ; 2011 Oct;16(9):1006–1023. doi: 10.1080/10810730.2011.571344

16. Greenleaf AR, Ahmed S, Moreau C, Guiella G, Choi Y. Cell phone ownership and modern contraceptive use in Burkina Faso: implications for research and interventions using mobile technology. *Contraception Elsevier*; 2019 Mar 1;99(3):170–174. doi: 10.1016/J.CONTRACEPTION.2018.11.006
17. LeFevre AE, Shah N, Bashingwa JJH, George AS, Mohan D. Does women's mobile phone ownership matter for health? Evidence from 15 countries. *BMJ Glob Heal BMJ Specialist Journals*; 2020 May 1;5(5):e002524. PMID:32424014
18. Iacoella F, Gassmann F, Tirivayi N. Which communication technology is effective for promoting reproductive health? Television, radio, and mobile phones in sub-Saharan Africa. Olapeju BB, editor. *PLoS One Public Library of Science*; 2022 Aug 17;17(8):e0272501. doi: 10.1371/journal.pone.0272501
19. Gross R, Ritz J, Hughes MD, Salata R, Mugenyi P, Hogg E, Wieclaw L, Godfrey C, Wallis CL, Mellors JW, Mudhune VO, Badal-Faesén S, Grinsztejn B, Collier AC. Two-way mobile phone intervention compared with standard-of-care adherence support after second-line antiretroviral therapy failure: a multinational, randomised controlled trial. *Lancet Digit Heal* 2019;1:26–34. Available from: www.thelancet.com/ [accessed Aug 26, 2019]
20. Lester RT, Ritvo P, Mills EJ, Kariri A, Karanja S, H Chung M, Jack W, Habyarimana J, Sadatsafavi M, Najafzadeh M, A Marra C, Estambale B, Ngugi E, Ball TB, Thabane L, J Gelmon L, Kimani J, Ackers M, Plummer FA. Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomised trial. *Lancet* 2010;376:1838–45. doi: 10.1016/S0140-6736(10)61997-6
21. Amankwaa I, Boateng Id D, Quansah DY, Akuoko CP, Evans C. Effectiveness of short message services and voice call interventions for antiretroviral therapy adherence and

- other outcomes: A systematic review and meta-analysis. *PLoS One* 2018;18(9). doi: 10.1371/journal.pone.0204091
22. Kliner M, Knight A, Mamvura C, Wright J, Walley J. Using no-cost mobile phone reminders to improve attendance for HIV test results: a pilot study in rural Swaziland. *Infect Dis Poverty BioMed Central*; 2013 Jun 14;2(1):12. doi: 10.1186/2049-9957-2-12
23. Chen S, Zhang Q, Chan C-K, Yu F-Y, Chidgey A, Fang Y, Mo PKH, Wang Z. Evaluating an Innovative HIV Self-Testing Service With Web-Based, Real-Time Counseling Provided by an Artificial Intelligence Chatbot (HIVST-Chatbot) in Increasing HIV Self-Testing Use Among Chinese Men Who Have Sex With Men: Protocol for a Noninferiority Randomized Controlled Trial. *JMIR Res Protoc JMIR Research Protocols*; 2023 Jun 30;12(1):e48447. PMID:37389935
24. Kaponda CPN, Jere DL, Chimango JL, Chimwaza AF, Crittenden KS, Kachingwe SI, McCreary LL, Norr JL, Norr KF. Impacts of a Peer-Group Intervention on HIV-Related Knowledge, Attitudes, and Personal Behaviors for Urban Hospital Workers in Malawi. *J Assoc Nurses AIDS Care* No longer published by Elsevier; 2009 May 1;20(3):230–242. doi: 10.1016/J.JANA.2008.12.005
25. Marino P, Simoni JM, Silverstein LB. Peer Support to Promote Medication Adherence Among People Living with HIV/AIDS. *Soc Work Health Care* NIH Public Access; 2007 Jul 2;45(1):67–80. PMID:17804348
26. Mbeba MM, Kaponda CPN, Jere DL, Kachingwe SI, Crittenden KS, McCreary LL, Norr JL, Norr KF. Peer Group Intervention Reduces Personal HIV Risk for Malawian Health Workers. *J Nurs Scholarsh* John Wiley & Sons, Ltd; 2011 Mar 1;43(1):72–81. doi: 10.1111/j.1547-5069.2011.01384.x
27. Ulm M, Widhalm P, Brandle N. Characterization of mobile phone localization errors with

- OpenCellID data. 2015 4th Int Conf Adv Logist Transp IEEE; 2015. p. 100–104. doi: 10.1109/ICAdLT.2015.7136601
28. Hawthorne TL, Kwan M-P. Using GIS and perceived distance to understand the unequal geographies of healthcare in lower-income urban neighbourhoods. *Geogr J* 2012;178(1). Available from: <https://www.jstor.org/stable/pdf/41475787.pdf?refreqid=excelsior%3A544a6e180ff6a37d012592efc800ea55> [accessed Nov 19, 2019]
29. Susan A, Karita E, N'gandu N, Tichacek A. The Evolution of Voluntary Testing and Counseling as an HIV Prevention Strategy. 2002. p. 87–108. doi: 10.1007/0-306-47157-4_5
30. Dwyer-Lindgren L, Cork MA, Sligar A, Steuben KM, Wilson KF, Provost NR, Mayala BK, VanderHeide JD, Collison ML, Hall JB, Biehl MH, Carter A, Frank T, Douwes-Schultz D, Burstein R, Casey DC, Deshpande A, Earl L, El Bcheraoui C, Farag TH, Henry NJ, Kinyoki D, Marczak LB, Nixon MR, Osgood-Zimmerman A, Pigott D, Reiner RC, Ross JM, Schaeffer LE, Smith DL, Davis Weaver N, Wiens KE, Eaton JW, Justman JE, Opio A, Sartorius B, Tanser F, Wabiri N, Piot P, Murray CJL, Hay SI. Mapping HIV prevalence in sub-Saharan Africa between 2000 and 2017. *Nature* Nature Publishing Group; 2019 Jun 15;570(7760):189–193. doi: 10.1038/s41586-019-1200-9
31. QGIS Development Team. QGIS Geographic Information System. Open Source Geospatial Foundation Project; 2016.
32. StataCorp LLC. Stata Statistical Software: Release 16. College Station, TX; 2019.
33. Colin Cameron A, Miller DL. A Practitioner's Guide to Cluster-Robust Inference. *J Hum Resour* University of Wisconsin Press; 2015 Apr 28;50(2):317–372. doi: 10.3368/jhr.50.2.317
34. Abadie A, Athey S, Imbens G, Wooldridge J. When Should You Adjust Standard Errors for

- Clustering? Cambridge, MA; 2017 Nov. Report No.: 24003. doi: 10.3386/w24003
35. Dinkelman T, Beaman L, Beuermann D, Busso M, Case A, Choi H, Dinardo J, Ferris A, Fisman R, Keyes B, Lam D, Levinsohn J, Martinez-A C, Mccrary J, Qian N, Posel D, Potter-Gunter S, Rosenzweig M, Smith J, Solon G, Tybout J, Thornton R, Wittenberg Substantially M. The Effects of Rural Electrification on Employment: New Evidence from South Africa. *Am Econ Rev* 2011; Available from: https://rpls.princeton.edu/sites/rpls/files/media/dinkelman_electricity_0810.pdf [accessed Jan 22, 2020]
 36. Monteiro C, Fernandez-Jimenez LA, Zorzano-Santamaria PJ, Garcia-Garrido E, Miranda V, Ramirez-Rosado JJ. GIS Spatial Analysis Applied to Electric Line Routing Optimization. *IEEE Trans Power Deliv* 2005;20(2):934–942. doi: 10.1109/tpwr.2004.839724
 37. Klonner S, Nolen P. Does ICT Bene...t the Poor? Evidence from South Africa. 2008. Available from: <https://pdfs.semanticscholar.org/0ce4/92f33b778dee001f29aff461a970a2bd4d3d.pdf> [accessed Dec 4, 2019]
 38. Batzilis D, Dinkelman T, Oster E, Thornton R, Zanera D. New Cellular Networks in Malawi: Correlates of Service Rollout and Network Performance. *African Successes, Vol III Mod Dev University of Chicago Press*; 2016. p. 215–245. doi: 10.7208/chicago/9780226315867.003.0006
 39. Buys P, Dasgupta S, Thomas TS, Wheeler D. Determinants of a Digital Divide in Sub-Saharan Africa: A Spatial Econometric Analysis of Cell Phone Coverage. *World Dev Pergamon*; 2009 Sep 1;37(9):1494–1505. doi: 10.1016/J.WORLDDEV.2009.01.011
 40. Pierskalla JH, Hollenbach FM. Technology and Collective Action: The Effect of Cell

- Phone Coverage on Political Violence in Africa. *Am Polit Sci Rev* 2013;107(2). doi: 10.1017/S0003055413000075
41. Berman N, Couttenier M, Rohner D, Thoenig M. This Mine is Mine! How Minerals Fuel Conflicts in Africa. *Am Econ Rev* 2017 Jun 1;107(6):1564–1610. doi: 10.1257/aer.20150774
42. Mayala B, Fish TD, Eitelberg D, Dontamsetti T. The DHS Program Geospatial Covariate Datasets Manual. 2018.
43. Bettinghaus EP. Health promotion and the knowledge-attitude-behavior continuum. *Prev Med (Baltim)* Academic Press; 1986 Sep 1;15(5):475–491. doi: 10.1016/0091-7435(86)90025-3
44. Mahmoud MM, Nasr AM, Gasmelseed DEA, Abdalelhafiz MA, Elsheikh MA, Adam I. Knowledge and attitude toward HIV voluntary counseling and testing services among pregnant women attending an antenatal clinic in Sudan. *J Med Virol* 2007 May;79(5):469–473. doi: 10.1002/jmv.20850
45. Rahbar T, Garg S, Tripathi R, Gupta V, Singh MM. Knowledge, attitude, behavior and practice (KABP) regarding HIV/AIDS among pregnant women attending PPTCT programme in New Delhi. *J Commun Dis* 2007;39(3):179–84. Available from: [https://www.semanticscholar.org/paper/%22Knowledge%2C-attitude%2C-behavior-and-practice-\(KABP\)-Rahbar-Garg/88376c0155c91f9e14138728bcc65fbb5c76e23b](https://www.semanticscholar.org/paper/%22Knowledge%2C-attitude%2C-behavior-and-practice-(KABP)-Rahbar-Garg/88376c0155c91f9e14138728bcc65fbb5c76e23b) [accessed Jun 14, 2020]
46. Broadhead RS, Heckathorn DD, Altice FL, van Hulst Y, Carbone M, Friedland GH, O'Connor PG, Selwyn PA. Increasing drug users' adherence to HIV treatment: results of a peer-driven intervention feasibility study. *Soc Sci Med Pergamon*; 2002 Jul 1;55(2):235–246. doi: 10.1016/S0277-9536(01)00167-8

47. Haignere CS, Freudenberg N, Silver DR, Maslanka H, Kelley J. One Method for Assessing HIV/AIDS Peer-Education Programs. *J Adolesc Heal* 1997;21(2):76–9. Available from: [https://research.steinhardt.nyu.edu/scmsAdmin/uploads/007/088/Haignere Freudenberg Silver et al.pdf](https://research.steinhardt.nyu.edu/scmsAdmin/uploads/007/088/Haignere_Freudenberg_Silver_et_al.pdf) [accessed Jun 15, 2020]
48. Pearlman DN, Camberg L, Wallace LJ, Symons P, Finison L. Tapping youth as agents for change: evaluation of a peer leadership HIV/AIDS intervention. *J Adolesc Heal Elsevier*; 2002 Jul 1;31(1):31–39. doi: 10.1016/S1054-139X(02)00379-8
49. Manacorda M, Tesei A. Liberation Technology: Mobile Phones and Political Mobilization in Africa. *Econometrica* 2020 Nov 1;88(2):533–567. doi: 10.2139/ssrn.3392757
50. World Bank. World Development Report 2016: Digital Dividends. Washington DC; 2016. doi: 10.1596/978-1-4648-0671-1
51. Mohlabane N, Tutshana B, Peltzer K, Mwisongo A. Barriers and facilitators associated with HIV testing uptake in South African health facilities offering HIV Counselling and Testing. *Heal SA Gesondheid* 2016 Oct 11;21:86–95. doi: 10.4102/hsag.v21i0.938
52. Ayieko J, Brown L, Anthierens S, Van Rie A, Getahun M, Charlebois ED, Petersen ML, Clark TD, Kamya MR, Cohen CR, Bukusi EA, Havlir D V., Camlin CS. “Hurdles on the path to 90-90-90 and beyond”: Qualitative analysis of barriers to engagement in HIV care among individuals in rural East Africa in the context of test-and-treat. Costa AB, editor. *PLoS One Public Library of Science*; 2018 Aug 30;13(8):e0202990. doi: 10.1371/journal.pone.0202990
53. Chikwari CD, Dringus S, Ferrand RA. Barriers to, and emerging strategies for, HIV testing among adolescents in sub-Saharan Africa. *Curr Opin HIV AIDS* 2018 May;13(3):257–264. PMID:29401121
54. Ling R, Horst HA. Mobile communication in the global south. Ling R, Horst HA, editors.

- New Media Soc SAGE PublicationsSage UK: London, England; 2011 May 31;13(3):363–374. doi: 10.1177/1461444810393899
55. Asongu SA, Nwachukwu JC. The Mobile Phone in the Diffusion of Knowledge for Institutional Quality in Sub-Saharan Africa. *World Dev Pergamon*; 2016 Oct 1;86:133–147. doi: 10.1016/J.WORLDDEV.2016.05.012
56. Wang W, Alva S, Wang S. HIV-Related Knowledge and Behaviors among People Living with HIV in Eight High HIV Prevalence Countries in Sub-Saharan Africa. Calverton, Maryland, USA; 2012. Available from: www.measuredhs.com [accessed Oct 7, 2019]
57. Aker JC, Ksoll C, Lybbert TJ. Can Mobile Phones Improve Learning? Evidence from a Field Experiment in Niger. *Am Econ J Appl Econ* 2012;4:94–120. doi: 10.1257/app.4.4.94
58. Qureshi S. What is the role of mobile phones in bringing about growth? *Inf Technol Dev Routledge* ; 2013 Jan 30;19(1):1–4. doi: 10.1080/02681102.2013.764597
59. Jadhav A, Weis J. Mobile phone ownership, text messages, and contraceptive use: Is there a digital revolution in family planning?.. *Contraception Elsevier*; 2019 Nov 26; doi: 10.1016/J.CONTRACEPTION.2019.10.004
60. Alami SO, Dulli L, Dal Santo L, Rastagar SH, Seddiqi S, Hemat S, Machlin Burke J, Todd CS. Getting health information to internally displaced youth in Afghanistan: can mobile phone technology bridge the gap? *Gates Open Res F1000 Research Limited*; 2019 Oct 23;3:1483. doi: 10.12688/gatesopenres.13008.2
61. Nie J, Unger JA, Thompson S, Hofstee M, Gu J, Mercer MA. Does mobile phone ownership predict better utilization of maternal and newborn health services? a cross-sectional study in Timor-Leste. *BMC Pregnancy Childbirth BioMed Central*; 2016 Dec 23;16(1):183. doi: 10.1186/s12884-016-0981-1
62. Venter W, Coleman J, Chan VL, Shubber Z, Phatsoane M, Gorgens M, Stewart-Isherwood


- L, Carmona S, Fraser-Hurt N. Improving Linkage to HIV Care Through Mobile Phone Apps: Randomized Controlled Trial. JMIR mHealth uHealth JMIR Publications Inc.; 2018 Jul 17;6(7):e155. PMID:30021706
63. Hampshire K, Porter G, Owusu SA, Mariwah S, Abane A, Robson E, Munthali A, DeLannoy A, Bango A, Gunguluza N, Milner J. Informal m-health: How are young people using mobile phones to bridge healthcare gaps in Sub-Saharan Africa? Soc Sci Med Elsevier Ltd; 2015 Oct 1;142:90–99. doi: 10.1016/j.socscimed.2015.07.033
64. Cilliers L, Viljoen KL-A, Chinyamurindi WT. A study on students' acceptance of mobile phone use to seek health information in South Africa. Heal Inf Manag J SAGE PublicationsSage UK: London, England; 2018 May 7;47(2):59–69. doi: 10.1177/1833358317706185
65. Naslund JA, Aschbrenner KA, Araya R, Marsch LA, Unützer J, Patel V, Bartels SJ. Digital technology for treating and preventing mental disorders in low-income and middle-income countries: a narrative review of the literature. The Lancet Psychiatry Elsevier; 2017 Jun 1;4(6):486–500. doi: 10.1016/S2215-0366(17)30096-2

Supplementary Files

Figures


Cell phone coverage in Sub Saharan African countries included in our study.

Figure 1. Cell phone coverage in Sub Saharan African countries included in our study



Cell towers distribution in Zimbabwe.


Figure 2. Cell towers distribution in Zimbabwe



Source: OpenCellid and Authors' elaboration on QGIS

Placebo test.

Figure 3. Placebo test



Note: Number of providers correspond, from left to right, to the 50th percentile of the sample, the 75th percentile, the 90th percentile, and the 100th percentile.

Multimedia Appendixes

Additional robustness tests.

URL: <http://asset.jmir.pub/assets/a3dfa9d6cadb4bac320260cb7dbf14cf.docx>

