

A pilot study of the role of social capital in a Parkinson's disease risk cohort: feasibility of a web application to enable online social interactions

Xiancheng Li, Aneet Gill, Pietro Panzarasa, Jonathan P Bestwick, Anette Schrag,
Alastair J Noyce, Anna De Simoni

Submitted to: JMIR Formative Research
on: August 18, 2023

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 28

..... 28



A pilot study of the role of social capital in a Parkinson's disease risk cohort: feasibility of a web application to enable online social interactions

Xiancheng Li¹ PhD; Aneet Gill² BSc, MSc; Pietro Panzarasa¹; Jonathan P Bestwick² BSc; Anette Schrag³; Alastair J Noyce²; Anna De Simoni⁴ MBBS PhD

¹School of Business and Management Queen Mary University of London London GB

²Preventive Neurology Unit Wolfson Institute of Population Health Queen Mary University of London London GB

³Queen Square Institute of Neurology University College London London GB

⁴Centre for Primary Care Wolfson Institute of Population Health Queen Mary University of London London GB

Corresponding Author:

Anna De Simoni MBBS PhD

Centre for Primary Care

Wolfson Institute of Population Health

Queen Mary University of London

58 Turner Street

Centre for Primary Care

London

GB

Abstract

Background: There is evidence that social interaction has an inverse association with the development of neurodegenerative diseases. PREDICT-PD is an online UK cohort study that stratifies participants for risk of future Parkinson's disease (PD).

Objective: This study aims to explore the methodological approach and feasibility of assessing the digital social characteristics of people at risk of developing PD and their social capital within the PREDICT-PD platform.

Methods: A web application was built to enable social interaction via the PREDICT-PD portal. Feedback from existing members of the cohort was sought and informed the design of the pilot. Dedicated staff used weekly engagement activities, consisting of PD-related research, facts, and queries to stimulate discussion. Data were collected by the hosting platform. We examined the pattern of connections generated over time through the cumulative number of posts and replies and ego networks using social network analysis. Both visualisation and analysis were conducted using Python. Descriptive analysis of the networks (i.e., number of users, posts, and posting frequency) was carried out using the Pandas library and Networkx. Relevant demographic data and Parkinson's risk scores (expressed as an odds 1:x) were analysed using descriptive statistics. As the PD risk score is the inverse of the risk of developing PD, users with lower risk PD score have higher risk of developing PD. Regression analysis was conducted to estimate the relationships between risk scores (after log transformation) and network measures.

Results: 219 participants took part in a 4-month pilot forum embedded in the study website. 201 people connected in a large group, where most pairs of users could reach one another either directly or indirectly through other users. 59% (20/34) of discussions were spontaneously started by participants. Participation was asynchronous, with some individuals acting as 'brokers' between groups of discussions. As more participants joined the forum and connected to one another through online posts, distinct groups of connected users started to emerge. This pilot showed that a forum application within the cohort web platform was feasible and acceptable, and fostered digital social interaction. Matching with PREDICT-PD previously collected data was feasible, showing potential for future analyses correlating online network characteristics with risk of PD over time, as well as testing digital social engagement as an intervention to modify the risk of developing neurodegenerative diseases.

Conclusions: These results show the potential of an online forum to investigate the role of social capital as a catalyst for moderating the risk of PD. Clinical Trial: n/a

(JMIR Preprints 18/08/2023:51977)

DOI: <https://doi.org/10.2196/preprints.51977>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

✓ **Only make the preprint title and abstract visible.**

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [http://www.jmir.org/](#)

Original Manuscript

A pilot study of the role of social capital in a Parkinson's disease risk cohort: feasibility of a web application to enable online social interactions

Li X¹, Gill A², Panzarasa P¹, Bestwick JP², Schrag A⁴, Noyce AJ², De Simoni A⁴

Affiliations:

1 School of Business and Management, Queen Mary University of London

2. Preventive Neurology Unit, Wolfson Institute of Population Health, QMUL

3 Department of Clinical and Movement Neurosciences, Institute of Neurology, University College London

4 Centre for Primary Care, Wolfson Institute of Population Health, Queen Mary University of London

Keywords: Pilot studies, network analysis, Parkinson's disease, risk factors

Total number of words 1,549

Abstract

Background

There is evidence that social interaction has an inverse association with the development of neurodegenerative diseases. PREDICT-PD is an online UK cohort study that stratifies participants for risk of future Parkinson's disease (PD).

Objective

This study aims to explore the methodological approach and feasibility of assessing the digital social characteristics of people at risk of developing PD and their social capital within the PREDICT-PD platform, making hypotheses about the relationship between online social engagement and potential predictive risk indicators of Parkinson's disease

Methods

A web application was built to enable social interaction via the PREDICT-PD portal. Feedback from existing members of the cohort was sought and informed the design of the pilot. Dedicated staff used weekly engagement activities, consisting of PD-related research, facts, and queries to stimulate discussion. Data were collected by the hosting platform. We examined the pattern of connections generated over time through the cumulative number of posts and replies and ego networks using social network analysis. We employed network metrics to describe the bonding, bridging, and linking social capital among participants on the platform. Both visualisation and analysis were conducted using Python. Descriptive analysis of the networks (i.e., number of users, posts, and posting frequency) was carried out using the Pandas library and Networkx. Relevant demographic data and Parkinson's risk scores (expressed as an odds 1:x) were analysed using descriptive statistics. As the PD risk score is the inverse of the risk of developing PD, users with lower risk PD score have higher risk of developing PD. Regression analysis was conducted to estimate the relationships between risk scores (after log transformation) and network measures.

Results

219 participants took part in a 4-month pilot forum embedded in the study website. 201 people (40% male and 57% female) connected in a large group, where most pairs of users could reach one another either directly or indirectly through other users. 59% (20/34) of discussions were spontaneously started by participants. Participation was asynchronous, with some individuals acting as 'brokers' between groups of discussions. As more participants joined the forum and connected to one another through online posts, distinct groups of connected users started to emerge. This pilot showed that a forum application within the cohort web platform was feasible and acceptable, and fostered digital social interaction. Matching with PREDICT-PD previously collected data was feasible, showing

potential for future analyses correlating online network characteristics with risk of PD over time, as well as testing digital social engagement as an intervention to modify the risk of developing neurodegenerative diseases.

Conclusions

The results from the pilot suggest that an online forum can serve as an intervention to enhance social connectedness and investigate whether patterns of online engagement can impact the risk of developing PD through long-term follow-up. This highlights the potential of leveraging online platforms to study the role of social capital in moderating PD risk and underscores the feasibility of such approaches in future research or interventions.

Introduction

Parkinson's disease (PD) is the second most common neurodegenerative disease after Alzheimer's disease [1]. It is associated with disability, loss of independence, carer burden, and premature death [2]. Life expectancy is reduced for all patients, especially in individuals with early onset PD [2]. More than 75% have dementia after eight years from diagnosis [3]. PD onset age is typically 60-65 years [4], with early onset being considered between 21 and 50 years [5].

Treatments and interventions for PD are more likely to be effective if they are applied before the current point of diagnosis, by which >50% brain cells have already been lost [6]. While neuroprotective agents for neurodegenerative disease have shown promise in experimental studies [7,8], clinical research on prevention of PD strategies in 'at-risk' individuals is relatively understudied [9].

Studies have suggested that lifestyle measures such as reducing stress and increasing social interaction may be protective factors for neurodegenerative diseases [10,11]. Social capital refers to any type of benefit that individuals or groups can achieve through their underlying social relationships [12]. Changes in social interactions (or variations in social capital) can be associated with variations in risk factor for early death, poor mental health, and the development of dementia [13,14]. There are three types of social capital, known as bonding, bridging, and linking [15]. *Bonding Social Capital* is characterised by intra-group connections among members of a network who are similar [16], often in terms of sociodemographic characteristics like family, age, or interests. *Bridging Social Capital* derives from ties across diverse otherwise disconnected groups, including individuals from varied communities or organisations [17]. *Linking Social Capital* bridges the gap between different power tiers, typically linking community members with influential individuals or

institutions with substantial resources [18]. Several studies have investigated the association between social capital and health-related outcomes, including health care accessibility and the provision of health education services [19,20].

PREDICT-PD is a web-based study of the UK general population to identify a group at higher risk of PD, using an algorithm that estimates the risk of developing PD from information that can be collected using online tools, before the symptoms appear [21]. More than 7,700 people across the UK were enrolled in PREDICT-PD when undertaking annual tests online, with in-person assessments, such as brain scans, happening in a smaller proportion. Traditional methods of evaluating social interactions have predominantly depended on self-administered surveys by participants [22]. By leveraging on the PREDICT-PD platform, we can directly observe participants' social engagement through their forum posts and responses, enabling the analysis of the social involvement of a broader participant group. While several studies have shed light on the impact of social capital and social networks on the well-being of elderly individuals with chronic non-communicable diseases [23] and those at risk for dementia [24], there is a limited body of research focusing specifically on the association between social capital and the risk of Parkinson's disease. A recent study found that social isolation was associated with greater patient-reported PD severity and lower quality of life. Participants who reported having an extensive circle of friends had 21% fewer symptoms compared to those with limited or no social connections [25]. This finding underscores a potential link between social capital and PD severity.

In the current pilot study, we aimed to assess the digital social characteristics of people at risk of developing PD and their social capital within the PREDICT-PD platform. We employed various network metrics to describe the bonding, bridging, and linking social capital among participants on the platform [20]. Our objective is to make and assess hypotheses about the relationship between online social engagement and potential predictive risk indicators of Parkinson's disease, specifically in the context of social capital [20]. The insights gained from this pilot study can inform future research or interventions. Specifically, beyond the main aim to test feasibility and acceptability of this approach, we set to test the following research questions:

- i) Whether the breadth of an individual's social circle, as represented by degree centrality, is positively or negatively correlated with the risk of PD.
- ii) Whether the proximity of an individual to others in the network, as represented by closeness centrality, is positively or inversely correlated with the risk of PD.
- iii) Whether a high local clustering coefficient, indicative of social embeddedness in a

closely-knit community, is associated with an increased or decreased risk of PD.

Methods

This study used a web application built to enable social interaction via the PREDICT-PD portal. The online forum was made public in April 2021 for a period of four months. Feedback on the web application was collated from five PREDICT-PD cohort 'champions', who were representatives of the PREDICT-PD participant focus group. Champions are participants of the PREDICT-PD study who work in partnership with the PREDICT-PD team through regular meetings, to ensure that the views of patients, carers and the public are considered when decisions are made. They have been taking part in the study for up to 13 years.

This feedback informed our plan for launching the platform and strategy for engagement, i.e., appointing dedicated staff to ensure the platform was successful and stayed focused on issues relevant to PREDICT-PD participants, with weekly engagement activities, consisting of PD-related research, facts, and queries to stimulate discussion.

Data were collected by the hosting platform. PREDICT-PD cohort participants were sent an email about the forum and imminent pilot study. The option 'Forum' appeared on the left-hand menu bar and anybody could access the Terms and Conditions of engagement and register a username. Registered users could choose to either write posts publicly or send private posts. In the latter case, posts were shared between two users only, whereas when posts were written publicly, a large number of users could become connected through threads of posts. Only posts that were shared publicly were collected and analysed. For this study, user identifiers (IDs) were anonymised, and no demographic information was collected. The data sets included posts and their metadata (i.e., the anonymised user ID numbers), user roles (e.g., user, administrator, or moderator), date of posting, the hierarchical level of the post within the corresponding thread, and the dates in which the users joined and left the community. The community was moderated by members of the research team.

Data Analysis

We looked at the number of users, the number of posts and connections per user, and posting frequency. A connection (i.e., a tie, link, or edge) was established from one user to another when the former replied to a post submitted by the latter. We examined the pattern of connections generated over time through the cumulative number of posts and replies. We were interested not just in the number of posts and responses but in who responded to whom, and when. To this end, we used social network analysis [26] to visualise and study the structure of the relationships among users.

Both visualisation and analysis were conducted using Python. The network analysis was carried out through additional custom computer code in Python. Descriptive analysis of the networks (i.e., number of users, posts, and posting frequency) was carried out using the Pandas library, an open-source library providing data structures and analysis tools for the Python programming language, and Networkx, another library facilitating analysis and calculations of network metrics. Taking each user as a node and replies as links, we constructed a user network. We focused on the largest connected component of this network and used several measures including:

‘(1) degree, defined as the number of links connected to the node. A high degree suggests that the user is directly connected to many other users, indicating a higher volume of social interactions in the OHC. In the context of social capital, it represents the quantity of direct interactions or the breadth of a users’ social interactions.

(2) betweenness centrality, which measures the number of times a node lies on the shortest path (i.e., a path between two nodes that has the smallest number of links) between other nodes. It indicates how often an individual acts as a ‘bridge’ or intermediary between others within the network, connecting different parts or groups, thus representing a form of *Bridging Social Capital*. An individual with high betweenness centrality is key in facilitating the flow of information and resources across diverse segments of the network, thus enhancing the network's overall diversity and integration.

(3) closeness centrality, which measures the average shortest distance (i.e., number of links in the shortest path) from each node to all other nodes in the network (thus indicating how close a node is to all other nodes in the network). A higher closeness centrality means that a user can reach many other users in fewer steps. In terms of social capital, this reflects how efficiently an individual can access or disseminate information within the community. An individual with high closeness centrality might be better positioned to gather diverse insights or support from the community.

We also constructed ego-centred networks for each user and measured

(4) the local clustering coefficient, which quantifies the degree to which the user’s neighbours tend to cluster together. Local clustering coefficient assesses how interconnected an individual’s neighbours are. A high value suggests that many of the user's contacts also interact with each other, creating a tightly-knit community. In the context of social capital, it represents the intensity of social interactions within a user's immediate circle, which aligns with the concept of *Bonding Social Capital*. A high clustering coefficient indicates density among a group, which might have implications for emotional and informational support [20].

(5) effective size, calculated by the number of neighbours that the user has, minus the average

number of ties that each of the neighbours has to other neighbours. Effective size quantifies the degree to which the user's ego-centered network is rich in structural holes, and thus provides the node with bridging potential and brokerage opportunities. Effective size can also be used as a measure of *Bridging Social Capital*. A higher value indicates more structural holes in the user's ego-centered network and a higher bridging ability.

By using regression analysis, we aim to investigate the association between each of the above forms of participants' social capital and their PD risks.'

Relevant demographic data and Parkinson's risk scores (expressed as an odds 1:x) were analysed using descriptive statistics. Regression analysis was conducted to estimate the relationships between PD risk scores (after log transformation) and network measures using statsmodels (a Python module that provides classes and functions for the estimation of statistical models).

Ethics

This study received ethical approval from Queen Square Ethics committee (REC reference: 10/H0716/85), IRAS project ID: 20769

Informed consent for participation, collection and analysis of engagement data was sought prior to users joining the online community through the online platform. Usernames were replaced with numerical codes and analyses performed anonymized and aggregated data. The PREDICT-PD website uses encrypted data capture and the AWS server is ISO270001 accredited. Profile information and research data are stored separately.

Results

Participants

219 participants from a cohort of 7,700 took part in the pilot forum, including 218 users and one administrator who posted a blog weekly. For the 218 users, age could be matched for 187/218, gender for 193/218, while PD risk score for 187/218 users with data previously collected from the PREDICT-PD database. Age and gender of forum participants were consistent with PREDICT-PD cohort members [21], with a slight predominance of females. Over four months, through posting to each other and to the administrator, 200/218 (91.7%) users became part of a connected group (the largest component) (see Figure 1). The largest community formed around the admin staff posting weekly (Figure 1, purple colour), though smaller communities were started by users themselves (Figure 1, green, pink, orange, green, blue, and black colour). Therefore, communities formed independently from the action of admin staff.

Figure 1 here

The characteristics of users in the largest component (200/218) have been summarised Table 1.

Table 1 here

Figure 2 illustrates the network of posts, where each node represents an individual post. Posts written by admin are in green and posts by users in red. The size of each circle is proportional to the number of replies received. Participants wrote a total of 364 posts, divided into 34 threads. 14/34 (41%) were started by the administrator. 20/34 (59%) threads were spontaneously started by users, attracting a total of 133 posts, written by 99 individual users (see Figure 2). Since the start of the pilot, users created small communities of discussion, independently from the administrator. Weekly blogs were effective in creating conversations among users.

Figure 2 here

To show the development of discussion in the pilot forum, we measured the time gap between the original post and the replies in each thread. The frequency of replies with corresponding time gaps are plotted in Figure 3. We found that participation was asynchronous (as is typical in online communities), with users replying to posts at different times. Most replies are written within a week after the original posts. The mean time gap between reply and the original posts is 5.67 days, with a standard deviation of 11.28 days. However, some users also replied to older posts with a maximum time gap of 82 days. This suggests that users were checking the forum and were motivated to participate in ongoing conversations. This pattern shows that the forum served not only as a platform for immediate exchange of information but also as a repository of knowledge and ideas that users revisited over time. These findings suggest a dual nature of the forum: as a space for timely interaction and as a valuable archive of discussions that continue to engage users over extended periods.

Figure 3 here

As network measures sometimes correlate with each other, we checked the correlations among the measures listed in Table 1. According to the correlation matrix in Table 2, betweenness centrality and effective size are significantly correlated with degree centrality. To avoid multicollinearity, we did not include the two measures in regression models.

Table 2 here

Regression results are given in Table 3. In model 1, we regressed the risk scores on the control variables age and gender. In model 2, 3 and 4, we added three network measures (i.e., degree centrality, closeness centrality and local clustering coefficient) separately in the regressions. In model

5, we added all three measures. The results showed ~~correlation~~ that the association between Parkinson's risk and degree centrality (i.e., high number of direct connections) is significant (see Table 3, model 2 and 5, $P < 0.1$). This may be related to the nature of the forum leading participants with lower PD risk scores (i.e., higher risks of developing PD) to reach out for help and suggestions, which results in more discussion and communication with other participants.

The non-significant results on closeness centrality and local clustering coefficient offer preliminary insights on the direction of their associations with Parkinson's risk. Local clustering coefficients tend to negatively correlate with the risk score, meaning that users with lower risk scores (i.e., higher risks of developing PD) were more likely to embed themselves into tightly-knit communities. This is possibly because participants with higher PD risks may share similar experiences and show interest in similar topics within the forum. On the opposite, users with higher scores (i.e., lower risks of developing PD) tended to display higher closeness centrality. This could suggest that users with closer access to many other members within the forum (i.e., those who could reach out to a large portion of the forum and receive information very quickly without necessarily being the hubs) are less likely to have higher risk of developing PD.

We performed tests for multicollinearity with variance inflation factor (VIF). Since each of the VIF values for the predictor variables in the model are close to 1, multicollinearity is not a problem in the model.

Table 3 here

This finding suggests that users' connections to others in the online community can be associated with the risk of developing PD, thus showing the potential of correlating network properties of users in online communities to risk of diseases.

Discussion

This 4-month pilot showed that a forum application within the PREDICT-PD web platform was feasible and acceptable, and fostered digital social interaction. Participants' activity over time showed that the forum was used as a repository of knowledge and ideas that users revisited over time.

Forum participants' characteristics were consistent with PREDICT-PD cohort members. Matching with PREDICT-PD previously collected data was feasible, showing potential for future analyses correlating online network characteristics with risk of PD over time, as well as testing digital social engagement as an intervention to modify the risk of developing neurodegenerative diseases.

Engagement with the forum was found to take place independently from the weekly engagement activities from admin staff. As more participants joined the forum and connected to one another through online posts, distinct groups of connected users started to emerge. These groups have fundamental implications for the effectiveness of processes of network dynamics such as information diffusion [27]. In the four-month pilot, the community underwent the formation of the 'largest component' of connected users. It is well known that components are critical for information diffusion [28,29]. Without a large, connected component, users would be members of small isolated islands, and information would be unable to flow from one island to another.

Based on the regression results we hypothesize that users with lower risk scores (i.e., higher risk of developing PD) may be those reaching out for help and suggestions within the forum, with a tendency of displaying higher degree centrality. Conversely, users who keep being active in the central community (i.e., with higher closeness centrality) tend to have higher risk scores (i.e., lower risk of developing PD). In this exploratory analysis of the associations between network characteristics and PD risk some of the findings did not reach statistical significance. This outcome, while representing a limitation, is not unexpected given the pilot nature of the study, designed primarily for feasibility and methodological exploration rather than definitive causal inference. These preliminary findings serve as an important stepping stone, highlighting the need for larger, more comprehensive studies. Future research should build on this foundational work, employing larger sample sizes and longer durations to adequately power the investigation of the potential impact of network characteristics on PD risk.

These results suggest that the online forum could serve as a platform for targeted interventions for individuals with higher PD risks to spontaneously gather and engage with one another. By monitoring network measures linked to their engagement, we might be able to identify users with higher PD risk and provide them with tailored resources or direct interventions.

This pilot study adopted an observational research approach. An important limitation inherent to such studies is the possibility of reverse causation influencing the detected associations, making it essential to consider the duration of follow-up. Specifically, we observed that local clustering coefficients tend to negatively correlate with the risk score, suggesting that individuals at higher risk of developing PD might preferentially engage in tightly-knit communities. This could reflect a

tendency among participants with higher PD risks to seek out and share experiences with peers facing similar challenges. Conversely, it is also possible that the formation of such tightly-knit communities and the homogeneous nature of the information exchanged within them could, if predominantly negative, contribute to increased stress or negative emotional states, potentially elevating the risk of PD. This dual interpretation underscores the complexity of the reverse causation issue, as it illustrates how the detected associations could be interpreted differently. A deeper understanding can be gained by analysing patterns of engagement in social activities over time and correlating them with the risk of PD. This can be facilitated using data from studies with repeated assessments of social participation and associated risk metrics, which will be the next step of this research.

Future research based on a larger data set would allow for more robust statistical analysis, potentially confirming the trends observed in this pilot study, and would allow us to investigate whether active online participation can be associated with modification of PD risk score and improved health outcomes, or whether it rather reflects a self-selection bias prompting individuals with higher risks to engage in forum discussions more intensely. Longitudinal studies could also determine whether changes in network centrality measures correlate with changes in users' health status in the context of neurodegenerative diseases. Such studies will aim to employ advanced statistical models like mixed-effects models or growth curve analysis to discern patterns of change and causality. Specifically, it would be insightful to investigate how early changes in social network dynamics precede the onset of PD symptoms, and whether increasing social connectivity can mitigate risk. Besides, by using the longitudinal data, temporal centrality measures can be calculated, and dynamic network analysis can significantly enhance the depth and breadth of insights, especially in tracking changes in social engagement and their impact on PD risk.

Additionally, qualitative analysis of the content of communications could provide deeper insights into the types of support and information being exchanged [20] and how these might relate to PD risk and management. Moreover, qualitative methodologies could enable us to uncover factors contributing to users' engagement with the forum, particularly when taking place independently of the admin staff's activities, e.g., specific user-driven initiatives or topics that lead to increased participation. Understanding these dynamics could provide insights for future digital social interventions targeted to people at risk of developing PD.

The results presented here show potential of an online forum as an intervention to trigger social connectedness and to investigate whether patterns of online engagement [20] have a causal impact upon the risk of developing PD through long-term follow-up.

Funding

The study was funded through a pump priming grant by Barts Charity (reference MGU04579).

XL, PP and ADS were partly funded by a NIHR Programme Grant for Applied Research (NIHR202037). Views are those of the authors and not necessarily those of the National Health Service, NIHR, or the Department of Health and Social Care.

Ethics

This study received ethical approval from Queen Square Ethics committee (REC reference: 10/H0716/85), IRAS project ID: 20769

Informed consent for participation, collection and analysis of engagement data was sought prior to users joining the online community through the online platform.

Acknowledgements

We would like to thank all the patient and public members involved with the research at PREDICT-PD.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of interest

None.




Figure 1. Cumulative network of individual participants in the largest component across the time span analysed. The size of the node correlates with the number of posts written by that user. Colours represent different groups of discussions.

Sample characteristics	N	Mean (range)
Number of posts		
Total number of posts (admin's posts excluded)	348	
Number of posts on the community /user		1.60 (1-6)
Number of users in the largest component	200	
Age	172	67.55
Unknown age (missing from matched PREDICT-PD data)	18	(25.84-83.60)
Gender		
Male	80	
Female	113	
Unknown (missing matched PREDICT-PD data)	7	
Risk of Parkinson's		
Risk Score (log transformed)	172	4.21 (IQR 4.29)
Network Characteristic*		
Degree	200	1.93 (1-18)
Closeness centrality	200	0.331 (0.005-0.417)
Betweenness centrality	200	0.006 (0-0.125)
Ego Network Characteristics		
Local clustering coefficient	200	0.082 (0-1)
Effective size	200	1.79 (1-17)

Table 1. Characteristics of participants in the largest component.

* analysis performed on 200/218 users connected in the largest component.




Figure 2. Representation of communities. Cumulative network of posts across the time span analysed. Circles represent individual posts: in green posts written by admin and in red posts by users.




Figure 3. Distribution of replies over time (days) after the first post in the corresponding thread.



	1	2	3	4	5	6	7	8
1. Risk score	1.0							
2. Age	-0.42***	1.0						
3. Gender	-0.32***	0.15	1.0					
4. Degree	-0.14	0.05	0.1	1.0				
5. Closeness	0.03	0	-0.09	0.11	1.0			
6. Betweenness	-0.21**	0.05	0.14	0.84***	0.12	1.0		
7. Clustering coefficient	-0.14	0.11	-0.07	0.13	0.24* *	-0.03	1.0	
8. Effective size	-0.13	0.03	0.12	0.98***	0.06	0.85***	-0.04	1.0
* p < 0.05, ** p < 0.01, ***p < 0.001								

Table 2. Correlation matrix.

	(1)	(2)	(3)	(4)	(5)
Intercept	15.64*** (1.56)	15.84*** (1.54)	15.52*** (1.80)	15.40*** (1.58)	14.90*** (1.75)
Age	-0.16*** (0.02)	-0.16*** (0.02)	-0.16*** (0.02)	-0.15*** (0.02)	-0.15*** (0.02)
Gender	-1.65*** (0.44)	-1.59*** (0.44)	-1.64*** (0.44)	-1.71*** (0.43)	-1.63*** (0.44)
Degree		-0.16* (0.08)			-0.14* (0.09)
Closeness			0.36 (2.98)		2.04 (2.96)
Clustering coefficient				-1.47 (1.03)	-1.46 (1.05)
Observations	172	172	172	172	172
Robust standard errors in parentheses. * p < 0.1, ** p < 0.05, ***p < 0.01					

Table 3. Logistic regression models showing the predictors of Parkinson's disease risk.

This analysis includes the 172/200 forum users in the largest component, whose data about age, gender, degree, closeness, clustering coefficient could be matched from the PREDICT-PD database.

References

1. Mhyre TR, Boyd JT, Hamill RW, Maguire-Zeiss KA. Parkinson's disease. Subcell Biochem. 2012;65:389-455. Doi: 10.1007/978-94-007-5416-4_16. PMID: 23225012; PMCID: PMC4372387.
2. Ishihara LS, Cheesbrough A, Brayne C, Schrag A. Estimated life expectancy of Parkinson's patients compared with the UK population. J Neurol Neurosurg Psychiatry 2007;78:1304-1309. Doi: 10.1136/jnnp.2006.100107
3. Aarsland D, Andersen K, Larsen JP, Lolk A. Prevalence and Characteristics of Dementia in Parkinson Disease: An 8-Year Prospective Study. Arch Neurol 2003;60:387-392. Doi: 10.1001/archneur.60.3.387

4. Ascherio A, Schwarzschild MA. The epidemiology of Parkinson's disease: risk factors and prevention. *Lancet Neurol.* 2016;15(12):1257-1272. doi: 10.1016/S1474-4422(16)30230-7. Epub 2016 Oct 11. PMID: 27751556.
5. Quinn N, Critchley P, Marsden CD. Young onset Parkinson's disease. *Mov Disord.* 1987;2(2):73-91. doi: 10.1002/mds.870020201. PMID: 3504266.
6. Fearnley JM, Lees AJ. Ageing and Parkinson's disease: substantia nigra regional selectivity. *Brain* 1991;114:2283-301. Doi: 10.1093/brain/114.5.2283
7. Güzelad, Ö., Özkan, A., Parlak, H. et al. Protective mechanism of Syringic acid in an experimental model of Parkinson's disease. *Metab Brain Dis* 36, 1003–1014 (2021). Doi: 10.1007/s11011-021-00704-9
8. Ogut, E., Armagan, K. & Gül, Z. The role of syringic acid as a neuroprotective agent for neurodegenerative disorders and future expectations. *Metab Brain Dis* 37, 859–880 (2022). Doi: 10.1007/s11011-022-00960-3
9. Noyce AJ, Lees AJ, Schrag AE. The prediagnostic phase of Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2016;87:871-8. Doi: 10.1136/jnnp-2015-311890
10. Livingston G, Sommerlad A, Orgeta V, Costafreda SG, Huntley J, Ames D, et al. Dementia prevention, intervention, and care. *Lancet* 2017;390:2673-2734. Doi: 10.1016/S0140-6736(20)30367-6
11. Guarnera J, Yuen E, Macpherson H. The Impact of Loneliness and Social Isolation on Cognitive Aging: A Narrative Review. *J Alzheimers Dis Rep.* 2023 Jun 29;7(1):699-714. Doi: 10.3233/ADR-230011. PMID: 37483321; PMCID: PMC10357115.
12. Lin, N. Social capital. A theory of social structure and action. 2001 Cambridge: Cambridge University Press. Doi: 10.1016/S0140-6736(17)31363-6
13. Allen C, Vassilev I, Kennedy A, Rogers A. Long-term condition self-Management support in online communities: A meta-synthesis of qualitative papers. *J Med Internet Res* 2016;18: , e61 Doi:10.2196/jmir.5260

14. Hobbs WR, Burke M, Christakis NA, Fowler JH. Online social integration is associated with reduced mortality risk. *Proc Natl Acad Sci U S A* 2016;113:12980-12984. Doi: 10.1073/pnas.1605554113
15. Derose KP, Varda DM. Social capital and health care access: a systematic review. *Medical care research and review: MCRR*. 2009;66(3):272. Doi: 10.1177/1077558708330428
16. Hawkins RL, Maurer K. Bonding, bridging and linking: How social capital operated in New Orleans following Hurricane Katrina. *British Journal of Social Work*. 2010;40(6):1777-93. Doi: 10.1093/bjsw/bcp087
17. Putnam RD. *Bowling alone: The collapse and revival of American community*. Simon and schuster; 2000. Doi: 10.1145/358916.361990. ISBN: 0743203046, 9780743203043
18. Szreter S, Woolcock M. Health by association? Social capital, social theory, and the political economy of public health. *International journal of epidemiology*. 2004;33(4):650-67. Doi: 10.1093/ije/dyh013.
19. Holtgrave DR, Crosby R. Is social capital a protective factor against obesity and diabetes? Findings from an exploratory study. *Annals of epidemiology*. 2006 May 1;16(5):406-8. Doi: 10.1016/j.annepidem.2005.04.017
20. Panzarasa P, Griffiths CJ, Sastry N, De Simoni A. Social medical capital: how patients and caregivers can benefit from online social interactions. *J Med Internet Res* 2020; 22:e16337 Doi: 10.2196/16337
21. Noyce AJ, R'Bibo L, Peress L, Bestwick JP, Adams-Carr KL, Mencacci NE, et al. PREDICT-PD: An online approach to prospectively identify risk indicators of Parkinson's disease. *Mov Disord*. 2017;32(2):219-226. Doi: 10.1002/mds.26898
22. Coleman JS. *Foundations of social theory*. Harvard university press; 1994. ISBN: 9780674312265
23. Christian AK, Sanuade OA, Okyere MA, Adjaye-Gbewonyo K. Social capital is associated with improved subjective well-being of older adults with chronic non-communicable disease in six low- and middle-income countries. *Globalization and Health*. 2020 ;16(1):1-1. Doi: 10.1186/s12992-019-0538-y

24. Abbott KM, Pachucki MC. Associations between social network characteristics, cognitive function, and quality of life among residents in a dementia special care unit: A pilot study. *Dementia*. 2017 Nov;16(8):1004-19. Doi: 10.1177/147130121663090
25. Subramanian I, Farahnik J, Mischley LK. Synergy of pandemics-social isolation is associated with worsened Parkinson severity and quality of life. *npj Parkinson's Disease*. 2020 Oct 8;6(1):28. Doi: 10.1038/s41531-020-00128-9
26. Wasserman S, Faust K, Granovetter M. *Social Network Analysis: Methods and Applications*. 1994 Cambridge: Cambridge University Press. ISBN: 0521387078, 9780521387071
27. Newman M. Newman. In: *Networks: An Introduction*. 2010 Oxford: Oxford University Press, USA. Doi: 10.1093/acprof:oso/9780199206650.001.0001. ISBN: 9780199206650
28. Yagan O, Qian D, Zhang J, Cochran D. Conjoining Speeds up Information Diffusion in Overlaying Social-Physical Networks. *IEEE J. Select. Areas Commun* 2013;31:1038-1048. Doi: 10.1109/JSAC.2013.130606
29. Lambiotte R, Panzarasa P. Communities, knowledge creation, and information diffusion. *Journal of Informetrics* 2009;3:180-190. Doi: 10.1016/j.joi.2009.03.007

Supplementary Files

Untitled.

URL: <http://asset.jmir.pub/assets/f1f0d2734b91a05b77f6c36b6171d5a7.docx>