

Development of an Electronic Health Record Self-Referral Tool for Lung Cancer Screening: A Pilot Study

Garrett Smith Stang, Nichole T Tanner, Ashley Hatch, Jakarri Godbolt, Benjamin A Toll, Alana M Rojewski

Submitted to: JMIR Formative Research
on: September 27, 2023

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 16

..... 16

Figures 17

 Figure 1..... 18

 Figure 2..... 19

 Figure 3..... 20

Multimedia Appendixes 21

 Multimedia Appendix 1..... 22

 Multimedia Appendix 2..... 22

 Multimedia Appendix 3..... 22

Development of an Electronic Health Record Self-Referral Tool for Lung Cancer Screening: A Pilot Study

Garrett Smith Stang^{1,2} BSc, MPH; Nichole T Tanner¹ MD, MSCR; Ashley Hatch¹ MPH; Jakarri Godbolt¹ MPH; Benjamin A Toll¹ PhD; Alana M Rojewski¹ PhD

¹Medical University of South Carolina Charleston US

²Brown University School of Public Health Providence US

Corresponding Author:

Garrett Smith Stang BSc, MPH
Brown University School of Public Health
121 S Main St
Providence
US

Abstract

Background: Approximately 14 million individuals in the U.S. are eligible for lung cancer screening (LCS), but only 5.8% completed screening in 2021. Given the low uptake despite the potential great health benefit of LCS, interventions aimed at increasing uptake are warranted.

Objective: This study sought to develop and pilot an electronic health record (EHR) patient-facing self-referral tool to an established LCS program in an academic medical center.

Methods: Guided by constructs of the Health Belief Model associated with LCS uptake (e.g., knowledge, self-efficacy), formative development of an EHR-delivered engagement message, infographic, and self-referring survey was conducted. The survey submits eligible self-reported patient information to a scheduler for the LCS program. The materials were pretested using an interviewer-administered mixed-methods survey captured through venue-day-time sampling in five network-affiliated pulmonology clinics. Materials were then integrated into the secure patient messaging feature in the EHR system. Next, a one-group posttest quality improvement pilot test was conducted.

Results: Seventeen individuals presenting for lung screening shared-decision visits completed the pretest survey. More than half were newly referred for LCS (n=10, 58.8%), and the remaining were returning patients. When asked if they would use a self-referring tool through their EHR messaging portal, 94.12% (n=16) reported yes. Fifteen participants provided oral feedback that led to refinement in the tool and infographic prior to pilot testing. When the initial application of the tool was sent to a convenience sample of 150 random patients, 13.3% (n=20) opened the self-referring survey. Of the 20, 45.0% (n=9) were eligible for LCS based on self-reported smoking data. Three self-referring individuals scheduled a LCS.

Conclusions: Pretest and initial application data suggest this tool is a positive stimulus to trigger the decision-making process to engage in self-referral process to LCS among eligible patients. This self-referral tool may increase the number of patients engaging in LCS and could also be used to aid in self-referral to other preventative health screenings.

(JMIR Preprints 27/09/2023:53159)

DOI: <https://doi.org/10.2196/preprints.53159>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.
Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/preprint/53159>, the full manuscript will be available to all users.



Original Manuscript

Development of an Electronic Health Record Self-Referral Tool for Lung Cancer Screening: A Pilot Study

Abstract

Background: Approximately 14 million individuals in the U.S. are eligible for lung cancer screening (LCS), but only 5.8% completed screening in 2021. Given the low uptake despite the potential great health benefit of LCS, interventions aimed at increasing uptake are warranted. The utilization of a patient-facing electronic health record (EHR) patient portal direct messaging tool offers a new opportunity to both engage eligible patients in preventative screening and provide a unique referral pathway for tobacco treatment.

Objective: This study sought to develop and pilot an electronic health record (EHR) patient-facing self-referral tool to an established LCS program in an academic medical center.

Methods: Guided by constructs of the Health Belief Model associated with LCS uptake (e.g., knowledge, self-efficacy), formative development of an EHR-delivered engagement message, infographic, and self-referring survey was conducted. The survey submits eligible self-reported patient information to a scheduler for the LCS program. The materials were pretested using an interviewer-administered mixed-methods survey captured through venue-day-time sampling in five network-affiliated pulmonology clinics. Materials were then integrated into the secure patient messaging feature in the EHR system. Next, a one-group posttest quality improvement pilot test was conducted.

Results: Seventeen individuals presenting for lung screening shared-decision visits completed the pretest survey. More than half were newly referred for LCS (n=10, 58.8%), and the remaining were returning patients. When asked if they would use a self-referring tool through their EHR messaging portal, 94.12% (n=16) reported yes. Fifteen participants provided oral feedback that led to refinement in the tool and infographic prior to pilot testing. When the initial application of the tool was sent to a convenience sample of 150 random patients, 13.3% (n=20) opened the self-referring survey. Of the 20, 45.0% (n=9) were eligible for LCS based on self-reported smoking data. Three self-referring individuals scheduled a LCS.

Conclusions: Pretest and initial application data suggest this tool is a positive stimulus to trigger the decision-making process to engage in self-referral process to LCS among eligible patients. This self-referral tool may increase the number of patients engaging in LCS and could also be used to aid in self-referral to other preventative health screenings. This tool has implications for clinical practice. Tobacco treatment clinical services or healthcare systems should consider utilizing EHR messaging for LCS self-referral. This approach may be cost-effective to improve LCS engagement and uptake. Additional referral pathways could be built into this EHR tool to not only refer patients who currently smoke to LCS but also simultaneously trigger a referral to clinical tobacco treatment.

Keywords: lung cancer screening (LCS); electronic health records (EHR); health belief model (HBM); self-refer; tobacco treatment

Introduction

Lung cancer is the second most common cancer worldwide and the leading cause of cancer-related death in the United States (U.S.), with higher mortality rates than breast, colon, and prostate cancer combined[1-3]. High-quality evidence demonstrates a 20-35% reduction in lung cancer mortality by screening those at high risk of lung cancer based on age and smoking history[4, 5]. Based on this, the U.S. Preventative Services Task Force (USPSTF) has given a grade B recommendation in favor of screening with low-dose computed tomography for individuals aged 50 to 80 years who currently smoke or have quit within the past 15 years with a minimum 20-pack-year smoking following a shared decision-making visit [1, 6]. Despite the benefits of screening and an estimated 14 million

individuals in the US eligible for lung cancer screening (LCS) [7], uptake of LCS has been low with only 12.8% having undergone screening [8]. Identifying approaches to increase LCS uptake would greatly benefit public health [9].

Barriers to LCS uptake have been identified at the system, provider, and patient levels. At the provider level, lack of familiarity with eligibility criteria, insufficient time or knowledge to conduct shared decision-making, skepticism about the benefits of screening, and familiarity with managing screen-detected findings have all been identified as barriers to uptake [10]. Patient-level barriers to LCS engagement include a need for more awareness of screening eligibility or programs, cost concerns, insurance status, and challenges in accessing imaging sites [10, 11]. In addition, LCS is the first preventative screening to target poor health behavior (i.e., smoking), and this population is less likely to have a primary care provider or engage in other screening services [11]. Thus, reducing barriers to LCS among this high-risk population who may not be engaging in health promoting behaviors is needed to increase uptake.

The present study aimed to develop an intervention to target the individual behavioral factors associated with low LCS uptake. A significant contributor to poor uptake is poor knowledge of the benefits of screening (e.g., perceived benefit) and a lack of perceived susceptibility and severity of lung cancer among screening-eligible patients [12-16]. Guided by the Health Belief Model (HBM) and leveraging an electronic health record (EHR) patient portal direct messaging feature, the tool is a multilevel theory-based conceptual model that targets behavior change in LCS uptake to engage potentially eligible patients in preventative screening [17, 18].

Methods

Theory-Guided Development

The HBM guided the development of the tool components. Research has shown that constructs within the HBM are associated with LCS uptake [14]. The tool components were designed to target knowledge of screening eligibility, the perceived threat of lung cancer, perceived benefits of LCS, self-efficacy, and prompt a cue to action for LCS self-referral.

Self-referral Tool Design

The tool was designed to be integrated into the electronic health record (Epic Systems); Epic Systems is a third-party, cloud-based EHR [19]. We utilized the patient-facing secure messaging application (MyChart) for self-referral to the medical center's centralized lung cancer screening program. A centralized lung screening program employs a dedicated lung screening navigator and a team that verifies eligibility, conducts shared decision-making, initiates tobacco treatment, orders, schedules imaging, manages results, and ensures adherence to follow-up. The tool components include an initial engagement message, a questionnaire, and an infographic highlighting the danger of lung cancer and LCS eligibility.

Initial Engagement Message

The initial engagement message conveyed the threat of lung cancer, the benefit of being screened, and a cue to action to self-assess their LCS eligibility. The initial engagement message mapped to the HBM constructs is presented in Figure 1.

Questionnaire

The MyChart-delivered questionnaire presents patients with the opportunity to assess their eligibility for LCS and self-refer to the LCS scheduler. The questionnaire assesses the individual's self-reported smoking status, years smoked, and packs per day, which map to the three LCS eligibility criteria.

Questions and responses were written at a presumed accepted general health literacy level. Figure 2 displays the questionnaire and associated branching logic. The results of this survey would then be sent to the LCS scheduler directly as an acute care LCS referral. Once the referral is received, the LCS scheduler will confirm eligibility and contact the patient directly to schedule their LCS.

Infographic

An infographic was created to expand on the emphasized perceived threat of lung cancer established in the initial engagement message. We determined that including an infographic in the tool lineup would increase questionnaire completion efforts by allowing individuals to visualize relevant statistics in an easily digestible format. The infographic mapped to the HBM constructs is presented in Appendix 1

Self-Referral Tool Pretest

The tool components were evaluated using a venue-day-time sampling interviewer-administered survey to gather feedback for iterative tool refinement prior to implementation.

Study Setting and Participants

Tool evaluation was conducted by participants meeting eligibility for lung cancer screening being seen in clinic as a new referral for LCS or returning patient with an active Epic MyChart account. Over four weeks, predetermined days and times were selected for data collection in five separate pulmonology clinics in the tri-county region of South Carolina where this investigation was conducted (Berkley, Charleston, and Dorchester).

Study Procedures

Eligible patients were identified through manual EHR review. Following the completion of the scheduled appointment, the LCS provider invited the patient to participate in the study. Those who agreed to participate then met with and gave verbal informed consent to the clinical research assistant (CRA). The participant was presented with copies of the self-referral questionnaire and the infographic. Then, a semi-structured interview and survey were conducted to gauge their response to the tool components.

Data Collection and Analysis

The survey was a mixed methods instrument that collected quantitative and qualitative data through open-ended questions. The survey measured health literacy, usability, perceived susceptibility, and self-efficacy using a 5-point Likert scale (1 = Strongly Agree, 2 = Agree, 3 = Neutral, 4 = Disagree, and 5 = Strongly Disagree). The survey also evaluated their overall willingness to complete the self-referral survey (Yes, No, and Unsure) and likelihood of completing the self-referral questionnaire without the accompanying message and infographic (Yes - I would complete any questionnaire sent to me via MyChart, Yes - but I would prefer to get those too, No - but I would if I also got the letter and infographic, No - I would not complete the questionnaire in MyChart, and Unsure). The qualitative free-responses items allowed participants to provide additional feedback on each tool component. Non-identifiable demographic data were also collected from a manual EHR chart review. The CRA took field notes of observations and quotes. Standard descriptive statistics were conducted to summarize quantitative variables. These analyses included a mean and standard deviation for age and percentages for the remaining categorical variables. Qualitative components (field notes and free responses) were analyzed using a content, thematic approach via inductive coding.

Electronic Health Record Integration

Following pre-test analysis, the self-referral tool components were refined and integrated into the

Epic system for broader testing. The self-referral tool was integrated into the Epic system secure patient messaging feature (i.e., In Basket) as a General Questionnaire task item. The initial engagement message was formatted as a SmartText option; this created a fixed message. The only modifiable element in the message was the patient's full name in the greeting, which is indicated by a coded field notation. The field notation auto-generated the patient's full name from their chart when sent to the patient. The infographic and questionnaire are sent as attachments to the message. The self-report questionnaire results are returned (like a reply) to the Epic account that sent the initial message. The reply or results can be securely forwarded to other healthcare professionals with an Epic account, and the LCS scheduler can message the patient directly on MyChart to discuss eligibility and scheduling.

Self-referral Tool Pilot Test

A one-group pilot test was conducted to determine the uptake of LCS in response to the self-referral tool.

Study Setting and Participants

A convenience sample of medical record numbers (MRNs) was identified from another ongoing quality improvement study in the MUSC LCS program. The other study examined the uptake of LCS among women who had completed a mammogram in the past year. The dataset included patients who were (1) network-affiliated; (2) female at birth; (3) between 50-77 years old; (4) undergone a mammogram in the past year; (5) had no previous LCS care or history of lung cancer; (6) had an active Epic MyChart account.

Study Procedures

MRNs for the pilot were extracted through random sampling. The MRNs listed on the dataset were assigned a sequential number. The sequential number boundaries were set on a random sequence generator software, which produced a random order list. From the randomized list, the first 150 assigned numbers (and accompanying MRN) were selected to receive the self-referral tool.

The tool components (the message in Figure 1 linking to the questionnaire in Figure 2) were delivered to eligible patients through MyChart from a research-designated Epic account linked to a CRA. Once sent, patients would receive a notification that they had a new MyChart message and could access it through their secure Epic MyChart account; this service is accessible on a mobile app or through a desktop web browser (displayed in Appendix 2 and 3).

Data Collection and Analysis

Responses to the self-referral questionnaire were collected and stored in the Epic account of the CRA, including tracking if the patient opened the message. The CRA tracked the following (1) the status of the message (open vs. unopened), (2) the status of the questionnaire (completed vs. not completed), (3) completed questionnaire status (eligible vs. ineligible), and (4) outcomes (LCS status and LC diagnosis, if applicable). Percentages for categorical variables were analyzed.

Ethical Considerations

Given that this was a quality improvement study, formal patient consent was not obtained. Participants verbally consented to participate in the pretest portion of the study, during which the study tool was evaluated, and informal interviews were conducted. Participants did not receive financial remuneration for their participation. All data reported in this publication have been de-identified.

Results

Pretest Results

Participant Demographics

Seventeen patients completed the survey for tool evaluation in October 2022. More than half were new patients (n=10, 58.8%), and the remaining were returning patients or annual screeners (n=7, 41.2%) (Table 1). The majority were white (n=15, 88.2%), female (n=11, 65.7%), self-reported former smokers (n=11, 64.7%), and were referred to the LCS program from a network-affiliated primary care provider (n=14, 82.4%). The average age was 63.5 (SD=8.10), ranging from 52 to 77. Health insurance coverage among participants included Medicare (n=9, 52.9%) and private (n=8, 47.1%).

Table 1. Demographics and Characteristics of Pretest Participants Surveyed for Self-referral Tool Evaluation

Demographic	Total
Age (years)	
N	17
Mean (SD)	63.5 (8.10)
Min,Max	52,77
Sex at Birth (n)	
Female	11 (64.7%)
Male	6 (35.5%)
Race (n)	
Black or African American	2 (11.8%)
White	15 (88.2%)
Appointment Type (n)	
New Patient Referral	10 (58.8%)
Returning (annual) Screener	7 (41.2%)
Smoking Status (n)	
Current Smoker	6 (35.2%)
Former Smoker	11 (64.7%)
Health Insurance Coverage (n)	
Medicare	9 (52.9%)
Private	8 (47.1%)
Referral Pathway to LCS Clinical Care Program (n)	
Network Affiliated PCP	14 (82.4%)
Network Affiliated non-PCP	1 (5.9%)
Self-Referral	2 (11.8%)

Questionnaire

When surveyed about their understanding of the health topic in the message, 100% reported Strongly Agree/Agree. When surveyed about their self-efficacy to complete the survey, 100% reported Strongly Agree/Agree with the easiness of completion.

Engagement Message

When surveyed about their understanding of the health topic in the message, 100% reported Strongly Agree/Agree. When surveyed about their understanding of the threat of lung cancer, 100% reported Strongly Agree/Agree. When surveyed about their understanding of the benefit of LCS, 100% reported Strongly Agree/Agree. Finally, when surveyed about their self-efficacy to take the LCS survey, 100% reported Strongly Agree/Agree.

Infographic

When surveyed about their understanding of the health topic in the infographic, 100% reported Strongly Agree/Agree. When surveyed about their perceived threat to lung cancer, 94.12% (N=16) reported Strongly Agree/Agree, and 5.88% (N=1) reported Neutral. When surveyed about their understanding of the benefit of LCS, 100% reported Strongly Agree/Agree. Finally, we surveyed their self-efficacy to scan a QR code to access additional resources to learn more about lung cancer and LCS, 82.35% (N=14) reported Strongly Agree/Agree, and 17.65% (N=3) reported Strongly Disagree.

Likelihood of Survey Completion

When asked if they would complete the self-referral survey sent via MyChart, the majority, or 94.12% (N=16), responded Yes, and 5.88% (N=1) reported Unsure. When asked if they would still fill it out if they did not receive the additional introductory message and infographic, the majority, 70.59% (N=12), reported that they would still complete the survey even without the additional materials; 41.18% (N=7) indicated they would prefer to receive the additional materials, and 29.41% (N=5) indicated they would complete any survey (i.e., not health screening related) sent to them in MyChart. 17.65% (N=3) indicated they were unsure, while the remaining 11.76% (N=2) indicated they would only complete the questionnaire if they received the additional materials.

Key Themes

Fifteen participants provided oral feedback when gauged about their perceptions of the tool during the pretest survey. The most common themes that emerged from the data were poor self-efficacy in utilizing the QR code, high self-efficacy in completing the survey, and heightened appreciation for information on LCS and the self-referral survey.

Pilot Test Results

Of the 150 patients sampled, 48.0% (n=72) opened the message. Of the 72 that opened the message, 20 completed the survey (27.78% completion rate). Based on the initial sample size, there was an overall response rate of 13.33%.

Of the 20 that completed the survey, 45.0% (n=9) self-reported LCS eligibility. Following referral to the LCS Navigator, who completed an additional manual EHR review prior to scheduling, determined that 6 of the self-referring individuals were ineligible. The remaining 3 self-referring individuals were scheduled for LCS. At the time of analysis, 2 had attended an LCS-shared decision-making appointment and underwent an LDCT. The findings of the two completed LDCTs were categorized into LUNG RADS I and LUNG RADS III.

A manual EHR review was conducted of the 52 individuals that did not complete the survey after opening the message. Of those patients, 71.15% (n=37) had no previous record of LDCT or established pulmonary care, 23.07% (n=12) had previously undergone LDCT and established pulmonary care, 0.02% (n=1) had contacted the LCS Program Navigator to schedule screening after opening the message, and 0.04% (n=2) had self-referred to pulmonary care (outside of the LCS care program) after opening the message.

A consort diagram of each patient's phase in the tool application is displayed in Figure 3.

Discussion

This formative process created an electronic health record (EHR) self-referral tool for an established LCS program. The pre-implementation evaluation of the tool indicates that the language, phrasing, and visualizations in the message, survey, and infographic were understood. Overall, a majority reported a favorable interpretation of the self-referral tool and related materials, indicating that this tool is a positive stimulus to trigger the decision-making process to accept a recommended health action (e.g., getting an LCS). Indeed, results from the pilot test revealed that the tool successfully aided the establishment of pulmonary care for three LCS patients.

While the materials were mostly favorably perceived by patients, there was minor confusion surrounding the QR code on the infographic. The data collected on the infographic suggested that patients perceived self-efficacy in using the QR code, but the qualitative data suggested poor self-efficacy. Further, there are logistical concerns about the QR code being viewed on a mobile phone which would result in people needing help to scan from the same device. The only modification to the pretested materials would involve removing the QR code from the infographic. A link could be provided in addition to the QR code which would allow the patient to directly navigate to the tool rather than using an additional device to do so.

Based on our experience with Epic Software and the initial application results, this tool can successfully be integrated into an EHR and produce self-referrals to LCS. Although the overall response rate was low, nearly half (48.0%) of our patients viewed the initial engagement message, consistent with other results found in other EHR-based messaging cancer screening tools [17]. Many patients that received the tool and did not complete the questionnaire are still eligible for LCS, and more work is needed to improve engagement with the self-referral tool and subsequent LCS uptake. However, this low-cost automated tool has the potential to add to a more considerable effort to increase LCS uptake.

There are several limitations to the study that could limit generalizability. Interviews were not recorded and transcribed verbatim for accuracy but instead collected through notetaking, which impacts response accuracy due to implementation errors. Survey-related effects that may have limited conclusions were response bias, specifically social desirability bias and anchoring bias. The social desirability bias became evident when differences were noted between survey and interview responses regarding the QR code. Additionally, anchoring bias is apparent as new patients were surveyed after their shared decision-making appointment, and their answers could have been influenced by knowledge of lung cancer and LCS obtained during that visit; moreover, this same bias is expected from returning patients surveyed. Returning patients were sampled in addition to new LCS patients to increase feedback and compensate for the poor number of new LCS referrals. Further, the survey was administered by an interviewer face-to-face and not self-administered, so there is the potential for mode effect. Another limitation could lie in the characteristics of the pilot test sample. Participants for this study were selected through convenience sampling of another project that only included a sample of women who completed a mammogram. Expanding the scope of pilot testing to engage racially, ethnically, and gender-diverse individuals who are referred to LCS from various clinical care teams or services (e.g., primary care, gerontology, etc.) could identify optimal tool delivery approaches to increase engagement with and applicability of the tool.

Conclusions

Pretest survey results indicate that the language, phrasing, and visualizations utilized in the tool are understood and increase the potential for engaging in LCS care. Most pretest participants reported favorably interpreting the self-referral tool and related components. This tool is a positive stimulus to trigger the decision-making process to accept a recommended health action (e.g., getting an LCS). Nearly half (48.0%) of the initial application test population opened the message, with an overall response rate of 13.3%. This EHR-integrated tool offers an additional opportunity for referral to

other clinical or ancillary services.

Acknowledgements/Contributions

GSS was responsible for conceptualization, methodology, formal analysis, finding interpretations, writing—original draft, writing—review and editing, and visualization; NTT for funding procurement, conceptualization, methodology, clinical care, writing—review and editing; AH for conceptualization, writing—review and editing, and visualization; JG for conceptualization and writing—review and editing; BAT for funding procurement and writing—review and editing; and AMR for conceptualization, writing—review and editing, and supervision.

Data Availability

De-identified data are available upon request to the corresponding author.

Conflicts of Interest

BAT testifies on behalf of plaintiffs who have filed litigation against the tobacco industry.

Abbreviations

CDC: Center for Disease Control and Prevention
CMS: Centers for Medicare & Medicaid Services
CRA: Clinical research assistant
EHR: Electronic health record
HBM: Health Belief Model
HCC: MUSC Hollings Cancer Center
ICD: International Classification of Diseases
LCS: Lung cancer screening
LDCT: Low-dose computer tomography scan
MRN: Medical record number
MUSC: Medical University of South Carolina
USPSTF: U.S. Preventive Services Task Force

Multimedia Appendix 1

Lung Cancer Infographic Mapped to HBM Constructs

Multimedia Appendix 2

MyChart Mobile App and Desktop Webpage – Patient Point of View

Multimedia Appendix 3

Self-referral Pathway from LCS Eligibility Questionnaire

References

1. Recommendation: Lung Cancer: Screening United States Preventive Services Taskforce (USPSTF); [05/24/2022]; Available from: <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/lung-cancer-screening>.
2. How Common is Lung Cancer? . American Cancer Society; [12/02/2022]; Available from: <https://www.cancer.org/cancer/lung-cancer/about/key-statistics.html>.
3. Ferlay J, Ervik M, Lam F, Colombet M, Mery L, Piñeros M, et al. Global Cancer Observatory: Cancer Today. 2020 [cited 2023]; Available from: <https://gco.iarc.fr/today>.
4. National Lung Screening Trial Research T, Church TR, Black WC, Aberle DR, Berg CD, Clingan KL, et al. Results of initial low-dose computed tomographic screening for lung cancer. *N Engl J Med*. 2013 May 23;368(21):1980-91. PMID: 23697514. doi: 10.1056/NEJMoa1209120.
5. de Koning HJ, van der Aalst CM, de Jong PA, Scholten ET, Nackaerts K, Heuvelmans MA, et al. Reduced Lung-Cancer Mortality with Volume CT Screening in a Randomized Trial. *N Engl J Med*. 2020 Feb 6;382(6):503-13. PMID: 31995683. doi: 10.1056/NEJMoa1911793.
6. Who Should Be Screened for Lung Cancer? : Center for Disease Control and Prevention; [07/13/2022]; Available from: https://www.cdc.gov/cancer/lung/basic_info/screening.htm.
7. World Lung Cancer Day. American Lung Association; [01/01/2023]; Available from: <https://www.lung.org/lung-force/about-lung-force/featured-campaigns/world-lung-cancer-day>.
8. Liu Y, Pan IE, Tak HJ, Vlahos I, Volk R, Shih YT. Assessment of Uptake Appropriateness of Computed Tomography for Lung Cancer Screening According to Patients Meeting Eligibility Criteria of the US Preventive Services Task Force. *JAMA Netw Open*. 2022 Nov 1;5(11):e2243163. PMID: 36409492. doi: 10.1001/jamanetworkopen.2022.43163.
9. State of Lung Cancer Report 2022.
10. Wang GX, Baggett TP, Pandharipande PV, Park ER, Percac-Lima S, Shepard JO, et al. Barriers to Lung Cancer Screening Engagement from the Patient and Provider Perspective. *Radiology*. 2019 Feb;290(2):278-87. PMID: 30620258. doi: 10.1148/radiol.2018180212.
11. Silvestri GA, Nietert PJ, Zoller J, Carter C, Bradford D. Attitudes towards screening for lung cancer among smokers and their non-smoking counterparts. *Thorax*. 2007 Feb;62(2):126-30. PMID: 17101739. doi: 10.1136/thx.2005.056036.
12. Lu MM, Zhang T, Zhao LH, Chen GM, Wei DH, Zhang JQ, et al. The relationship between demands for lung cancer screening and the constructs of health belief model: a cross-sectional survey in Hefei, China. *Psychol Health Med*. 2018;23(8):934-51. PMID: WOS:000437202400004. doi: 10.1080/13548506.2018.1428757.

13. Reihani AR, Ahari, J., Manning, E. P., Cain, H. C., Bade, B. C., & Zimmerman, H. G. . Barriers and facilitators to lung cancer screening in the United States: A systematic review of the qualitative literature. *Journal of Health and Social Sciences*. 2021;6(3):333–48.
14. Ryan BM. Lung cancer health disparities. *Carcinogenesis*. 2018 May 28;39(6):741-51. PMID: 29547922. doi: 10.1093/carcin/bgy047.
15. Triplette M, Thayer JH, Pipavath SN, Crothers K. Poor Uptake of Lung Cancer Screening: Opportunities for Improvement. *J Am Coll Radiol*. 2019 Apr;16(4 Pt A):446-50. PMID: 30777648. doi: 10.1016/j.jacr.2018.12.018.
16. van der Aalst CM, ten Haaf K, de Koning HJ. Lung cancer screening: latest developments and unanswered questions. *Lancet Resp Med*. 2016 Sep;4(9):749-61. PMID: WOS:000382417700022. doi: Doi 10.1016/S2213-2600(16)30200-4.
17. Sequist TD, Zaslavsky AM, Colditz GA, Ayanian JZ. Electronic patient messages to promote colorectal cancer screening: a randomized controlled trial. *Arch Intern Med*. 2011 Apr 11;171(7):636-41. PMID: 21149743. doi: 10.1001/archinternmed.2010.467.
18. Halket D, Dang J, Phadke A, Jayasekera C, Kim WR, Kwo P, et al. Targeted Electronic Patient Portal Messaging Increases Hepatitis C Virus Screening in Primary Care: a Randomized Study. *J Gen Intern Med*. 2022 Oct;37(13):3318-24. PMID: 35230622. doi: 10.1007/s11606-022-07460-1.
19. Epic Systems. 2023.

Supplementary Files

Untitled.


URL: <http://asset.jmir.pub/assets/a9900ea5401e9601ebaafa2293b4d351.docx>

Figures


Initial MyChart Engagement Message Mapped to HBM Constructs.

	Field Notation
Dear @Name@,	
Did you know that most people at risk for lung cancer never get screened? Lung cancer kills more people in the US than breast, colon, and prostate cancer combined.	Perceived Threat
The good news is that screening for lung cancer with a low-dose CT scan can find it early, which is the best chance of a cure.	Perceived Benefit
[HEALTHCARE SYSTEM/HOSPITAL] and the [CLINIC/DEPARTMENT /PROGRAM] care about your preventative health. Due to your patient status at [HEALTHCARE SYSTEM/HOSPITAL], you are eligible to complete this 5 question Lung Cancer Screening Eligibility Survey using your MyChart.	Cue to Action
If you are eligible, an [CLINIC/DEPARTMENT/PROGRAM] team member will contact you.	
Sincerely,	
[CLINICAL TEAM MEMBER(S)] [CLINIC/DEPARTMENT/PROGRAM] [HEALTHCARE SYSTEM/HOSPITAL]	

MyChart Lung Cancer Screening Questionnaire Branching Logic.



Initial Application Test Consort Diagram.



Multimedia Appendixes

Lung Cancer Infographic Mapped to HBM Constructs.

URL: <http://asset.jmir.pub/assets/773f6ce5b5f68c1a9ea33c141cd8444a.pdf>

MyChart Mobile App and Desktop Webpage – Patient Point of View.

URL: <http://asset.jmir.pub/assets/4adddd9e4ae79614749bccfab4232972.docx>

Self-referral Pathway from LCS Eligibility Questionnaire.

URL: <http://asset.jmir.pub/assets/41f96d845c6d5717d0a8fc7c5d10a8c1.pdf>

