

Adding Virtual Reality Mindful Exposure Therapy to a Cancer Center's Tobacco Treatment Offerings: An Acceptability Pilot Study

Riley Walton Jackson, Ann Cao-Nasalga, Amy Chieng, Amy Pirkel, Annemarie D. Jagielo, Cindy Xu, Emilio Goldenhersch, Nicolas Rosencovich, Cristian Waitman, Judith J. Prochaska

Submitted to: JMIR Formative Research
on: November 22, 2023

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5
Supplementary Files..... 20
 Figures 21
 Figure 1..... 22
 Figure 2..... 23

Adding Virtual Reality Mindful Exposure Therapy to a Cancer Center's Tobacco Treatment Offerings: An Acceptability Pilot Study

Riley Walton Jackson¹ MS; Ann Cao-Nasalga² MBA; Amy Chieng³ BA; Amy Pirk² BA; Annemarie D. Jagielo⁴ MS; Cindy Xu² MPH; Emilio Goldenhersch⁵ BS; Nicolas Rosencovich⁶ BS; Cristian Waitman⁷ BA; Judith J. Prochaska³ MPH, PhD

¹MS program in Community Health and Prevention Research Department of Medicine Stanford University Stanford US

²Health Education, Engagement and Promotion Stanford Health Care Stanford US

³Stanford Prevention Research Center Department of Medicine Stanford University Palo Alto US

⁴PGSP-Stanford Psy.D. Consortium Palo Alto University Palo Alto US

⁵Laboratorio de Investigación en Neurociencia y Ciencias Sociales Universidad de Flores Buenos Aires AR

⁶Escuela de Ingeniería Biomédica Universidad Nacional de Córdoba Córdoba AR

⁷Universidad Siglo XXI Córdoba AR

Corresponding Author:

Judith J. Prochaska MPH, PhD
Stanford Prevention Research Center
Department of Medicine
Stanford University
3180 Porter Drive
Palo Alto
US

Abstract

Background: Smoking contributes to 1 in 3 cancer deaths. At the Stanford Cancer Center, tobacco cessation medication management and counseling are provided as a covered benefit. Patients charted as using tobacco are contacted by a tobacco treatment specialist and offered cessation services. A novel addition, the current study examined the acceptability of a virtual reality (VR) mindful exposure therapy app for quitting smoking called MindCotine.

Objective: Determine the feasibility and acceptability of offering 6-weeks of MindCotine treatment as a part of Stanford's Tobacco Treatment Services for patients seen for cancer care.

Methods: As part of a pilot study, the MindCotine VR program was offered to English or Spanish-speaking patients interested in quitting smoking. Given the visual interface, epilepsy was a medical exclusion. Viewed from a smartphone with an attachable VR headset, MindCotine provides a virtual environment with audiovisual content guiding mindfulness exercises (e.g., breathing techniques, body awareness, thought recognition), text-based coaching, and cognitive behavioral therapy-based self-reflections for quitting smoking. Interested patients providing informed consent were mailed a MindCotine headset and asked to utilize the app 10+ minutes a day. At the end of 6 weeks, participants completed a feedback survey.

Results: Of the 357 patients reached by the tobacco treatment specialist, 62 (17.3%) were ineligible, 190 (53.2%) were not interested in tobacco treatment services, and 78 (21.8%) preferred other tobacco treatment services. Among the 105 eligible and interested in assistance with quitting, 27 (25.7%) were interested in MindCotine, of whom 20 completed the informed consent, 9 used the program, and 8 completed their end of treatment survey. Participants using MindCotine completed, on average, 13 (SD=20.2) program activities, 19 (SD=26) journal records, and 11 (SD=12.3) coaching engagements. Of the 9 participants who used MindCotine, 4 reported some dizziness with app use that resolved (44%), and 7 would recommend MindCotine to a friend (78%). Two participants quit tobacco (22.2% reporting, 10% overall), two others reduced their smoking by 50% or more, and two quit for 24-hours and then relapsed.

Conclusions: In an acceptability pilot study of a novel VR tobacco treatment app offered to patients at a cancer center, 4 of 9 (44%) reporting and 4 of 20 (20%) overall, substantially reduced or quit using tobacco after 6-weeks and most would recommend the app to others. Further testing in a larger sample is warranted.

(JMIR Preprints 22/11/2023:54817)

DOI: <https://doi.org/10.2196/preprints.54817>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

✓ **Only make the preprint title and abstract visible.**

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/>

Original Manuscript

Original Paper**Adding Virtual Reality Mindful Exposure Therapy to a Cancer Center's Tobacco Treatment Offerings: An Acceptability Pilot Study**

Riley Walton Jackson, MS,¹ Ann Cao-Nasalga, MBA,² Amy Chieng, BA,³ Amy Pirkel, BA,² Annemarie D. Jagielo, MS,⁴ Cindy Xu, MPH,² Emilio Goldenhersch, BS,⁵ Nicolas Rosencovich, BS,⁶ Cristian Waitman⁷, BA, Judith J. Prochaska, PhD, MPH³

¹ MS program in Community Health and Prevention Research, Department of Medicine, Stanford University, Stanford, CA, USA

²Health Education, Engagement and Promotion, Stanford Health Care, Stanford, CA, USA

³Stanford Prevention Research Center, Department of Medicine, Stanford University, Stanford, CA, USA

⁴PGSP-Stanford Psy.D. Consortium, Palo Alto University, Palo Alto, CA, USA

⁵Laboratorio de Investigación en Neurociencia y Ciencias Sociales, Universidad de Flores, Ciudad Autónoma de Buenos Aires, Argentina

⁶Escuela de Ingeniería Biomédica, Universidad Nacional de Córdoba, Córdoba, Argentina

⁷Universidad Siglo XXI, Córdoba, Argentina

Corresponding Author: Judith Prochaska, PhD, MPH

3180 Porter Dr #A105

Palo Alto, CA 94304-1212

Phone: 650-724-3608

Email: jpro@stanford.edu

Abstract

Background: Smoking contributes to 1 in 3 cancer deaths. At the Stanford Cancer Center, tobacco cessation medication management and counseling are provided as a covered benefit. Patients charted as using tobacco are contacted by a tobacco treatment specialist and offered cessation services. A novel addition, the current study examined the acceptability of a virtual reality (VR) mindful exposure therapy app for quitting smoking called MindCotine.

Objective: Determine the feasibility and acceptability of offering 6-weeks of MindCotine treatment as a part of Stanford's Tobacco Treatment Services for patients seen for cancer care.

Methods: As part of a pilot study, the MindCotine VR program was offered to English or Spanish-speaking patients interested in quitting smoking. Given the visual interface, epilepsy was a medical exclusion. Viewed from a smartphone with an attachable VR headset, MindCotine provides a virtual environment with audiovisual content guiding mindfulness exercises (e.g., breathing techniques, body awareness, thought recognition), text-based coaching, and cognitive behavioral therapy-based self-reflections for quitting smoking. Interested patients providing informed consent were mailed a MindCotine headset and asked to utilize the app 10+ minutes a day. At the end of 6 weeks, participants completed a feedback survey.

Results: Of the 357 patients reached by the tobacco treatment specialist, 62 (17.3%) were ineligible, 190 (53.2%) were not interested in tobacco treatment services, and 78 (21.8%) preferred other tobacco treatment services. Among the 105 eligible and interested in assistance with quitting, 27 (25.7%) were interested in MindCotine, of whom 20 completed the informed consent, 9 used the program, and 8 completed their end of treatment survey. Participants using MindCotine completed, on average, 13 (SD=20.2) program activities, 19 (SD=26) journal records, and 11 (SD=12.3) coaching engagements. Of the 9 participants who used MindCotine, 4 reported some dizziness with app use that resolved (44%), and 7 would recommend MindCotine to a friend (78%). Two participants quit tobacco (22.2% reporting, 10% overall), two others reduced their smoking by 50% or more, and two quit for 24-hours and then relapsed.

Conclusions: In an acceptability pilot study of a novel VR tobacco treatment app offered to patients at a cancer center, 4 of 9 (44%) reporting and 4 of 20 (20%) overall, substantially reduced or quit using tobacco after 6-weeks and most would recommend the app to others. Further testing in a larger sample is warranted.

Keywords: tobacco cessation; virtual reality; exposure therapy; cancer care

Introduction

Tobacco use, and in particular cigarette smoking, has taken a devastating toll on public health. While tobacco smoking has declined significantly over the past several decades, it remains the leading cause of preventable disease, disability, and death in the United States [1] and worldwide. Smoking contributes to one in three cancer deaths [2, 3], and more than half of all people who smoke long term will die from a tobacco-related disease, losing ten years of life on average [2].

Quitting smoking proves to be a difficult task, even when individuals have a strong desire to stop. Nearly 7 in 10 people who smoke want to quit, and 55% of people who smoke, quit for at least one day in a given year [1, 2]. Yet, 60% of those who quit for a day end up relapsing within one week [2]. Of the 7% of initial quit attempts that last for 6 months, 45% of those result in relapse [2].

Nicotine's effects are transient and within an hour of having a cigarette, withdrawal symptoms can begin. These include irritability, anxiety, depressed mood, and difficulty concentrating [2]. There also is a behavioral component of tobacco use disorder [2, 4]. The act of smoking becomes associated with the pleasurable effects of nicotine, and functions as a coping mechanism when individuals feel negative emotions [2, 4, 5]. Furthermore, environmental and situational factors, such as being around friends who smoke, or watching television, can become linked with nicotine and act as triggers or cues for usage [2, 6]. Thus, effective treatments often combine multiple strategies to target both the behavioral and physiological aspects of nicotine dependence [2, 4, 5].

Studies of smoking and different forms of cancer indicate that between 35% and 72% of patients who smoke continue to smoke after cancer diagnosis and treatment [3, 7]. Continuing to smoke during treatment can decrease the effectiveness of radiotherapy, wound healing efficiency, and survival time [3, 7]. Smoking also can increase the risk of treatment side effects and complications, cancer recurrence, and second primary malignancies [3, 7]. Quitting smoking can improve treatment outcomes and survival [3]. Cancer screening, diagnosis, and hospitalization can motivate and support a quit attempt, making it a critical time for engagement with cessation resources and treatment [1, 3, 7].

Traditional treatment strategies for quitting smoking include behavioral counseling (individual or group) and cessation medications [1, 2, 4], such as nicotine replacement therapy (NRT). These evidence-based treatments increase quit rates when utilized, and clinical practice guidelines recommend using them in combination [4, 5]. However, not all individuals are receptive to or able to take cessation medications [8]. Behavioral counseling might not be available or preferable to patients due to travel, time, and cost constraints, in addition to concerns about privacy or feelings of shame (e.g., in group counseling) [2, 9].

To address these challenges, additional delivery methods for smoking cessation interventions have been developed. As of 2021, 85% of American adults own a smartphone, and there is an immense potential for reach, accessibility, flexibility, and cost-effectiveness of mobile health interventions [2, 10, 11]. Mobile health interventions can ideally be combined with pharmacotherapy and counseling, or they may be used as standalone therapies for patients who are not interested in or candidates for cessation medications and traditional behavioral counseling. There is evidence for the efficacy of Internet, Quitline, and text-based cessation resources as well as preliminary evidence for mobile apps [1, 2, 9, 10]. Evidence for the efficacy of technology-based interventions in smoking cessation is limited, yet promising [10, 12].

An emerging field for the treatment of mental health disorders, including substance use disorders, is

virtual reality (VR). VR generally refers to an interactive, computer generated experience that consists of a simulated, immersive environment [9, 13], [14]. VR can be used to create virtual environments that elicit challenging sensations and emotions, which can be utilized to create opportunities for learning and practicing skills to manage them [6, 9, 13, 14]. VR immersion and interactivity can vary with different types of headsets (smartphone app based, standalone, tethered) and apply different theoretical approaches (e.g. cognitive behavioral, gamified behavioral, cue exposure) with varying outcomes [15]. The most established evidence for utilization of VR in treatment is with exposure-based treatment for anxiety disorders [13, 14]. There is a small but growing body of evidence indicating that VR is effective in inducing cravings for cigarettes, thus allowing for the opportunity to practice virtually presented coping strategies [6, 9, 13, 14, 15].

A randomized controlled pilot study on the efficacy of a smartphone app-based VR mindful exposure therapy (VR-MET) smoking cessation program developed by MindCotine was conducted in 2018. VR-MET incorporated mindfulness training and cue exposure as part of the VR intervention. While VR sessions typically occur in a lab or hospital-based setting with supervising staff, MindCotine VR sessions are able to be self-administered and delivered remotely [6, 9]. With a headset that connects to a smartphone, MindCotine creates an immersive environment in a cost-effective and scalable manner [9]. This contrasts with other forms of VR that are less immersive (utilize a computer monitor only) or highly expensive such as head-mounted systems with separate displays for each eye [6, 14]. Preliminary evidence shows that mindfulness interventions can be efficacious for smoking cessation in traditional, in-person, and technology-based formats [9, 12]. Additionally, evidence supports that CBT is effective as a smoking cessation treatment when delivered in a traditional format [10, 12]. CBT focuses on identifying and changing maladaptive thoughts and behavioral patterns to alleviate distress. The app utilizes a series of CBT reflective questions and journaling for participants to evaluate changes in their relationship with smoking and offers text-based coaching. In this randomized controlled study, a sample of 120 individuals who smoked at least 5 cigarettes a day living in Buenos Aires, Argentina, was recruited via Facebook and television advertisements, with 93% of the sample completing the full 21-day program [9]. Individuals in the control condition received a self-help manual on smoking cessation by the Argentine Ministry of Health. At post-intervention, the treatment group participants reported a significant decrease in cigarettes per day, lower cravings, increased readiness to quit, and higher abstinence rates compared to the control group [9]. Overall, the MindCotine intervention was found to be feasible and acceptable, with preliminary evidence for supporting smoking cessation in a nonclinical community sample. Unexplored is use of the VR-MET app in a clinical sample and among people with lower levels of smoking. With a focus on integrated cancer care, we aimed to test VR-MET as a novel tobacco cessation treatment addition to our clinic's offerings.

In 2018, the Stanford Cancer Institute integrated tobacco treatment services within cancer care, as part of the Cancer Center Cessation Initiative (C3I) [3]. Collaboration with three pilot clinics began in 2019 (head and neck, thoracic, gastrointestinal), and by 2020, the Stanford Tobacco Treatment Service had successfully extended into more than 20 clinics at the cancer center [3]. The service continues to explore expansion opportunities to improve patient reach and engagement during cancer care, including collaborating with MindCo Health in exploring VR smoking cessation with this pilot study.

The aim of the current study was to determine the feasibility and acceptability of a VR tobacco cessation treatment among patients seen for cancer care who smoke. If feasible and acceptable, the intervention could expand the treatment options available to patients, potentially improve cessation rates, and help improve cancer care outcomes. A secondary objective was to study efficacy of the

intervention in a clinical sample. Furthermore, the COVID-19 pandemic has altered traditional healthcare delivery. Many patients are immunocompromised due to their anticancer treatment or the disease itself, and there has been an increase in the use of telehealth services for cancer care to avoid unnecessary exposure to COVID-19 [16]. Thus, the implementation of a remote delivery smoking cessation treatment may be particularly beneficial for patients undergoing cancer treatment.

This pilot differs from the initial efficacy trial by sample. The initial efficacy trial studied adults with a high level of daily smoking residing in Buenos Aires, Argentina, while this study focused on adult patients of the Stanford Cancer Center who smoked daily or non-daily [9]. Patients with cancer face unique barriers as participants in research studies, including lasting side effects of treatment and coordinating many different care providers [17]. Participating in a research study may seem overwhelming and difficult to manage during cancer treatment. However, the Tobacco Treatment Services at Stanford have been able to reach 69% of patients identified as using tobacco and engage 38% of the patients reached in cessation treatment services [3, 7]. Treating smoking can greatly improve cancer care outcomes [7].

Methods

Study Design

This was a single group pilot study designed for a maximum of 20 participants assessing outcomes after 6-weeks of use. The primary objective was to determine the feasibility and acceptability of this VR intervention. Feasibility and acceptability metrics were at least (i) 20% interested in using the app; (ii) 75% of interested patients onboarded and using the app; and (iii) 70% recommending the app to others. The secondary objective was to capture efficacy by evaluating quit attempts, smoking reduction, and abstinence. Goals were at least 50% reducing tobacco use and 10% quitting. The metrics and tobacco usage were assessed via survey at enrollment and end of treatment.

Ethical Considerations

Study approvals were obtained from the Stanford University Institutional Review Board on May 18, 2021 (protocol number 60253) and the Stanford Cancer Center Scientific Review Committee on September 19, 2021. There was no compensation for participation in the study. The study was offered as an option as part of the Stanford Tobacco Treatment Service. All study materials were provided to participants at no cost and the VR kits did not need to be returned.

The study had a waiver of documentation to allow for telephone consent to screen and the eligibility questionnaire. The eligibility questionnaire also functioned as the baseline questionnaire for those eligible for the study. Informed consent procedures used the Adobe Sign platform and were compliant with HIPAA and 21CFR11. Study staff was available via phone to go through consent. Each potential participant was able to take time to consider participation and could drop out at any time knowing their participation was voluntary. No other tobacco treatment options were withheld. Materials were available in English and Spanish. Stanford Health Care Interpreter Services could be used during consent as needed.

Assessment data were collected and stored via REDCap, a HIPAA compliant survey administration tool. The MindCotine app was reviewed by Stanford's Data Risk Assessment team prior to recruitment start. The study also underwent annual reviews by the Stanford Cancer Center Data Safety Monitoring Committee.

The Stanford clinical team handled the outreach portion and informed the research team which patients would like to participate in the study. The Stanford research team managed mailing of kits, end of treatment, and data analysis. MindCotine had access to the app user data, which was provided to the Stanford research team through secure email. The Stanford research team provided a summary of outcomes (no PHI) to MindCotine.

Recruitment

Study recruitment occurred between November 16, 2021 and August 18, 2022. Patients seen for cancer care at the Stanford Cancer Center, who were charted as using tobacco products in the electronic health record, were included in a weekly report that the health system tobacco treatment specialist used to reach out to patients offering tobacco treatment services. Patients were offered a variety of treatment options including individual counseling, group counseling, cessation medications (e.g., NRT, varenicline, bupropion), and the MindCotine VR intervention. The MindCotine intervention was described to patients as a “mobile-based virtual reality program for quitting smoking,” and they were told that “the program teaches skills for quitting smoking and is very visual. The treatment can be done at home from your phone.”

Patients who expressed interest in MindCotine completed a phone screen to determine eligibility. Inclusion criteria were smoking daily or non-daily (at least 9 cigarettes per week), fluency in English or Spanish, owning a compatible smartphone, open to assistance with quitting smoking, and willing to use the MindCotine app at least 10 minutes daily for 6 weeks and to complete survey assessments. Exclusion criteria were history of seizures or an epilepsy diagnosis.

Study Procedures

Patients deemed eligible went through an informed consent process and were sent a password protected link to read and e-sign the informed consent form. Once consent was completed, a MindCotine VR kit was mailed to participants, which included the cardboard or plastic headset and an activation code to gain full access to the app for a year at no cost. The app shared suggestions, such as sitting comfortably and using headphones before the VR experience. Users could also schedule meetings with the MindCotine app coaching team, where further explanation on regulating the focus with the VR headset could be provided. Instructions were delivered by phone or video call. At the end of the 6 weeks, participants were contacted by a research team member to complete an end of treatment assessment.

Intervention

This study utilized the MindCotine smartphone app for the VR smoking cessation intervention. The MindCotine intervention consists of audiovisual media with VR and non-VR content drawing on mindfulness-based techniques and cognitive behavioral therapy (CBT) strategies for addressing smoking cravings [9]. Treatment components within the MindCotine app are summarized in Table 1. VR sessions focus on the act of smoking and cravings, stress at work, and bodily sensations. The VR intervention included 7 VR-MET sessions, each lasting 10 minutes. These experiences consist of a combination of animated environments at the beginning and end of each session lasting 4 minutes total to induce meditative states, and 6 minutes of cue-exposure therapy filmed with a 360 camera. The reason behind this integrated approach is based on studies showing the use of VR in natural environments to create relaxation and VR with real people to simulate familiar situations to elicit cravings and bodily sensations [18, 19]. Non-VR sessions focus on formal mindfulness-based exercises such as deep breathing, body scan, and recognizing thoughts, emotions, and the impulse of smoking.

Table 1. MindCotine App Components for Treating Tobacco Use.

App Component	Description of Component	Virtual Reality
Program Activity	Virtual Reality – Mindful Exposure Therapy (VR-MET) exercises and guided mindfulness-based two-dimensional video, audio, and reflection exercises.	Yes
Journal Record	Cognitive Behavioral Therapy (CBT) based self-reflections. Includes tracking cigarette use per day.	No
Coaching Engagement	Communicating with a coach through the MindCotine mobile application.	No
Lifesaver Use	Brief informational intervention via chatbot.	No
Input Activity	Inputting an emotion experienced.	No
Commitment Activity	Setting a quit date.	No

Participants were encouraged to engage with MindCotine for 6 weeks, with a different concept of focus each week (Table 2). A daily session of MindCotine varied between self-reflecting CBT questions, mindfulness audios or videos, and VR experiences, including daily journaling of cigarette consumption. The time varied between 1 to 10 minutes of usage. Due to the nature of VR-MET, every VR experience was embedded as part of the 6-week intervention to address the Concepts (Table 2) under situations that were familiar to the patient and could train their emotional and practical coping skills in the comfort of their home, while identifying the associated-sensations elicited by the specific cues. All VR experiences were immersive but were not interactive in order to reduce potential dizziness.

Table 2: Overview of 10-minute weekly concepts in the six-week MindCotine intervention [16].

Week	Concept	Description
1	Self-awareness	Breathing techniques, the act of smoking, and reasons / motivations to start the program.
2	Sensations	Focusing on bodily sensations in relation to smoking and triggers.
3	Emotions	Managing cravings born out of stress and feelings related to familiar situations.
4	Cognition	Exploring cognitive distortions, beliefs, and activities led by curiosity and reflections.
5	Readiness	Practicing empathy and past victories to determine future actions.
6	Preparation	Setting new structures that can stimulate resilience and behavior activation tasks.

Statistical Analysis and Outcome Goals


Descriptive statistics were utilized to summarize tobacco use characteristics, changes in tobacco use, MindCotine app usage, and feedback on the MindCotine app. Feasibility (primary objective) was defined as (i) at least 20% of patients open to assistance with quitting reporting interest in using the MindCotine app, (ii) 75% of the interested patients being fully onboarded and using the app, and (iii) 70% of participants who complete end of treatment survey indicating that they would recommend the app to others. Efficacy (secondary objective) was defined as at least 50% of participants reducing their tobacco use and 10% or more quitting.

Results

Eligibility, Engagement, & Retention

Of 357 patients reached by the tobacco treatment specialist, 62 (17.3%) were ineligible, 190 (53.2%) were not interested in treatment for quitting tobacco, and 78 (21.8%) preferred other cessation treatment services (Figure 1). Of the 27 participants interested in MindCotine (25.7% of the 105 eligible and interested in quit assistance), 20 (74.1%) completed the consent process. Of the 20 participants, 9 (45%) used the MindCotine program, and 8 (40%) completed the end of treatment assessment.

Figure 1: Participant flow diagram of the MindCotine pre to post 6-week smoking cessation intervention among Stanford Cancer Center patients.




Baseline Sample Characteristics

Of the 20 participants who consented to participate, there were 11 females and 9 males; the mean age was 52 years [SD=13 years, range 35 to 75 years]; 15 identified as White, 2 as Asian, 1 as Black, 1 as Multiracial, and 1 was unreported. All were non-Hispanic. At study start, 18 of 20 (90%) were smoking cigarettes daily and averaged 14.7 cigarettes (SD=7.2) per day (median=15, range: 4-30). Among the 2 (10%) who reported smoking on some days, they averaged 25 cigarettes per week (SD=7, range: 20-30). At baseline, 50% (10/20) were not using any other cessation support; 7 (35%) were using a combination of counseling services and cessation medication; 2 (10%) were only using cessation medications; 1 (5%) was using counseling services only.

MindCotine Use

Participants using MindCotine completed, on average, 13 (SD=20.2, range=2-65; median=4) program activities, 19 (SD=26, range=0-87; median=11) journal records, and 11 (SD=12.3, range=0-40; median=7) coaching engagements (Figure 2). In general, without limiting to the 6-week treatment period, active use of the MindCotine app ranged from 3 to 266 days (mean=63, SD=80; median=48).

Figure 2. N=9 Participant usage of MindCotine VR-MET program components during the study period.



Feasibility and Acceptability

Two of three conditions for the primary objective of feasibility and acceptability were met: (1) 27.5% of those eligible and interested in help with quitting smoking were interested in MindCotine (surpassing goal of 20%); and (2) 78% of those who used MindCotine would recommend the app to a friend (surpassing goal of 70%). Less than half (45%, 9/20) of the consented and interested patients, however, were fully onboarded and used the app (goal of 75% not met).

Of the 9 participants who used the MindCotine app, 8 completed the end of treatment survey and provided feedback about the program. 7 of the 8 would recommend MindCotine to a friend (88% of those reporting, 78% of those who used the app). Five of 8 participants (62.5%) provided positive comments about their experience with the app, noting that it was enjoyable, fun, and helpful for people who want to quit smoking. Additionally, there were several points recommended for improvement or consideration when using the program. For example, 3 of 8 participants (37.5%) described headset discomfort and issues of alignment between the headset and smartphone that negatively impacted their experience. Two participants (25%) reported having difficulty getting started with the program and required assistance from clinic staff or MindCotine staff to set up the program. Three out of 8 (38%) completing the end of treatment survey reported mild to moderate VR related symptoms (e.g. headache, disorientation, dizziness, motion sickness, nausea) with app use. One additional participant who withdrew mentioned experiencing dizziness and lightheadedness. In all cases, removal of the headset or discontinuing use of the app resolved the symptoms as expected.

Smoking Outcomes

One of the two conditions for the secondary objective of efficacy was met. Smoking outcomes were available from both the MindCotine app and the end of treatment survey. Of the 9 participants with smoking outcomes, two participants quit tobacco (22% reporting, 10% overall [2/20]; goal: 10%), two additional participants reduced their smoking by 50% or more (44.4% reporting, 20% overall [4/20]; goal: 50%), and two quit for 24-hours then relapsed. Mean change in cigarette use was -4.1 cigarettes per day (SD=8; range: -20 to 9). Of the 8 completing their end of treatment survey, 5 (62%; 25% overall) were using cessation medications (e.g. nicotine replacement therapy, varenicline, bupropion).

Discussion

Principal Results

Two of three conditions for the primary objective of feasibility and acceptability were met: (1) 27.5% of those eligible and interested in help with quitting smoking were interested in MindCotine (surpassing goal of 20%); and (2) 78% of those who used MindCotine would recommend the app to a friend (surpassing goal of 70%). One of the two conditions for the secondary objective of efficacy was met: Two of the nine participants with smoking outcomes quit tobacco (22% reporting, 10% overall [2/20]; goal: 10%). Outcomes for feasibility, acceptability, and efficacy show promise but with less than half of the consented patients using the app, further testing of MindCotine in a larger sample is warranted.

In the current study, we were successful in reaching 357 patients to offer cessation services. A majority (57.5%) declined assistance with quitting smoking and 62 (17.4%) were not eligible for the study. Worth highlighting is that the patients contacted about the study were proactively called by the program's tobacco treatment specialist after being identified as using tobacco by the Stanford Cancer Center. Barriers to accessing conventional cessation treatments such as cost, travel, and coordination were removed [1, 2, 9]. Therefore, it is unsurprising that not being ready to quit or declining cessation treatment were among the most common reasons for not participating in the MindCotine pilot. A higher level of participation would be anticipated in settings reliant on patients calling in (i.e., reactive rather than proactive outreach model) who ostensibly would present with higher interest in quitting smoking. Those who opted for cessation support other than MindCotine instead chose a healthcare provider, counseling, and/or pharmacotherapy (19.6%, 70/357). In Stanford's Tobacco Treatment Program, patients initially start with one treatment option (typically 1:1 counseling) and may include another option later as suggested by their individual counselor. This

likely explains why a large proportion of patients not interested in our study cited their engagement with another cessation treatment or resource as the reason. Uptake of the MindCotine app could improve over time and could be optimized if integrated within counseling sessions. Adding a brief motivational video also could aid adoption of MindCotine. Though quit attempts are significantly more likely to be successful when individuals utilize cessation pharmacotherapy and behavioral counseling [5, 20, 21], most US adults who smoke try to quit without support and most quit attempts fail [1]. Further research is needed to understand hesitancy to engage in support for quitting smoking [20].

Study Limitations

The low completion rate of the intervention made it difficult to draw conclusions about the efficacy of the MindCotine VR treatment. The small sample had limited demographic diversity. The study sample was not large enough to explore the effects of other smoking cessation options with or without the VR-MET treatment, which is worth exploring in future studies especially for patients who have exhausted current treatment options as monotherapies. The field would benefit from more studies of VR treatment comparisons [15]. Additionally, this study had a short follow up time (follow-up occurred immediately after program completion), and information on tobacco usage at baseline and end of treatment was self-reported rather than biochemically verified.

Study Strengths

This study offered VR-MET as one of multiple treatment options for smoking cessation in a clinical sample undergoing cancer care. The menu of options allowed patients to choose what may work best for them without concern of being limited to one type of treatment, demonstrating practical application and outcomes in a healthcare setting.

While the scope of this pilot study is limited, there was demonstrated interest among patients undergoing cancer care in trying VR-MET for smoking cessation. Including people who have cancer in research studies may be challenging given coordination of care and lasting side effects from treatment, but it remains extremely important for health and treatment recovery. Also worth testing is VR for smoking cessation and relapse prevention in other clinical samples where treatment outcomes are acutely affected by tobacco use, such as in the perioperative setting.

Comparison with Prior Work

This pilot of MindCotine differed from an earlier efficacy trial in enrollment and onboarding. The first was that participants in the efficacy trial were invited to meet with the research team in-person for consent, onboarding, and distribution of the MindCotine headset kit [9]. Most (80%, 120/150) of those invited to use the app confirmed their participation and completed the consent process, compared to 74% (20 of 27) in our pilot [9]. In our pilot, all aspects of the study occurred remotely. The remote method may have contributed to the slightly lower proportion of patients who completed the consent process.

In terms of scale and costs, smartphone app-based VR treatments have these benefits over standalone and tethered VR sets. The MindCotine VR-MET was a smartphone app-based treatment, which allowed for freedom of wherever the user decided to engage in smoking cessation treatment, and shipping of the VR kit did not involve handling fragile parts. Standalone or tethered VR kits generally have better audio-visual quality and interactivity but are often expensive, with tethered sets limited to locations with electrical outlets [22]. The MindCotine VR-MET environment included cue-exposure therapy using still images filmed with a 360 camera, and animations for inducing meditative states. Interactivity beyond seeing and practicing cognitive behavioral strategies were not necessary for this intervention. It may be worthwhile to see the efficacy of CBT strategies pre and post intervention with interactivity in the VR environment (e.g., users having options of picking up a virtual cigarette vs virtual substitute). However, this could also be done with smartphone app-based headsets where looking at an object long enough would count as a decision to use the object, as demonstrated in a smartphone app-based VR escape game [23].

Prior evaluations of VR interventions for smoking cessation largely studied adults outside of clinical settings and supported VR cue exposure therapy; more work is needed to explore multiple and combined treatment modalities and integration within clinical settings [15]. Krebs et al.'s study, similar to our pilot in analytic sample size (n=8), but with a focus on patients recently hospitalized for cancer care, aimed to prevent relapse among those who recently quit smoking [6]. Krebs et al.'s team used a virtual world simulation game for treatment delivery to build skills to cope with smoking cues. Their game was found to cue cravings as designed and virtual practice of coping was viewed as helpful.

Our evaluation of the MindCotine VR intervention found that remote treatment and low-cost headsets for VR immersion are largely feasible and acceptable among patients receiving cancer care with varying levels of cigarette use. Quality testing onboarding procedures and adjustments with the headset are recommended prior to scaling to a remote study of feasibility, acceptability, and efficacy in a larger demographically diverse sample of patients undergoing cancer care.

Conclusions

In this initial clinical pilot, the MindCotine VR intervention was of interest to more than one in four patients open to assistance with quitting smoking. Though remote onboarding and use of the technology proved challenging for some, most would recommend MindCotine to others and a substantial proportion quit or reduced their smoking. The positive feedback and outcomes in reducing smoking from those who completed the intervention indicate that further testing of MindCotine in a larger sample is warranted. Further, uptake and sustained use are anticipated to be greater among people who are actively seeking strategies to quit smoking and perhaps among young adults and those familiar with digital games and VR environments.

Acknowledgements

Funding from National Cancer Institute Moonshot Supplements 3P30CA124435-11S2/12S1 and 3P30CA124435-13S. We thank Sharon Hall, PhD, for her guidance and feedback as a reader of the master's thesis version of this manuscript.

Conflicts of Interest

Emilio Goldenhersch, Nicolas Rosencovich, and Cristian Waitman are all employees of MindCo Health.

Abbreviations

CBT: Cognitive behavioral therapy

NCI: National Cancer Institute

NRT: Nicotine Replacement Therapy

TTS: Tobacco Treatment Specialist

VR: Virtual reality

VR-MET: virtual reality mindful exposure therapy

Data Availability

Access to the data is restricted for patient privacy protections.

References


- [1] U.S. Department of Health and Human Services, "Smoking Cessation: A Report of the Surgeon General (Executive Summary)," *US Dep. Health Hum. Serv. Cent. Dis. Control Prev. Natl. Cent. Chronic Dis. Prev. Health Promot. Off. Smok. Health*, p. 28, 2020.
- [2] J. J. Prochaska and N. L. Benowitz, "Current advances in research in treatment and recovery: Nicotine addiction," *Sci. Adv.*, vol. 5, no. 10, p. eaay9763, Oct. 2019, doi: 10.1126/sciadv.aay9763.
- [3] K. Gali *et al.*, "Integration of Tobacco Treatment Services into Cancer Care at Stanford," *Int. J. Environ. Res. Public. Health*, vol. 17, no. 6, p. 2101, Mar. 2020, doi: 10.3390/ijerph17062101.
- [4] R. L. Corelli and K. S. Hudmon, "Integrating Medications into Smoking Cessation Treatment: The Basics," UCSF Smoking Cessation Leadership Center. Accessed: Feb. 01, 2022. [Online]. Available: <https://ucsf.app.box.com/s/iagrat06fq4bvwrafufidtvewutnkwwb>
- [5] C. D. Morris and A. Sripipatana, "A Team Approach: Integrating Tobacco Dependence Treatment into Routine Clinical Practice," UCSF Smoking Cessation Leadership Center. Accessed: Feb. 23, 2022. [Online]. Available: <https://smokingcessationleadership.ucsf.edu/webinar/team-approach-integrating-tobacco-dependence-treatment-routine-clinical-practice>
- [6] P. Krebs *et al.*, "Development of a Virtual Reality Coping Skills Game to Prevent Post-Hospitalization Smoking Relapse in Tobacco Dependent Cancer Patients," *J. Virtual Worlds Res.*, vol. 2, no. 2, p. 470, Aug. 2009.
- [7] J. C. de Bruin-Visser, A. H. Ackerstaff, H. Rehorst, V. P. Retèl, and F. J. M. Hilgers, "Integration of a smoking cessation program in the treatment protocol for patients with head and neck and lung cancer," *Eur. Arch. Otorhinolaryngol.*, vol. 269, no. 2, pp. 659–665, Feb. 2012, doi: 10.1007/s00405-011-1673-0.
- [8] Office of the Commissioner, "Want to Quit Smoking? FDA-Approved Products Can Help," *FDA*, Sep. 2020, Accessed: Mar. 17, 2022. [Online]. Available: <https://www.fda.gov/consumers/consumer-updates/want-quit-smoking-fda-approved-products-can-help>
- [9] E. Goldenhersch, J. Thrul, J. Ungaretti, N. Rosencovich, C. Waitman, and M. R. Ceberio, "Virtual Reality Smartphone-Based Intervention for Smoking Cessation: Pilot Randomized Controlled Trial on Initial Clinical Efficacy and Adherence," *J. Med. Internet Res.*, vol. 22, no. 7, p. e17571, Jul. 2020, doi: 10.2196/17571.
- [10] C. Tudor-Sfetea *et al.*, "Evaluation of Two Mobile Health Apps in the Context of Smoking Cessation: Qualitative Study of Cognitive Behavioral Therapy (CBT) Versus Non-CBT-Based Digital Solutions," *JMIR MHealth UHealth*, vol. 6, no. 4, p. e9405, Apr. 2018, doi: 10.2196/mhealth.9405.
- [11] Pew Research Center, "Demographics of Mobile Device Ownership and Adoption in the United

- States,” Pew Research Center: Internet, Science & Tech. Accessed: Mar. 17, 2022. [Online]. Available: <https://www.pewresearch.org/internet/fact-sheet/mobile/>
- [12] C. Vinci, “Cognitive Behavioral and Mindfulness-Based Interventions for Smoking Cessation: a Review of the Recent Literature,” *Curr. Oncol. Rep.*, vol. 22, no. 6, p. 58, May 2020, doi: 10.1007/s11912-020-00915-w.
- [13] J. Worley, “Virtual Reality for Individuals With Substance Use Disorders,” *J. Psychosoc. Nurs. Ment. Health Serv.*, vol. 57, no. 6, pp. 15–19, Jun. 2019, doi: <http://dx.doi.org/10.3928/02793695-20190430-01>.
- [14] D. Freeman *et al.*, “Virtual reality in the assessment, understanding, and treatment of mental health disorders,” *Psychol. Med.*, vol. 47, no. 14, pp. 2393–2400, Oct. 2017, doi: 10.1017/S003329171700040X.
- [15] M. Keijsers, M. C. Vega-Corredor, M. Tomintz, and S. Hoermann, “Virtual Reality Technology Use in Cigarette Craving and Smoking Interventions (I ‘Virtually’ Quit): Systematic Review,” *J. Med. Internet Res.*, vol. 23, no. 9, p. e24307, Sep. 2021, doi: 10.2196/24307.
- [16] J. van de Haar *et al.*, “Caring for patients with cancer in the COVID-19 era,” *Nat. Med.*, vol. 26, no. 5, Art. no. 5, May 2020, doi: 10.1038/s41591-020-0874-8.
- [17] J. Easley *et al.*, “Coordination of cancer care between family physicians and cancer specialists,” *Can. Fam. Physician*, vol. 62, no. 10, pp. e608–e615, Oct. 2016.
- [18] I. Miralles, D. Castilla, C. Botella, I. Miralles, D. Castilla, and C. Botella, “Desarrollo de un Entorno Virtual para la Inducción Emocional,” in *CEUR Workshop Proceedings*, 2014, pp. 230–238. Accessed: Apr. 20, 2024. [Online]. Available: <https://producciocientifica.uv.es/documentos/5ecfad70299952687c554904?lang=en>
- [19] M. Ferrer-García, O. García-Rodríguez, J. Gutiérrez-Maldonado, I. Pericot-Valverde, and R. Secades-Villa, “Efficacy of virtual reality in triggering the craving to smoke: its relation to level of presence and nicotine dependence,” *Stud. Health Technol. Inform.*, vol. 154, pp. 123–127, 2010.
- [20] S. S. Khariwala *et al.*, “‘Cold turkey’ or pharmacotherapy: Examination of tobacco cessation methods tried among smokers prior to developing head and neck cancer,” *Head Neck*, vol. 41, no. 7, pp. 2332–2339, 2019, doi: 10.1002/hed.25708.
- [21] M. C. Fiore and H. K. Koh, “50th Anniversary of the Surgeon General’s Report on Smoking and Health: Where Are We Now?,” UCSF Smoking Cessation Leadership Center. Accessed: Feb. 22, 2022. [Online]. Available: <https://smokingcessationleadership.ucsf.edu/webinar/50th-anniversary-surgeon-generals-report-smoking-and-health-where-are-we-now>
- [22] S. Stein, “Best VR Headset of 2024,” CNET. Accessed: Apr. 20, 2024. [Online]. Available: <https://www.cnet.com/tech/gaming/best-vr-headset/>
- [23] K. Kanaiwa, “VR Escape Game.” Jul. 04, 2020. [Apple iOS]. Available: <https://apps.apple.com/us/app/vr-escape-game/id1516134033>

Supplementary Files

Figures

Participant flow diagram of the MindCotine pre to post 6-week smoking cessation intervention among Stanford Cancer Center patients.



N=9 Participant usage of MindCotine VR-MET program components during the study period.

