

“Just because it's hard, it doesn't mean that our patients can't benefit from it”: Evaluation of a Remote Patient Monitoring Program During the COVID-19 Pandemic

Rose Gunn, Shelby L. Watkins, Dave Boston, A. Gabriela Rosales, Stefan Massimino, Suparna Navale, Stephanie L. Fitzpatrick, John Dickerson, Rachel Gold, George Lee, Carmit K. McMullen

Submitted to: JMIR Formative Research
on: December 21, 2023

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 28

 Figures 29

 Figure 1..... 30

 Figure 2..... 31

 Multimedia Appendixes 32

 Multimedia Appendix 1..... 33

 Multimedia Appendix 2..... 33

 CONSORT (or other) checklists..... 34

 CONSORT (or other) checklist 0..... 34

“Just because it's hard, it doesn't mean that our patients can't benefit from it”: Evaluation of a Remote Patient Monitoring Program During the COVID-19 Pandemic

Rose Gunn¹ MA; Shelby L. Watkins¹ MPH; Dave Boston¹ MD; A. Gabriela Rosales² MS; Stefan Massimino² MS; Suparna Navale¹ PhD; Stephanie L. Fitzpatrick² PhD; John Dickerson² PhD; Rachel Gold^{1,2} MPH, PhD; George Lee³ MD; Carmit K. McMullen² PhD

¹OCHIN, Inc. Portland US

²Kaiser Permanente Center for Health Research Kaiser Permanente Portland US

³Asian Health Services Oakland US

Corresponding Author:

Rose Gunn MA
OCHIN, Inc.
PO Box 5426
Portland
US

Abstract

Background: Community health center (CHC) patients experience a disproportionately high prevalence of chronic conditions and barriers to accessing technologies that might support the management of these conditions. One such technology includes tools utilized for remote patient monitoring (RPM), the use of which surged during the COVID-19 pandemic.

Objective: The aim of this study was to assess how one CHC implemented an RPM program during the COVID-19 pandemic.

Methods: This retrospective case study used a mixed methods explanatory sequential design to evaluate one CHC's implementation of a suite of RPM tools during the COVID-19 pandemic. Analyses used EHR-extracted health outcomes data and semi-structured interviews with CHC staff and patients participating in the RPM program.

Results: The CHC enrolled 147 patients in a hypertension RPM program. After 6 months of RPM use, mean systolic blood pressure (BP) was 13.4 mmHg lower and mean diastolic BP 6.4 mmHg lower, corresponding with an increase in hypertension control (BP<140/90) from 33.3% of patients to 81.5%. Considerable effort was dedicated to standing up the program, reinforced by organizational prioritization of chronic disease management, and by a clinician who championed program implementation. Noted barriers to implementation of the RPM program were limited initial training, lack of sustained support, and complexities related to the RPM device technology.

Conclusions: While RPM technology holds promise for addressing chronic disease management, successful RPM program requires substantial investment in implementation support and technical assistance.

(JMIR Preprints 21/12/2023:55732)

DOI: <https://doi.org/10.2196/preprints.55732>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://preprints.jmir.org/preprint/55732>



Original Manuscript

Original Paper

“Just because it's hard, it doesn't mean that our patients can't benefit from it”:

Evaluation of a Remote Patient Monitoring Program During the COVID-19 Pandemic

Rose Gunn, MA¹; Shelby L. Watkins, MPH¹; Dave Boston, MD¹; A. Gabriela Rosales, MS²; Stefan Massimino, MS²; Suparna Navale, PhD; Stephanie L. Fitzpatrick, PhD²; John Dickerson, PhD²; Rachel Gold, PhD, MPH^{1,2}; George Lee, MD³; Carmit K. McMullen, PhD²

¹ OCHIN, Inc. Portland, Oregon, USA

² Kaiser Permanente Center for Health Research, Kaiser Permanente, Portland, Oregon, USA

³ Asian Health Services, Oakland, California, USA

Corresponding Author:

Rose Gunn, MA
OCHIN, Inc.
PO Box 5426
Portland, OR USA 97228-5426
Email: gunnr@ochin.org
Tel: 503.943.2512
Fax: 503.943.2501

ABSTRACT

Background. Community health center (CHC) patients experience a disproportionately high prevalence of chronic conditions and barriers to accessing technologies that might support the management of these conditions. One such technology includes tools utilized for remote patient monitoring (RPM), the use of which surged during the COVID-19 pandemic.

Objective. The aim of this study was to assess how one CHC implemented an RPM program during the COVID-19 pandemic.

Methods. This retrospective case study used a mixed methods explanatory sequential design to evaluate one CHC's implementation of a suite of RPM tools during the COVID-19 pandemic. Analyses used EHR-extracted health outcomes data and semi-structured interviews with CHC staff and patients participating in the RPM program.

Results. The CHC enrolled 147 patients in a hypertension RPM program. After 6 months of RPM use, mean systolic blood pressure (BP) was 13.4 mmHg lower and mean diastolic BP 6.4 mmHg lower, corresponding with an increase in hypertension control (BP<140/90) from 33.3% of patients to 81.5%. Considerable effort was dedicated to standing up the program, reinforced by organizational prioritization of chronic disease management, and by a clinician who championed program implementation. Noted barriers to implementation of the RPM program were limited initial training, lack of sustained support, and complexities related to the RPM device technology.

Conclusions. While RPM technology holds promise for addressing chronic disease management, successful RPM program requires substantial investment in implementation support and technical assistance.

Key Words: Patient-generated health data, telemedicine, telehealth, diabetes mellitus, hypertension, self-management, patient portal, implementation science

BACKGROUND

Chronic conditions including type 2 diabetes mellitus (DM) and hypertension (HTN) affect six in ten adults in the United States and are leading contributors to disability and mortality.[1, 2] Community

health centers (CHC) serve low-income populations whose members experience a disproportionately high prevalence of these conditions.[3] CHC patients also face barriers to accessing and using promising new technologies that can improve chronic disease management, such as telehealth and home monitoring devices.[4-6] Varying terminology is used for such technologies. Here, 'Remote Patient Monitoring' (RPM) describes patient-operated external biometric measurement devices with associated software and hardware to allow for patient-facing reporting and EHR integration, used to record personal health data in one location that is received by a provider elsewhere.[7]

RPM has been shown to benefit patients and their care teams by improving access to care, supporting clinical decision-making, and promoting chronic disease self-management.[8-13] Its use has also been associated with improved health outcomes, reduced emergency department visits and hospital admissions, and lower costs of care.[14-16] Despite these promising potential benefits, little has been reported on how best to implement RPM programs in primary care settings and what elements of RPM effectively engage patients and clinic staff.[12] This is particularly significant in health centers like CHCs, which serve a high volume of patients who are uninsured or Medicaid-insured for whom health disparities might be exacerbated by health centers' delayed adoption of digital health technologies.[17]

The Federal Communications Commission's (FCC) COVID-19 Telehealth Program distributed \$200 million under the Coronavirus Aid Relief and Economic Security (CARES) Act to help community-based health care organizations.[18] The program supported the provision of telehealth services and Wi-Fi connected devices, including those used for RPM. This provided a unique opportunity to understand how CHCs implemented an RPM program during a period when in-person care was greatly restricted.

OCHIN, a nonprofit health technology organization hosting a central instance of the OCHIN Epic electronic health record (EHR) system, received FCC COVID-19 Telehealth Program funds to distribute technology to support RPM to 24 health centers. The technology included Bluetooth-enabled glucometers, blood pressure (BP) cuffs, weight scales, pulse oximeters, and smartphones with one year of pre-paid wireless data. OCHIN offered the 24 participating health centers support with distributing these devices through a help desk, device setup and troubleshooting videos, printed guides for patients and clinic staff, and facilitation of a monthly peer-to-peer learning collaborative meeting. Health centers selected which of the offered devices to distribute, determined which of their patients would receive them, and decided whether the devices should be given to patients permanently or temporarily and then re-distributed. The RPM programs were implemented in different ways at the participating sites, one of which stood out as a success story. This manuscript describes implementation of the RPM program in this health center, including identified barriers and facilitators to RPM adoption, and patterns of improvement in participating patients' BP.

METHODS

We used a mixed methods explanatory sequential design to evaluate one CHC's implementation of an RPM program made available through the FCC COVID-19 Telehealth Program.[19] When we gathered information about the FCC grant program, we learned from informal interviews and formal program evaluation data that the program did not generally result in successful RPM adoption. Using a positive deviance case study design,[20, 21] we identified one participating CHC that experienced unusual success in distributing and gathering data from RPM devices. These analyses quantitatively evaluate the program's impact on BP and qualitatively explain that impact through interviews with CHC personnel and patients.

Study Setting

The setting was a single CHC health system that implemented an RPM program in fall 2020. This urban CHC system provides wraparound services (e.g., medical, dental, behavioral health) to over 50,000 patients, most of whom speak languages other than English, primarily Chinese dialects. Medicare/Medicaid and self-pay are the two most common forms of payment at this CHC.

Participants

Study participants include clinic personnel and patients who received RPM devices. Clinic personnel were recruited purposefully based on their knowledge of the RPM implementation effort. Patients eligible for this analysis were adults who received an RPM device as part of the FCC program and used it at least once to upload data to the EHR. Details about how the clinic provided devices to patients are given in the Results; in brief, clinicians had discretion to select patients for the RPM program. The RPM implementation team decided to prioritize distributing RPM devices to individuals with uncontrolled HTN, substantial barriers to care, no evidence of prior smartphone use, and no previous experience with virtual office visits.

Data Collection

Device distribution data collected by the CHC were used to confirm participants and dates of device disbursement and return, if available. BP measurement data were extracted from the CHC's EHR. Baseline BP measurements from RPM devices were collected from the time the patients received the devices through day seven. Seven days were added to the initial device distribution date to account for patients' initial learning curve using the RPM device. Subsequent device use data were collected at months 1, 3, and 6.

Qualitative data included semi-structured interviews with two clinicians and two patients participating in the RPM program at the CHC. See Appendix A and B for interview guides. Interviews were conducted from June 2022 to July 2022 and lasted 30-60 minutes. A clinician champion[22] who led and advocated for the RPM program identified staff and patients to interview based on their RPM involvement. Interviews were held virtually and audio-recorded with permission. Patient interviews were conducted through a professional interpreter. We also gathered background information by conducting informal interviews with OCHIN staff involved with administering and supporting the FCC program and by reviewing documents pertaining to program roll out. Recorded interviews were professionally transcribed, de-identified, and uploaded to NVivo (Release 1.7) for analysis.

Data Analysis

Quantitative data were analyzed descriptively to assess patterns of RPM use in the six months following adoption of the tools and patient characteristics associated with these patterns. Paired differences between BP measurements from the RPM devices were assessed using non-parametric Wilcoxon signed-rank tests because of the non-normality of the data. We assessed how the device's BP data changed at 0-7, 30-37, 90-97, and 182-189 days post-distribution.

Per explanatory sequential design, we assessed what qualitative data explained the quantitatively measured improvement in BP control associated with participation in the RPM program. Qualitative data were analyzed thematically, informed by the Consolidated Framework for Implementation Research (CFIR).[23-25] First, an initial code list was developed based on key CFIR constructs (e.g., process, inner setting, outer setting, intervention characteristics), with additional codes added inductively. Emerging themes were shared with the OCHIN FCC Program staff, CHC staff interviewees, and the full cross-disciplinary research team to confirm early findings and foster

interpretation through diverse perspectives. Results are conveyed using a case study reporting approach characterized by chronological narrative framing.

Ethical Considerations

The Institutional Review Board at Kaiser Permanente Northwest approved this study (#00000405). All methods were carried out in accordance with relevant guidelines and regulations. BP measurement data were extracted from the OCHIN Epic EHR, as allowable by an agreement with OCHIN member health centers. Informed verbal consent was obtained from interview participants who were notified of their right to refuse to participate and the study team's procedures for deidentifying data. Patient participants were provided with a \$50 gift card in appreciation of their time. The CHC received \$400 as an impact payment in recognition of their participation in the evaluation of their RPM program.

RESULTS

The CHC in this case study received 50 Bluetooth-enabled blood pressure cuffs and 50 iPhones with unlimited wireless data for one year and implementation support as described above. They opted to use a lending model of device distribution to maximize patient reach.


Device Use & Blood Pressure

Blood pressure readings were received from 147 patients, indicating that each of the 50 devices was loaned to an average of 3 patients over the course of the RPM intervention. Patients whose records included device return dates used their devices for an average of 195 days. The CHC provided the study team with a list of patients who received devices (n=152). This list was cross-referenced with

EHR data indicating whether the patient used the device. Five patients did not have evidence of device use, leaving 147 patients in the sample. Among the 147 patients, 63% were female, 98% were Asian, and none were Hispanic. The mean age was 70 and 90% of the patients were at or below the federal poverty level.


Average BP steadily dropped from baseline to six-month follow-up, with a reduction of systolic (-13.4 mmHg) and diastolic (-6.4 mmHg) measurements among those having BP measurements at six-month follow-up (Figure 1).

Figure 1. Distribution of mean systolic and diastolic by months.



This translated into more than doubling of hypertension control (BP<140/90). At baseline, 33.3% of patients with BPs at baseline had controlled hypertension, and 81.5% of patients with BPs at six-month follow up had controlled hypertension (Figure 2).

Figure 2. Percent change in hypertension control.



RPM Program Implementation

Insights from qualitative data about RPM program implementation illuminated how the clinic successfully implemented RPM. When OCHIN received the FCC grant, it broadly communicated the RPM opportunity across its membership and invited health centers to submit proposals to participate in the RPM program. Proposals were scored based on several domains, including prioritizing health centers and patients most disproportionately affected by COVID-19.

Leaders at the case study CHC convened a team to oversee implementation of the RPM program. This 'RPM team' consisted of four clinicians with an interest in telehealth who agreed to pilot the

program, and members of the CHC's health coaching, quality improvement (QI), and health information technology teams. The team met weekly for three months to plan how to implement the RPM program. The QI director developed meeting agendas, tracked task status, and enforced timelines.

The RPM team viewed the program as a 'launch point' for more broadly shifting their approach to care delivery from episode- or encounter-based to a continuous model. Their envisioned new model involved focusing on how patients were doing between visits and enabling care teams to support chronic condition management in a manner convenient for patients. The team saw the value of RPM as part of this vision but knew it required significant planning. Implementation was complicated by the concurrent rollout of a patient EHR portal that did not include the languages predominantly spoken by this CHC's patients. The RPM team considered which functions of the portal could be useful to RPM patients who were unable to read the available languages given that portal activation was required to send BP data to the care team. The team decided to prioritize distributing RPM devices to patients with multiple challenges to HTN control, as described previously. A clinician reflected,

I was very skeptical because I was like, "It is going to be so hard to do it with our patients." And at the same time, though, there's this kind of like, "Well, if we're not going to try it, then who will try it?" Just because it's hard, it doesn't mean that our patients can't benefit from it.

[MD1]

Given the unique needs of the population they were trying to reach, the CHC did not use the implementation support materials provided by OCHIN. Instead, they developed printed materials to

support device adoption and use in different languages and literacy levels, and culturally tailored hypertension educational materials. One of the clinicians piloting the program shared,

There's a language barrier for our patients, and so we had to figure out what is the most simple document that we can give people that's very pictorial, has a lot of arrows, and not a lot of text. We just came up with a lot of the materials in-house. I think our team was really used to coming up with teaching materials for staff, and so they helped us to make it even more simple. [MD1]

The team also developed device distribution and return workflows (i.e., handing out and receiving devices back), and an EHR-based dashboard that tracked recruitment, interactions about device troubleshooting, and coaching calls, as well as reports on incoming device data. The RPM team was concerned that patients would not have enough IT support and/or would experience substantial language and technology literacy barriers, so they trained the clinic's health coaches, who are also medical assistants, to support patients as required. The CHC's health IT team provided backup support.

Members of the RPM team with QI roles developed systems for tracking program metrics, which alerted CHC leaders to their initial investment in the program and to the resources likely needed to sustain it. This was primarily done using structured EHR data elements, including when the devices were distributed and returned, how many coaching calls each patient received, and how much troubleshooting each patient required.

Clinicians involved in participating in the RPM program initially reached out to eligible patients to offer the devices; if interest was expressed, a health coach scheduled an hour-long appointment to set

up and sync the devices and provide patient education on device use. Health coaches supported patients with portal sign-up, then synced the blood pressure cuff to the provided iPhone using the iHealth app, which interfaced with the Apple Health app that interfaced with the patient portal. Blood pressure data automatically went through the patient portal directly to the EHR's BP data, which could be reviewed by care team members. A member of the IT team developed reports showing average weekly and monthly BP over time. Clinicians and health coaches met regularly to review the BP readings sent from the RPM devices to the EHR, and made plans for patient follow-up, which included potential medication adjustments and virtual appointments with a health coach. Patients were eligible for health coaching if they demonstrated ability to use the devices by having at least five BP readings submitted from the devices.

The CHC experienced notable barriers to implementing the RPM program, particularly given the many competing demands related to clinics' COVID-19 response. Care teams faced challenges in setting up and syncing devices; patients reported difficulty in maintaining synced connections and staying logged in. After some testing and refinement, CHC leaders were pleased with the promising results of the RPM program, a sentiment echoed by the patients. One clinician shared,

"We had pretty fantastic results in terms of people's blood pressure becoming under control. We also were able to pick up on things that we never thought we would. I was alerted because [one of my patients] had a very low heart rate, and I couldn't figure out why, so they ended up calling him. It turned out he was prescribed a different medication by another physician that was a beta-blocker that lowers your heart rate. Another one of my patients had been fairly well controlled for a while and then all of a sudden had some spikes in his blood pressure. Our health coach called him, and it turned out he was having a gout attack."

And so, our nurse said, ‘You should come in and be treated.’ So he got treated. But we found out through blood pressure. So, I think it definitely works well.” [MD2]

As care teams noticed improvements in patients’ BP, the CHC added two RPM team members to support the program: a digital health advocate to work with patients on device setup and technical support and a program manager to support project management of the RPM activities. Based on the organization’s prioritization of RPM, their grants department wrote a grant proposal to fund the digital health advocate position. They received multiple grants to support the digital health advocate role but also acknowledged the lack of sustainability of grant-funded positions.

Patients said they felt prepared to use the devices after the initial visit with the health coach and were able to call the coach for support with their devices. One patient shared her and her husband’s experience using the devices,

When we first received the devices, the nurse taught me how to use the device in person. And they know that this technology is a little bit difficult for us, so they made sure that we learned the steps and how to use it. And then after a couple of days, my husband forgot how to use it. And then I called the clinic and asked the nurse. And the nurse taught us on the phone again until we all felt comfortable using the devices. [Patient 1]

Health center clinicians noted that the drop-out rate for the program was high, as approximately 30% of patients who initially said they wanted a device chose not to participate after the initial related visits. Despite implementation and technology barriers, however, the CHC was satisfied with their decision to prioritize RPM and are expanding the RPM program to more care teams in their health system and considering how to sustain the program over time.

DISCUSSION

Among adult patients selected to receive a home BP cuff and smartphone, BP reductions were demonstrated during device use, likely due to a combination of factors including increased staff attention to lowering BP, educating patients how to properly use the devices, and motivating patients to actively participate in RPM and hypertension management. In addition to measuring BP, motivated patients may have adhered better to lifestyle recommendations and medication prescriptions. Neither the relative contributions of these and other factors, nor long-term durability of hypertension control can be accurately assessed from these data, but these would be helpful to analyze in future investigations.

Limited prior research has assessed factors affecting RPM implementation and use in CHCs.[26] These results indicate that successful program implementation in CHCs may involve not just technology and training but also substantial upfront planning and commitment, driven by a clinical champion or leader. This may include establishing an RPM team, developing related workflows, and creating data reports to monitor clinical care decisions, in collaboration with patients. This may be challenging for health centers that lack a robust staffing model with QI, EHR support, health coaching and other resources available to support pilot testing and material development. For such clinics, a more modest pilot phase might be more feasible, but may not yield the same clinical outcomes.

Prior research showed the potential for technology innovations such as RPM to exacerbate health disparities,[27-29] exemplified by inequitable expansion of RPM technologies during the COVID-19 pandemic.[30, 31] It is therefore important to document successes like the CHC described here, which saw positive outcomes from its RPM program despite its focus on patients facing considerable

barriers.[32, 33] Elements of its success included developing multi-lingual materials and utilizing uncomplicated RPM devices to support patient accessibility. Maintaining regular outreach with participating patients (i.e., co-intervention) also likely helped, as has been demonstrated by others. [13, 34-36] Future research is needed to examine the impact of different implementation strategies on the uptake and sustained use of RPM devices, particularly among patients facing barriers to care.

Limitations

A single case study limits the generalizability of findings; the case study approach was used here to provide a comprehensive view of this multifaceted topic.[37] Also, retrospective studies are limited in that they provide a snapshot in time after an intervention has been implemented but do not provide a detailed understanding of implementation in real time. This potential was mitigated here by collecting data as close to the project end date as possible to optimize the timeliness of reflections about implementation. Finally, because this primarily is a qualitative evaluation, quantitative analysis was limited to descriptive statistics. We compared average BP values of patients with BP data at the various timepoints rather than tracking individual patients. This likely biased our results by excluding those patients who were less actively participating. And since there was no usual care control group with which to compare effectiveness of RPM on BP reduction, more robust statistical analysis could not be performed.

CONCLUSION

The experience of one CHC indicates that while RPM technology holds promise for addressing chronic disease management among safety net patients, successful and equitable RPM implementation will require substantial investment in implementation support and technical assistance. Further exploration of these issues in a prospective evaluation is currently underway.

Acknowledgements

The authors appreciate the participating health centers and are grateful for their time and expertise. This study was supported by the National Institute of Nursing Research (R01NR020305). The study sponsor did not play a role in the study design, collection, analysis, and interpretation of data, or the submission of this manuscript. The content provided is solely the responsibility of the authors and does not necessarily represent the official views of the funders. This work was conducted with the Accelerating Data Value Across a National Community Health Center Network (ADVANCE) Clinical Research Network (CRN). ADVANCE is led by OCHIN in partnership with Health Choice

Network, Fenway Health, and Oregon Health & Science University. ADVANCE is funded through the Patient-Centered Outcomes Research Institute (PCORI), contract number RI-OCHIN-01-MC.

Data Availability

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Authors' Contributions

CM, DB, R Gold, SF, and JD conceived and designed the study. R Gunn and SW collected qualitative data and analyzed it with CM. SN extracted EHR data; AGR, JD, and SF conducted the quantitative analysis. GL validated qualitative and quantitative findings and served as case study advisor. All authors played a role in result interpretation and preparation of the manuscript. All authors approved the final version of this manuscript.

Conflicts of Interest

None declared.

REFERENCES

1. National Center for Chronic Disease Prevention and Health Promotion (NCCHPHP). Chronic Diseases in America. Centers for Disease Control and Prevention. 2022. <https://www.cdc.gov/chronicdisease/resources/infographic/chronic-diseases.htm>. Accessed 15 June 2023.
2. Watson KB CS, Loustalot F, et al. Chronic Conditions Among Adults Aged 18–34 Years — United States, 2019. Centers for Disease Control and Prevention. MMWR Morb Mortal Wkly Rep. 2022. ;71:964–970. [DOI: 10.15585/mmwr.mm7130a3]
3. Corallo B, Proser M, Nocon R. Comparing Rates of Multiple Chronic Conditions at Primary Care and Mental Health Visits to Community Health Centers Versus Private Practice

- Providers. J Ambul Care Manage. 2020;43(2):136-47. [DOI: 10.1097/JAC.0000000000000324]
4. Shi L, Lee DC, Haile GP, Liang H, Chung M, Sripipatana A. Access to Care and Satisfaction Among Health Center Patients With Chronic Conditions. J Ambul Care Manage. 2017;40(1):69-76. [DOI: 10.1097/JAC.0000000000000153]
 5. Weigel G, Ramaswamy, A., Sobel, L., Salganicoff, A., Cubanski, J., Freed, M. . Opportunities and Barriers for Telemedicine in the U.S. During the COVID-19 Emergency and Beyond. Kaiser Family Foundation. 2020. <https://www.kff.org/womens-health-policy/issue-brief/opportunities-and-barriers-for-telemedicine-in-the-u-s-during-the-covid-19-emergency-and-beyond/>. Accessed 15 June 2023.
 6. Lee G, Chang A, Pal A, Tran TA, Cui X, Quach T. Understanding and Addressing the Digital Health Literacy Needs of Low-Income Limited English Proficient Asian American Patients. Health Equity. 2022;6(1):494-9. [DOI: 10.1089/heq.2022.0045]
 7. The Office of the National Coordinator for Health Information Technology (ONC). Telemedicine and Telehealth. 2023. <https://www.healthit.gov/topic/health-it-health-care-settings/telemedicine-and-telehealth>. Accessed 15 June 2023.
 8. Omoloja A, Vundavalli S. Patient generated health data: Benefits and challenges. Curr Probl Pediatr Adolesc Health Care. 2021;51(11):101103. [DOI: 10.1016/j.cppeds.2021.101103]
 9. Vicente MA, Fernández C, Guilabert M, Carrillo I, Martín-Delgado J, Mira JJ, et al. Patient Engagement Using Telemedicine in Primary Care during COVID-19 Pandemic: A Trial Study. Int J Environ Res Public Health. 2022;19(22). [DOI: 10.3390/ijerph192214682]
 10. Salehi S, Olyaeemanesh A, Mobinizadeh M, Nasli-Esfahani E, Riazi H. Assessment of remote patient monitoring (RPM) systems for patients with type 2 diabetes: a systematic review and meta-analysis. J Diabetes Metab Disord. 2020;19(1):115-27. [DOI:

10.1007/s40200-019-00482-3]

11. Bunch A, Ardila F, Castaño R, Quiñonez S, Corzo L. Through the Storm: Automated Peritoneal Dialysis with Remote Patient Monitoring during COVID-19 Pandemic. *Blood Purif.* 2021;50(3):279-82. [DOI: 10.1159/000511407]
12. Noah B, Keller MS, Mosadeghi S, Stein L, Johl S, Delshad S, et al. Impact of remote patient monitoring on clinical outcomes: an updated meta-analysis of randomized controlled trials. *NPJ Digit Med.* 2018;1:20172. [DOI: 10.1038/s41746-017-0002-4]
13. Sheppard JP, Tucker KL, Davison WJ, Stevens R, Aekplakorn W, Bosworth HB, et al. Self-monitoring of Blood Pressure in Patients With Hypertension-Related Multi-morbidity: Systematic Review and Individual Patient Data Meta-analysis. *Am J Hypertens.* 2020;33(3):243-51. [DOI: 10.1093/ajh/hpz182]
14. Taylor ML, Thomas EE, Snoswell CL, Smith AC, Caffery LJ. Does remote patient monitoring reduce acute care use? A systematic review. *BMJ Open.* 2021;11(3):e040232. [DOI: 10.1136/bmjopen-2020-040232]
15. De Guzman KR, Snoswell CL, Taylor ML, Gray LC, Caffery LJ. Economic Evaluations of Remote Patient Monitoring for Chronic Disease: A Systematic Review. *Value Health.* 2022;25(6):897-913. [DOI: 10.1016/j.jval.2021.12.001]
16. Niu B, Mukhtarova N, Alagoz O, Hoppe K. Cost-effectiveness of telehealth with remote patient monitoring for postpartum hypertension. *J Matern Fetal Neonatal Med.* 2022;35(25):7555-61. [DOI: 10.1080/14767058.2021.1956456]
17. Richardson S, Lawrence K, Schoenthaler AM, Mann D. A framework for digital health equity. *NPJ Digit Med.* 2022;5(1):119. [DOI: 10.1038/s41746-022-00663-0]
18. Federal Communications Commission (FCC). COVID-19 Telehealth Program - Frequently Asked Questions (FAQs) 2020. <https://www.fcc.gov/covid-19-telehealth-program-frequently-asked-questions-faqs>. Accessed 20 March 2023.

19. Fetters MD, Curry LA, Creswell JW. Achieving integration in mixed methods designs- principles and practices. *Health Serv Res.* 2013;48(6 Pt 2):2134-56. [DOI: 10.1111/1475-6773.12117]
20. Rose AJ, McCullough MB. A Practical Guide to Using the Positive Deviance Method in Health Services Research. *Health Serv Res.* 2017;52(3):1207-22. [DOI: 10.1111/1475-6773.12524]
21. Bradley EH, Curry LA, Ramanadhan S, Rowe L, Nembhard IM, Krumholz HM. Research in action: using positive deviance to improve quality of health care. *Implementation Science.* 2009;4(1):25. [DOI: 10.1186/1748-5908-4-25]
22. Morena AL, Gaias LM, Larkin C. Understanding the Role of Clinical Champions and Their Impact on Clinician Behavior Change: The Need for Causal Pathway Mechanisms. *Front Health Serv.* 2022;2:896885. [DOI: 10.3389/frhs.2022.896885]
23. Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for the analysis of qualitative data in multi-disciplinary health research. *BMC Med Res Methodol.* 2013;13:117. [DOI: 10.1186/1471-2288-13-117]
24. Braun V, Clarke V, Hayfield N, Terry G. Thematic Analysis. In: Liamputtong P, editor. *Handbook of Research Methods in Health Social Sciences.* Singapore: Springer Singapore; 2019. p. 843-60.
25. Damschroder L, Hall C, Gillon L, Reardon C, Kelley C, Sparks J, et al. The Consolidated Framework for Implementation Research (CFIR): progress to date, tools and resources, and plans for the future: *Implement Sci.* 2015 Aug 20;10(Suppl 1):A12. [DOI: 10.1186/1748-5908-10-S1-A12. eCollection 2015.
26. Yi SS, Tabaei BP, Angell SY, Rapin A, Buck MD, Pagano WG, et al. Self-blood pressure monitoring in an urban, ethnically diverse population: a randomized clinical trial utilizing the electronic health record. *Circ Cardiovasc Qual Outcomes.* 2015;8(2):138-45. [DOI: 10.1161/

CIRCOUTCOMES.114.000950]

27. Veinot TC, Mitchell H, Ancker JS. Good intentions are not enough: how informatics interventions can worsen inequality. *J Am Med Inform Assoc.* 2018;25(8):1080-8. [DOI: 10.1093/jamia/ocy052]
28. Brewer LC, Fortuna KL, Jones C, Walker R, Hayes SN, Patten CA, et al. Back to the Future: Achieving Health Equity Through Health Informatics and Digital Health. *JMIR Mhealth Uhealth.* 2020;8(1):e14512. [DOI: 10.2196/14512]
29. Agarwal S, Simmonds I, Myers AK. The Use of Diabetes Technology to Address Inequity in Health Outcomes: Limitations and Opportunities. *Curr Diab Rep.* 2022;22(7):275-81.[DOI: 10.1007/s11892-022-01470-3]
30. Price JC, Simpson DC. Telemedicine and Health Disparities. *Clin Liver Dis (Hoboken).* 2022;19(4):144-7. [DOI: 10.1002/cld.1171]
31. Ortega G, Rodriguez JA, Maurer LR, Witt EE, Perez N, Reich A, et al. Telemedicine, COVID-19, and disparities: Policy implications. *Health Policy Technol.* 2020;9(3):368-71. [DOI: 10.1016/j.hlpt.2020.08.001]
32. Nouri S, Khoong EC, Lyles CR, Karliner L. Addressing Equity in Telemedicine for Chronic Disease Management During the Covid-19 Pandemic. *NEJM Catalyst non-issue content.* 2020;1(3). [DOI: 10.1056/CAT.20.0123]
33. Bailey JE, Gurgol C, Pan E, Njie S, Emmett S, Gatwood J, et al. Early Patient-Centered Outcomes Research Experience With the Use of Telehealth to Address Disparities: Scoping Review. *J Med Internet Res.* 2021;23(12):e28503. [DOI: 10.2196/28503]
34. Tucker KL, Sheppard JP, Stevens R, Bosworth HB, Bove A, Bray EP, et al. Self-monitoring of blood pressure in hypertension: A systematic review and individual patient data meta-analysis. *PLoS Med.* 2017;14(9):e1002389. [DOI: 10.1371/journal.pmed.1002389]
35. Beger C, Rüegger D, Lenz A, Wagner S, Haller H, Schmidt-Ott KM, et al. Blood pressure

- dynamics during home blood pressure monitoring with a digital blood pressure coach-a prospective analysis of individual user data. *Front Cardiovasc Med*. 2023;10:1115987. [DOI: 10.3389/fcvm.2023.1115987]
36. Hellem AK, Whitfield C, Casetti A, Robles MC, Dinh M, Meurer W, et al. Engagement in Self-measured Blood Pressure Monitoring Among Medically Underresourced Participants (the Reach Out Trial): Digital Framework Qualitative Study. *JMIR Cardio*. 2023;7:e38900. [DOI: 10.2196/38900]
37. Crowe S, Cresswell K, Robertson A, Huby G, Avery A, Sheikh A. The case study approach. *BMC Medical Research Methodology*. 2011;11(1):100. [DOI: 10.1186/1471-2288-11-100]


Abbreviations

Health Information Technology (HIT), Electronic Health Record (EHR), Community Health Center (CHC), Remote Patient Monitoring (RPM)


Supplementary Files

Figures

Distribution of mean systolic and diastolic by months.



Percent change in hypertension control.



Multimedia Appendixes

Clinic Staff Interview Guide.

URL: <http://asset.jmir.pub/assets/88b2c90f6aab05af3d387903ed7e91ba.docx>

Patient Interview Guide.

URL: <http://asset.jmir.pub/assets/28e3987b25f2f389d37f5d6084881e9a.docx>



CONSORT (or other) checklists

COREQ Checklist.

URL: <http://asset.jmir.pub/assets/ccd4545f620d0b5a183b409397c4aa7c.pdf>