

Factors associated with risky drinking decisions in a virtual reality alcohol prevention simulation: A structural equation model

Robert Hrynyschyn, Julie Dalgaard Guldager, Daniel Schulze, Patricia Bianca Lyk, Gunver Majgaard, Christiane Stock

Submitted to: JMIR XR and Spatial Computing (JMXR)
on: January 09, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 24

 Figures 25

 Figure 1..... 26

 Figure 2..... 27

 Multimedia Appendixes 28

 Multimedia Appendix 1..... 29

 Multimedia Appendix 2..... 29

 Multimedia Appendix 3..... 29

 TOC/Feature image for homepages 30

 TOC/Feature image for homepage 0..... 31

Factors associated with risky drinking decisions in a virtual reality alcohol prevention simulation: A structural equation model

Robert Hrynyschyn^{1,2} MPH; Julie Dalgaard Guldager^{3,4} PhD; Daniel Schulze⁵ PhD; Patricia Bianca Lyk⁶ PhD; Gunver Majgaard⁶ PhD; Christiane Stock^{1,4} PhD

¹Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Health and Nursing Science Berlin DE

²Leibniz ScienceCampus Bremen Digital Public Health Bremen DE

³Research Department University College South Denmark Haderslev DK

⁴Unit for Health Promotion Research Department of Public Health University of Southern Denmark Esbjerg DK

⁵Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Biometry and Clinical Epidemiology Berlin DE

⁶The Maersk Mc-Kinney Moller Institute Game Development and Learning Technology University of Southern Denmark Odense DK

Corresponding Author:

Robert Hrynyschyn MPH

Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Institute of Health and Nursing Science

Augustenburger Platz 1

Berlin

DE

Abstract

Background: Risky alcohol consumption among adolescents is a significant public health concern in most Western countries. Efforts to tackle this challenge have been successful through skills training to deal with social pressures. However, interventions of this nature require significant resources. Technological solutions, such as virtual reality, offer advantages as they enable immersive experiences that replicate real-life scenarios. However, a persistent question pertains to the fidelity of real-world behaviors within virtual environments.

Objective: This study is exploratory and aims to ascertain if the established drinking motives and factors for risky drinking decisions are transferrable to the virtual environment in the simulation game VR FestLab and to uncover determinants linked to risky drinking decisions within the simulation.

Methods: Data from 161 Danish participants (aged 14-18 years) in a longitudinal intervention study were analysed. Confirmatory factor analysis and structural equation modelling were utilized to examine diverse latent variables (including alcohol and drug use, sensation seeking, refusal skills and knowledge & communication skills) in relation to virtual risk decisions.

Results: The results indicate that individuals with prior alcohol experiences exhibited a fourfold increased chance (odds ratio = 4.32, $p = .01$) of engaging in virtual risk decisions within the simulated environment. However, other latent variables did not demonstrate significant associations with virtual risk decisions. Neither gender nor age exhibited moderation effects on the outcome.

Conclusions: The immersive and lifelike properties of VR partially reflected risk-related decisions. However, it remains unclear which factors favour the mapping of real-world behaviours in virtual simulations. Therefore, future research should address the mechanisms underlying behavioural dynamics in virtual simulations and explore the translation of virtual behaviours into real behaviours to gain a comprehensive understanding of the potential of virtual simulations for alcohol prevention.

(JMIR Preprints 09/01/2024:56188)

DOI: <https://doi.org/10.2196/preprints.56188>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

✓ **No, I do not wish to publish my submitted manuscript as a preprint.**

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/preprint/56188>



Original Manuscript

Original Paper

Factors associated with risky drinking decisions in a virtual reality alcohol prevention simulation: A structural equation model

Abstract

Background: Risky alcohol consumption among adolescents is a significant public health concern in most Western countries. Various motives and factors (e.g. sensation seeking, gender, reduced self-efficacy) are known in the literature that are associated with risky drinking decisions in real-life. Efforts to tackle risky drinking decisions in real-life have been successful through skills training to deal with social pressures. However, interventions of this nature require significant resources. Technological solutions, such as virtual reality, offer advantages as they enable immersive experiences that replicate real-life scenarios. However, a persistent question pertains to the fidelity of real-world behaviors within virtual environments.

Objective: This study is exploratory and aims to ascertain if the established drinking motives and factors for risky drinking decisions are transferrable to the virtual environment in the simulation game VR FestLab and to uncover determinants linked to risky drinking decisions within the simulation.

Methods: The study analyzed data from the intervention arm of a cluster-randomized study of 161 Danish students aged 14-18 years who tested the virtual alcohol prevention simulation VR FestLab. At baseline and before playing VR FestLab, independent variables such as age, gender, alcohol consumption, use of other drugs, sensation seeking, drinking refusal skills, knowledge of blood alcohol concentration and refusal communication skills were recorded. The dependent variable, virtual risk decisions, was measured immediately after the game play. Confirmatory factor analysis and structural equation modeling were used to examine the latent variables in relation to virtual risk decisions. Moderation analyses for age and gender in relation to the latent characteristics and the primary outcome were also conducted.

Results: The data indicate that 73.9% (n= 119) of the participants have engaged in binge drinking at least once in their lifetime. The confirmatory factor analysis demonstrated a good fit of the items for their respective constructs, and therefore, they were adopted without modification in the structural equation model. The data suggests that individuals with prior alcohol experience are four times more likely to engage in virtual risk decisions within the simulated environment (OR = 4.31 (CI = 1.70-10.84), $P = .01$). Knowledge/Awareness of blood alcohol concentration was associated with a lower chance to engage in virtual risk decisions (OR = 0.32 (CI = 0.11-0.93), $P = 0.04$). However, no significant associations were found between virtual risk decisions and other latent variables. Gender or age did not moderate the associations.

Conclusions: The immersive and lifelike properties of VR partially reflected risk-related decisions. However, it remains unclear which factors favor the mapping of real-world behaviors in virtual simulations. Therefore, future research should address the mechanisms underlying behavioral dynamics in virtual simulations and explore the translation of virtual behaviors into real behaviors to gain a comprehensive understanding of the potential of virtual simulations for alcohol prevention.

Keywords: alcohol; prevention; virtual reality; risk behavior; structural equation model

Introduction

Alcohol prevention continues to represent a pertinent public health concern worldwide [1]. Despite witnessing a reduction on a global scale since the establishment of the Millennium Development Goals in 1995, alcohol consumption still contributes to 5% of disability-adjusted life years and 5% of total fatalities [2]. Furthermore, harmful alcohol consumption is pivotal in over 200 diseases and injury conditions [3]. Notably, alcohol prevention remains significant, particularly among adolescents, as indicated by the latest European School Survey Project on Alcohol and Other Drugs report, which reveals that approximately 80% of 15-16 year old students have experimented with alcohol at least once [4]. Denmark surpasses the European average, with roughly 92% of adolescents having consumed alcohol at least once in their lifetimes [4].

To effectively deter premature substance use among adolescents, targeted alcohol prevention programs prove indispensable [5]. In this context, it is important to consider the motives underlying risky consumption patterns among adolescents. A systematic review conducted by Adan et al. [6] reveals an association between risky drinking behavior and specific personality traits. For instance, binge drinking is correlated with increased impulsivity and sensation seeking [6], while Stautz et al. [7] and Percy et al. [8] established an association between high sensation seeking and an increased likelihood of heavy episodic drinking. Other studies have identified binge drinking as particularly linked to male gender and reduced self-efficacy [9]. Moreover, engagement in other legal or illegal substances, apart from alcohol, correlates with increased risk behavior. Creamer et al. [10] demonstrated that using various tobacco products corresponds to heightened risk behavior. Additionally, a correlation exists between cannabis use, drinking-related risk behavior [11], and the frequency of alcohol-related consequences [12].

Given the known drinking motives among adolescents, prevention programs anchored in concepts like inoculation theory [13] and social learning theory [14] advocate for skill enhancement that strengthens resilience against social influences, such as peer pressure for risky behavior. Research involving life skills training in educational institutions underscores the value of cultivating social resistance skills and broader personal and social competencies, leading to decreased cigarette use and enhanced anti-drinking attitudes [15]. Moreover, participants demonstrate higher substance use awareness and skill-related knowledge [15]. Effective refusal communication skills and risk-related knowledge are crucial components for enhancing the personal and social skills of young people. Refusal communication, which involves the ability to say 'no' to substance offers, is essential in reducing substance use and risky behavior. Several studies have demonstrated that individuals with strong refusal skills regarding alcohol misuse exhibit reduced abusive alcohol use [16-18], possess greater knowledge of alcohol misuse prevention, are less susceptible to peer pressure, and have better internal health control [19]. However, in addition to communication skills, it is also crucial for young people to have knowledge about substance use and its effects on the body to assess and avoid risks. Individuals with greater knowledge of substance-related topics may be better equipped to handle risky situations, potentially reducing the likelihood of substance abuse. Various studies have explored the relationship between substance-related knowledge and the reduction of risk behaviors, such as alcohol consumption. Teesson, Newton [20] conducted a cluster-randomized study in schools and found that combining digital prevention programs that increased alcohol-related knowledge resulted in a reduction in binge drinking. Hasking and Schofield [21] demonstrated that health and alcohol knowledge can strengthen the intention-behavior relationship. Individuals with more alcohol-related knowledge and experience are likely to be better informed about the consequences. Conversely, better-informed adolescents are likely to feel better prepared to minimize the risks of alcohol consumption [21]. Padget, Bell [22] discovered that increased awareness of the detrimental effects of alcohol on the brain resulted in improved perceptions of harm and subsequent attitudes of alcohol aversion. These improvements had a significant impact on the intention not to use alcohol, but they did not result in a significant reduction in short-term alcohol consumption. Therefore, it is important

to note that knowledge about alcohol may be only one of many factors that can influence risk-taking decisions. Risk behaviors are frequently caused by multiple factors, and knowledge about alcohol may only have a partial impact on the development or absence of such behaviors.

While traditional skills training involving rational alcohol consumption often relied on labor-intensive and costly in-person role-playing, contemporary technological solutions such as virtual reality (VR) have emerged. An intrinsic advantage of VR lies in its capacity to deliver an immersive encounter that faithfully mimics real-life scenarios. By replicating authentic situations, VR prompts participants to enact genuine behaviors within virtual environments [23, 24]. Whereas studying real-world behaviors within experimentally controlled settings posed challenges, VR now allows one to scrutinize behaviors within genuine settings and uncover determinants of behavioral intentions [25].

The convergence between VR and real-life behavior might be attributed to presence and immersion. Individuals immersed in VR experiences can subjectively experience a sense of “being there” in the virtual realm. This sense of presence fosters more authentic and realistic behavioral responses akin to real-world conduct [26]. Alcañiz et al. [27] reported that comparable neural mechanisms can be triggered in individuals immersed in a virtual world, paralleling experiences in the physical world. Additionally, the increasing realism and interactivity of VR technology play a role. As VR systems advance, they provide heightened sensory input and feedback, including lifelike visuals, haptic responses, and precise motion tracking. These immersive, lifelike elements contribute to greater congruence between VR and real-life behaviors [28]. Moreover, psychological factors such as social presence and adherence to social norms influence VR behavior, as people replicate real-world behaviors due to a sense of social presence and the desire to conform, even within virtual contexts [29].

The immersive, authentic portrayal of virtual environments implies that genuine risk behaviors are likely to manifest in these settings. This conjecture is supported by the findings that, for example, children exhibiting higher risk behavior in road traffic replicate this behavior in a virtual cycling simulation [30].

Nonetheless, reviews indicate that VR’s role in substance use prevention remains still limited [31, 32]. Thus, the co-created development of a virtual alcohol prevention simulation (VR FestLab) was developed in 2020 [33]. The VR FestLab application, an educational game simulation, aims to enhance the refusal self-efficacy of adolescents aged 15-18 who experience social pressure to consume alcohol. Given the scarceness of VR-based alcohol prevention applications [31, 32], this study is exploratory and aims to ascertain if the established drinking motives and factors for risky drinking decisions are transferrable to VR FestLab. A structural equation model was employed to uncover determinants linked to risky drinking decisions within the simulation. To this end, the following hypotheses were formulated for testing:

1. Higher sensation seeking is associated with increased virtual risk decisions in the simulation.
2. Enhanced knowledge/awareness of blood alcohol concentration, refusal communication skills, and drinking refusal skills are linked to reduced virtual risk decisions in the simulation.
3. Prior alcohol experiences are associated with increased virtual risk decisions in the simulation.

Methods

Study design

The data were collected as part of a longitudinal study that investigated the efficacy of the VR FestLab application. The comprehensive procedure and outcomes of the primary study can be found elsewhere [34]. To achieve this objective, a total of 13 Danish schools were allocated in a 1:1 ratio to either the intervention or control group. The schools assigned to the intervention group engaged with the VR FestLab application, whereas those in the control group played the VR game First Steps (Meta Platforms Inc., Menlo Park, USA). The dataset in this current study is derived from the seven schools that were selected as intervention schools only. Data collection transpired within the school

premises between August and December 2020 and then again from April to May 2021, the latter period was necessitated by COVID-19 restrictions.

Participants

To enroll adolescents aged between 14 and 18, initial contact was established with the administrations of the seven schools through email. Once the school administrations granted their approval, the study's objective was presented to the respective classes, emphasizing the confidential and anonymous nature of data collection. In accordance with the principles outlined in the Declaration of Helsinki, all students were duly informed that their participation was voluntary, and they gave written consent prior to their involvement. The study adhered to Danish standards for the ethical conduct of scientific studies and was approved by the Research Ethics Committee of the University of Southern Denmark in March 2020 (case no 20/5348). In accordance with the Ethics Committee of the University of Southern Denmark, parental consent was waived as it is only mandatory for adolescents under the age of 15 in Denmark. For the original study [34], a sample size calculation was performed using STATA 15 with a two-sample t-test. This resulted in a sample size of 135 individuals for the control group and 135 individuals for the intervention group to yield an intervention effect of Cohen's $d = 0.44$ with a power of 0.80 using a two-sided alpha of 0.05. The sample size was calculated based on an estimated intra-class correlation for drinking refusal self-efficacy of 0.01 and 45 students per school. Taking into account an estimated attrition of 35%, 420 participants were planned to be recruited for the study. Further information can be found elsewhere [34].

The virtual reality simulation VR FestLab

The Danish VR application, VR FestLab, immerses users in a 360-degree filmed simulation and was specifically designed for adolescents aged between 15 and 18 years. A total of 128 distinct scenes allows users to engage with diverse simulation sequences. This interactive experience commences at the home of a school friend, where they both join in a birthday celebration. Within this simulation, users can navigate through two virtual rooms using eye movements. As they move within these spaces, they encounter various scenes, such as engaging in beer pong or participating in a flirting scenario, wherein they are presented with choices of both alcoholic and non-alcoholic beverages. Adhering to the taxonomy outlined in the Behavior Change Wheel [35], the simulation effectively integrates various behavior change functions, including education, training, modelling, and coercion/incentivization. Notably, individuals within the simulation can interact with role models who demonstrate the refusal to consume alcohol. If the user decides to consume alcohol and accepts an alcoholic drinking choice, the choice results in an incremental increase of a blood alcohol concentration (BAC) bar, calculated via an algorithm, which is filled at the top of the screen (see Figure 1, yellow BAC bar at the top of the screen).




Figure 1. Screenshot of one scene from the VR simulation FestLab

This algorithm considers factors like gender, alcohol content in grams, drinking pace, and the average weight of a 16-year-old boy or girl to calculate the BAC [36]. To calculate the exact BAC score, a BAC calculation by Becker and Nielsen [37] was used. The calculation for girls and boys is as follows:

$$\text{Female BAC} : \frac{\text{Alcohol consumed} \in \text{grams}}{(\text{body weight} \in \text{kg} \times 60\%) - (0,15 \times \text{hours} \text{ drinking start})}$$

$$\text{Male BAC} : \frac{\text{Alcohol consumed} \in \text{grams}}{(\text{body weight} \in \text{kg} \times 70\%) - (0,15 \times \text{hours} \text{ drinking start})}$$

Should the user consume an excessive amount of alcohol within a short span, as determined by the algorithm, they suffer a blackout. This blackout is first portrayed by the camera shaking and a magnifying display of the BAC bar. Following this, within the game the screen turns black, with the user subsequently awakening in a bedroom. In this bedroom, they receive messages that they have blacked out, concluding the simulation.

VR FestLab was pre-installed on Oculus Quest (Meta Platforms Inc., Menlo Park, USA) VR devices and handed out to the participants. Before engaging with the simulation, adolescents received instructions on device operation and navigation within the simulation. Following this, adolescents experienced VR FestLab for a maximum of 15 minutes in a classroom session. Depending on the simulation decisions, several rounds could be played during this time. After that, a structured 45-minute group reflection period was moderated by a trained study assistant in the classroom.

Measures

Data were gathered through electronic questionnaires administered during school hours within

classroom settings. The questionnaire was developed using the English versions of the respective scales (sensation seeking, drinking refusal skills, refusal communication skills) because no Danish versions were available (Multimedia Appendix 1). For the other measures (alcohol consumption, other drug consumption, knowledge/awareness of BAC) without existing scales in the literature, the project team created their own questions and items. The English questionnaires were translated to Danish and pre-tested in 31 students to determine their psychometric characteristics.

Adolescents were tasked with completing the questionnaire before (T0) and after the intervention (T1). Independent variables, including age, gender, alcohol consumption, other drug consumption, sensation seeking, drinking refusal skills, knowledge/awareness of BAC, and refusal communication skills, were assessed at T0 before the VR FestLab intervention. The dependent variable, virtual risk behavior, was surveyed at T1 following the intervention.

Virtual risk decisions

The primary outcome was virtual risk decisions, evaluated through a self-developed question: *“Did you pass out at any time during the party?”*. Answer options were dichotomized (yes/no). Passing out can only be achieved in the simulation when the number and types of drinks accepted in a given time period has resulted in a BAC of 2.0 permille, thus representing several risk decisions. The scenes and participant selections within the simulation were intentionally not recorded or tracked to afford participants the utmost freedom in their interactions. As a result, the blackout experience in the game could only be assessed verbally. If participants responded affirmatively, it was inferred that the simulation terminated prematurely due to excessive alcohol consumption in a condensed timeframe during the simulation.

Age and gender

Participant gender was dichotomously determined by the question *“Are you a girl or a boy? (State what you most identify as right now)”* while age was quantified using the question *“How old are you?”*.

Perceived family affluence, school performance and health

To gather information on the socio-demographic factors of the participants, the following questions were adapted from the Health Behavior in School-aged Children study [38]: *“How well-off do you think your family is?”*, *“What does your class teacher(s) think about your school performance compared to your classmates?”* and *“Would you say your health is...?”*

Alcohol consumption

Inquiries designed by the researchers were employed to estimate alcohol consumption with three single questions. The questions *“Have you ever drunk alcohol?”*, *“Have you ever been drunk?”* and *“Have you ever had 5 or more drinks on a single occasion?”* could be answered dichotomously (yes/no) by the participants.

Other drug consumption

To assess the consumption of substances other than alcohol, customized five questions were utilized. Participants were queried about their usage of cigarettes, hookah, e-cigarettes, snuff, or cannabis. Response options were dichotomized (yes/no).

Sensation seeking

Sensation seeking was measured using the eight-item Brief Sensation Seeking (BSSS-8) scale with Cronbach’s alpha 0.76 by Hoyle et al. [39]. Participants encountered eight statements, responding on a five-point Likert scale, ranging from “strongly disagree” to “strongly agree”. The 8-item scale

contained four subscales with 2 items each. Subscales, including “thrill and adventure seeking”, “experience seeking”, “disinhibition”, and “boredom susceptibility”, were calculated following the criteria outlined by Hoyle et al. [39]. Cronbach’s alpha of the BSSS-8 was 0.76 in our sample.

Drinking refusal skills – social pressure subscale

To evaluate drinking refusal skills within the context of peer pressure, the five-item social pressure subscale of the Drinking Refusal Self-Efficacy Questionnaire (DRSEQ) with Cronbach’s alpha 0.87 by Young et al. [40] was employed. Participants responded on a six-point Likert scale, ranging from “*I am very sure I could not resist drinking*” to “*I am very sure I could resist drinking*”. Cronbach’s alpha of the DRSEQ social pressure subscale was 0.88 in our sample.

Knowledge/awareness of blood alcohol concentration

Knowledge/awareness pertaining to BAC was measured using two self-constructed items. Participants were prompted to rate the statements “*It is easy for me to estimate my own alcohol tolerance*” and “*I know how much alcohol I can drink before I get drunk*” on a five-point Likert scale, spanning from “strongly disagree” to “strongly agree”.

Refusal communication skills

Refusal communication skills were assessed using two items drawn from the Alcohol Misuse Prevention Knowledge Questionnaire by Shope et al. [19]. Only two items from the Shope et al. [19] questionnaire were used because the other questionnaire items were not covering relevant aspects of VR FestLab [34]. Participants were asked to evaluate the statements “*If someone offers me a drink of alcohol and I say no, I can make them take no for an answer*” and “*If my best friends want me to drink beer with them and I don’t want to, I have ways to say no*” on a five-point Likert scale, ranging from “strongly disagree” to “strongly agree”.

Statistical analysis

Data analysis was performed using the R Studio software package (version 2022.07.2). The “two-step” methodology, as outlined by Herting & Costner [41], was applied to calculate model fit as a first step and then formulate the structural equation model as a second step. Initially, a confirmatory factor analysis (CFA) utilizing the lavaan package [42] was carried out to determine adequacy of the measurement models of the questionnaires. For this purpose, different single-factor models were created and calculated using a diagonally weighted least squares estimator that is suitable for categorical items. Model fit was checked by using a chi-square test in combination with other fit indices such as comparative fit index (CFI) ($CFI \geq .95$), root mean square error of approximation (RMSEA) ($RMSEA \leq .06$), Tucker-Lewis index (TLI) ($TLI \geq .95$) and standardized root mean square residual (SRMR) ($SRMR \leq .08$) [43].

Subsequently, a structural model with latent variables was calculated to investigate the influence of the constructs (alcohol consumption, other drug consumption, sensation seeking, drinking refusal skills and knowledge/awareness of BAC and refusal communication skills) on the main outcome virtual risk decisions and the correlation of the constructs with age and gender as covariates. In addition, a moderation analysis for gender and age was run. Estimated scores for each latent variable were calculated and subsequently incorporated into a logistic regression model to predict virtual risk decisions.

Results

A total of 268 students from seven schools were assigned to the intervention group. Of these, 183 students participated in the baseline survey (T0). The main reasons for dropout between allocation and baseline survey were that one complete school ($n=36$) dropped out, one complete school class

(n=15) dropped out, and 34 students were not willing to participate or did not provide consent. After the intervention, two additional participants dropped resulting in 181 individuals completing the first follow-up questionnaire (T1). The subsequent analysis was based on a complete dataset from 161 participants. Additional information on allocation and dropouts can be found elsewhere [34]. To test whether there were differences between respondents at T1 and those who provided complete information at T1 and were included in the analysis, independent t-tests for metric and chi-squared tests were conducted for nominal and ordinal scaled sociodemographic variables. The analyses showed no differences between the two groups in terms of age ($t(179) = 0.61, P = .54$), gender ($\chi^2(1) = 0.31, P = .58$), and Family Affluence Scale (FAS) ($\chi^2(1) = 0.25, P = .42$). For the chi-square analysis of the FAS between completer and non-completer, the prerequisite of cell frequencies above 5 was violated, which is why Fisher's exact test was applied.

Sample characteristics

Table 1 provides an overview of demographic attributes. Gender distribution was equal, constituting 78 out of 161 (48.4%) being female. The mean age of the sample was 15.6 (SD= 0.72). Most respondents 88.8% (n=143) reported low to moderate perceived family affluence, and approximately 62.1% (n=100) rated their perceived school performance as good to very good. Additionally, 79.5% (n=128) reported good to very good health. The majority of respondents 73.9% (n=119) reported having engaged in binge drinking at some point in their lives.

Table 1. Characteristics of the study population (n=161).

| Characteristics | Value |
|---|-------------|
| Age in Years , mean (SD) | 15.6 (0.72) |
| Gender, female, n (%) | 78 (48.4) |
| Perceived Family Affluence, n (%) | |
| Low to medium | 143 (88.8) |
| High to very high | 18 (11.2) |
| Perceived School Performance, n (%) | |
| Good to very good | 100 (62.1) |
| Below average to average | 61 (37.9) |
| Perceived Health and Well-being, n (%) | |
| Good to excellent | 128 (79.5) |
| Poor to fair | 33 (20.5) |
| Lifetime Binge Drinking, n (%) | |
| Yes | 119 (73.9) |
| No | 42 (26.1) |

Calculations of model fit

The single-factor models generally showed a good fit in the CFA and could therefore be transferred to the structural equation model without adjustments. Sensation seeking provided a mediocre fit with CFI and RMSEA beyond their cut-offs, while the 90%-CI of the RMSEA included the cut-off (see Table 2). The factors “knowledge/awareness of BAC” and “refusal communication skills” were combined for the CFA for statistical reasons, as otherwise it would not have been possible to determine the fit, as they only contained two items each. Finally, the combination of the two factors showed high loadings on the individual and superordinate factors, which resulted in good fit values in the CFA. The results of the single-factor models are shown in Table 2 and Figure 2.

Table 2. The goodness of fit indices of the confirmatory factor analysis for single-factor models.

| Model | Items | χ^2 -test | P- | CFI | TLI | SRMR | RMSEA | P-value |
|-------|-------|----------------|----|-----|-----|------|-------|---------|
|-------|-------|----------------|----|-----|-----|------|-------|---------|

| | | | value | | | | (90% CI) | | |
|--|--|---|-------|-----|------|------|----------|---------------------|-----|
| Alcohol consumption | | 3 | 2.24 | .33 | 1.00 | 1.00 | 0.04 | 0.03 (0.00-0.16) | .46 |
| Other drug consumption | | 5 | 3.02 | .70 | 1.00 | 1.00 | 0.02 | 0.00 (0.00-0.08) | .84 |
| Sensation seeking | | 4 | 7.38 | .03 | 0.92 | 0.76 | 0.05 | 0.13 (0.04-0.23) | .07 |
| Drinking Refusal Skills – social pressure subscale | | 5 | 19.47 | .01 | 0.97 | 0.95 | 0.03 | 0.13 (0.08-0.20) | .01 |
| Knowledge of BAC & Refusal communication skills | | 4 | 1.42 | .23 | 1.00 | 0.99 | 0.02 | 0.05 (0.00-0.22) | .32 |

χ^2 = Chi-squared, CFI = comparative fit index, TLI = Tucker-Lewis index, SRMR = standardized root mean square residual, RMSEA = root mean square error of approximation, CI = confidence interval, BAC = blood alcohol concentration

Determinants linked to risky drinking decisions

A general overview of the bivariate correlations of all variables used in this study can be found in the Multimedia Appendix 2. Age and gender were included as covariates without moderation in the structural equation model, as shown in Figure 2. The path model used to test the formulated hypotheses showed an unsatisfying model fit, χ^2 -test (238, 161) = 422.59, $P < .001$, CFI = 0.94, SRMR = 0.12, RMSEA = 0.07, CIRMSEA = 0.06 – 0.08. Nevertheless, it was decided to retain the model, because all included measurement models were a-priori tested and showed good fit. Due to the explorative nature of the study, the structural model included all possible paths. Therefore, the reason for misfit can be attributed to the cross-loadings of various scales only. As the structural part of the model was the aim of the study, it was decided to retain it. Figure 2 shows the tested model. Prior alcohol experiences and knowledge/awareness on BAC were significantly associated with virtual risk decisions in VR FestLab. Participants who reported prior alcohol consumption had a 4.31-fold higher chance of showing virtual risk decisions in the simulation (OR = 4.31 (CI = 1.70-10.84), $P = .01$). Therefore, hypothesis (3) can be accepted. Apart from that, adolescents with high knowledge/awareness of BAC at baseline were 0.32 less likely to make virtual risk decisions in the simulation (OR = 0.32 (CI = 0.11-0.93), $P = .04$). Higher social pressure drinking refusal skills (OR = 0.89 (CI = 0.69-1.16), $P = .39$), other consumption of drugs (OR = 0.72 (CI = 0.36-1.45), $P = .36$), sensation seeking (OR = 0.69 (CI = 0.36-1.34), $P = .27$), or refusal communication skills (OR = 1.45 (CI = 0.68-3.10), $P = .34$) showed no significant relationship to virtual risk decisions in the simulation. Therefore, hypotheses (1) must be rejected and hypothesis (2) can only be partially confirmed.




Figure 2. Path model of the relation of drinking-refusal skills – social pressure, alcohol consumption, drug consumption, sensation seeking, knowledge/awareness of blood alcohol concentration and refusal communication skills with virtual risk decisions.

DR-SP = Drinking refusal –social pressure, Alcohol Consum = alcohol consumption, Other Consum = other consumption of drugs, SeSe = Sensation seeking, Know BAC = Knowledge/awareness of blood alcohol concentration, Refu Comm Skills = Refusal communication skills, P = significance value, OR = Odds ratio

Moderation effects of gender or age regarding virtual risk decisions

Subsequently, it was analyzed whether age and gender had a moderating influence on the latent variables. The results of this moderation analysis are presented in Multimedia Appendix 3. All individual moderation analyses for the variables (alcohol consumption, other drug consumption, sensation seeking, drinking refusal skills and knowledge/awareness of BAC & refusal communication skills) showed no significant moderation. Accordingly, age or gender did not significantly alter the effects of the tested constructs on virtual risk decisions in the simulation. Age and gender also did not significantly influence the dependent variable virtual risk decisions (Multimedia Appendix 3).

Discussion

Principal Results

In summary, the structural equation model exhibited a significant association between prior alcohol experiences and knowledge/awareness of BAC with virtual risk decisions, supporting hypothesis 3. The other initially hypothesized factors (hypotheses 1 and 2), including drinking refusal skills, sensation seeking, refusal communication skills, and consumption of other drugs, did not demonstrate significant associations with virtual risk decisions. Upon exploring moderating factors such as age and gender, no moderation on virtual risk decisions was identified. The present study has underscored that prior alcohol experiences are notably linked to virtual risk decisions. Participants who have encountered alcohol and engaged in binge drinking appear to perceive the VR FestLab game as realistically simulating their personal behaviors, leading them to enact these behaviors within the game. Apart from that, it appears that higher knowledge/awareness of BAC is a protective factor and leads to fewer virtual risk decisions in the simulation.

Comparison with Prior Work

These findings align with qualitative insights collected from focus group investigations involving adolescents discussing the simulation [44, 45]. Adolescents reported finding the VR simulation remarkably realistic, evoking sensations akin to being present at an actual party [44, 45]. The initial participation in VR FestLab possibly aimed to ascertain whether the simulated party aligned with their expectations and whether outcomes matched real-world drinking behaviors. The study's dataset does not offer insights into potential variations in behavior between several attempts with VR FestLab. However, it seems plausible that distinct behaviors and strategies would be attempted in subsequent trials, as highlighted by focus group participants in qualitative interviews. The participants expressed the view that VR serves as a medium for experimenting with various behaviors and that repeated engagements with the VR simulation allow for exploring different strategies, such as drinking versus abstaining, while observing the reactions of simulation characters [45]. This points to VR's potential benefit in alcohol prevention, allowing participants to experiment with diverse approaches and behaviors in risky scenarios. The present study found a relatively high lifetime prevalence of binge drinking in the group of adolescents aged 14-18 years, at about 73.9% (n=119). In this context, it would be interesting for future studies to investigate the association between previous alcohol consumption and the likelihood of risk decisions in different social environments. These results may also provide hints for further virtual scenarios that can be integrated into the VR FestLab. Hadley et al. [46], who combined VR environments with emotion regulation and a risk reduction intervention, also arrived at similar conclusions, indicating that VR, through better simulated contextual cues of risky situations, facilitates the application of different emotion regulation strategies among adolescents [46]. In the real world, such experimentation of behavioral strategies is constrained by the necessity of personally experiencing risky behavior and its consequences, often without the possibility of multiple trials (e.g. in the case of blackout).

The hypothesis that individuals with high sensation seeking tendencies would exhibit elevated virtual risk decisions within the game could not be substantiated by the study's findings. The lack of significant association between sensation seeking and virtual risk decisions could potentially be attributed to the design of VR FestLab. The game's preventive nature might not have resonated sufficiently with individuals displaying high sensation-seeking behavior. Given that sensation seeking characterizes those seeking diverse, potentially risky experiences [47], the simulated risk behavior in VR FestLab might not have provided compelling cues for such participants. Participants were aware of the virtual and simulated nature of the risk decisions, possibly leading to the observed absence of virtual risk decisions in the game. This absence could also stem from the absence of personal consequences, such as intoxication, emotional arousal, and sensory perceptions, conveyed through VR FestLab. This lack of stimuli might not engage adolescents with a high sensation-seeking drive. On the contrary, sensation seeking is a multifaceted construct, not exclusively tied to risk behavior. The study of Ravert et al. [48] found that sensation seeking, manifested as a search for stimulation, could also be positively linked to psychological well-being. Furthermore, sensation seeking might have been manifest in non-alcohol-related scenes within VR FestLab (e.g., interactions with game characters), rendering the presumed virtual risk decisions unappealing to participants with pronounced sensation-seeking tendencies.

Likewise, drinking refusal self-efficacy skills failed to display an association with reduced virtual risk decisions in this study. Those with higher baseline drinking refusal self-efficacy skills might have opted for less risky drinking decisions within VR FestLab, possibly bypassing activities like beer pong. These observations align with the study of Guldager et al. [34], assessing the efficacy of VR FestLab. While effects were insignificant, a more pronounced increase in drinking refusal skills was noted among participants whose baseline skills lay below the median compared to those above the median. Despite the documented real-world associations between higher drinking refusal skills, drug consumption, and refusal communication skills with risky behavior [49, 50], these correlations did not manifest in the virtual world.

Although it was originally hypothesized that higher refusal communication skills would be associated with a reduction in virtual risk decisions, this hypothesis was not supported by the present study. It is possible that participants were more exploratory when playing VR FestLab to "just see what happens next" and did not use their refusal communication skills in the simulation to say no to drinking offers. Qualitative interviews with adolescents who played VR FestLab revealed that they felt the peer pressure in the simulation was weaker than in real life [44], which could be a reason why refusal communication skills were not associated with reduced virtual risk decisions in the present study.

On the other hand, the present study has shown that individuals with higher knowledge/awareness of BAC are less likely to make risky decisions in virtual environments. It is reasonable to assume that those who understand how different drinks affect BAC are less likely to engage in risky decisions in VR FestLab compared to those who cannot accurately assess the effects of BAC. These assumptions are supported by literature, which indicates that individuals with greater knowledge of low-risk alcohol consumption are more likely to reduce their alcohol intake and make fewer risky decisions [51]. Increased awareness of BAC could enhance risk perception and enable better anticipation and assessment of the consequences of risky decisions. It is possible that individuals without knowledge/awareness of BAC were more willing to experiment in the simulation, leading to an increase in virtual risk-taking decisions. VR FestLab is a prevention program that uses an in-game BAC bar to illustrate the effects of alcohol-related decisions on BAC. The in-game presentation may help adolescents to avoid risky decisions by increasing their knowledge of BAC. Other research [52] using VR in e-cigarette prevention have also shown that virtual prevention simulations can increase substance-related knowledge and harmful perceptions about e-cigarettes.

To our knowledge, this study is the first attempt to examine the relationship between real-life behavior and attitudes and virtual risk decisions in the context of alcohol prevention making it

difficult to contextualize the findings with other virtual alcohol prevention studies. The findings illuminate VR's capacity to replicate real-life behavior within simulated environments. Other studies [53-55] using VR in other thematic areas have also concluded that there are links between simulated behavior and real behavior. The combination of prevention and VR within research is relatively novel, necessitating further exploration into leveraging simulated settings for risk behavior prevention and health promotion. Existing studies underscore an increasing adoption of VR for prevention and health promotion [56]. Building behavior change interventions and predicting behavioral transformations is challenging in real-world scenarios [57]. Immersive technologies, such as VR, can harness specific attributes like training and realism to facilitate future behavioral change. Nevertheless, the translation of real-life behavior into virtual simulations, along with the potential impact of virtually acquired behavioral patterns on subsequent real-world behavior, remains uncertain. Research suggests that altering an individual's avatar representation in VR influences their behavior and psychological disposition in the real world, an occurrence termed the Proteus effect [24, 58]. This phenomenon could also be probed in the context of alcohol prevention, where avatars could shape participants' self-image and preventive behaviors. Addressing these unresolved questions in future studies while delving into the mechanisms at play in virtual simulations could position VR as a valuable asset in prevention and health promotion.

Limitations

In the present study, certain limitations must be acknowledged when interpreting the results. Firstly, the applicability of the findings warrants consideration. Participants were drawn from diverse Danish schools, making the results pertinent exclusively to Danish students. Additionally, the recruitment was limited to a specific region within Denmark, implying that adolescents from other regions within Denmark or other countries might exhibit disparate outcomes.

Secondly, the results should be cautiously approached due to the reliance on self-reported data, which is susceptible to common biases such as respondent bias, extreme responding, and social desirability bias [59]. These potential biases could have influenced and skewed the reported results. The timing of the follow-up data collection (T1), conducted immediately after the intervention, may have influenced the results. Furthermore, this study did not survey risk behaviors that may occur prior to a blackout, such as engaging in sexual activity or driving under the influence of alcohol. Future studies should consider measuring these risk behaviors before a blackout occurs in order to establish a relationship. Specifically, the question about certain risk decisions may have been biased (e.g. experiencing a blackout). The potential for a Hawthorne effect should be considered when interpreting the results, because the students in the intervention group were not blinded and tested the new VRFestLab application. Their knowledge of participation in the study and the aim of VR FestLab can influence their self-assessment of whether they experienced a blackout in the application, which could potentially distort the results. Since there was no content tracking during participants' engagement with the simulation, the navigation patterns within the simulation remain uncharted. Integrating a content log in future analyses could offer deeper insights into participants' usage behavior.

Thirdly, organizational limitations limited the playing time to 15 minutes for each participant. This restriction could have led to variations in the number of rounds played within the simulation, potentially affecting the reported outcomes. Additionally, the controlled setting of the VR FestLab intervention study might have curtailed the expression of extreme behaviors that could naturally occur among adolescents. This controlled environment may have influenced the observed results.

Conclusions

Despite these limitations, this study illuminates the link between real-life behavior and virtual simulations in alcohol prevention. The significance of prior alcohol experiences in shaping virtual

risk decisions underscores the potential of VR to mimic real behaviors. The study highlights VR's potential as a valuable tool for behavior change interventions and health promotion. To harness this potential, future research should delve into mechanisms underpinning behavioral expression in virtual environments. As VR evolves, it presents a promising avenue for health promotion strategies, offering a unique medium to experiment with behaviors in risk-prone scenarios. This research signifies the importance of advancing our understanding of the intersection between human behavior, virtual simulations, and their real-world implications.

Acknowledgements

We thank Kristoffer Drue Knudsen for his literature research and data preparation and Satayesh Lavasani Kjær, Nørgaard Lehnsted and Maria Busk Damgaard for data collection. We would also like to thank the students, teachers and principals who supported our research through their active participation.

Data Availability

The analysed data can be accessed from the Zenodo database (doi: 10.5281/zenodo.8276272).

Funding

This research was supported by TrygFonden, Denmark (grant number ID 129372). The funder did not have any involvement in the study design, conduct, or publication of results.

Author contributions

Conceptualization: RH, CS, JG

Data curation: RH, JG, CS,

Formal Analysis: RH, CS, DS

Funding acquisition: CS

Investigation: JG, PL, GM, CS

Methodology: RH, DS, CS

Project administration: CS

Resources: CS, PL, GM

Software: PL, GM

Supervision: CS

Validation: RH, JG, DS, CS

Visualization: RH, DS, CS

Writing – original draft: RH, JG, DS, CS

Writing – review & editing: RH, JG, DS, PL, GM, CS

Conflicts of Interest

None declared.

Abbreviations

BAC: Blood alcohol concentration

BSSS: Brief sensation seeking scale

CFA: Confirmatory factor analysis

CFI: Comparative fit index

DRSEQ: Drinking Refusal Self-Efficacy Questionnaire

TLI Tucker-Lewis index

VR: Virtual Reality

RMSEA: Root mean square error of approximation

SRMR: Standardized root mean square residual

Multimedia Appendix 1

Investigator-developed questionnaire.

Multimedia Appendix 2

Bivariate correlations of all variables.

Multimedia Appendix 3

Results of the moderation analyses.

References

1. Jernigan DH. Alcohol and Public Health: Failure and Opportunity. *Milbank Q.* 2023;101(S1):552-78. doi: 10.1111/1468-0009.12631.
2. Shield K, Manthey J, Rylett M, Probst C, Wettlaufer A, Parry CD, et al. National, regional, and global burdens of disease from 2000 to 2016 attributable to alcohol use: a comparative risk assessment study. *Lancet Public Health.* 2020;5(1):e51-e61. doi: 10.1016/S2468-2667(19)30231-2.
3. World Health Organization. Global status report on alcohol and health 2018. Geneva: World Health Organization; 2018.
4. ESPAD Group. ESPAD Report 2019: Results from the european school survey project on alcohol and other drugs. Luxembourg: EMCDDA Joint Publications, Publications Office of the European Union; 2020.
5. Foxcroft DR, Tsertsvadze A. Universal school-based prevention programs for alcohol misuse in young people. *Cochrane Database Syst Rev.* 2011 (9):CD009308. doi: 10.1002/14651858.CD009113.
6. Adan A, Forero DA, Navarro JF. Personality traits related to binge drinking: A systematic review. *Front Psychiatry.* 2017;8:134. doi: 10.3389/fpsy.2017.00134.
7. Stautz K, Cooper A. Impulsivity-related personality traits and adolescent alcohol use: a meta-analytic review. *Clinical psychology review.* 2013;33(4):574-92. doi: 10.1016/j.cpr.2013.03.003.
8. Percy A, McKay MT, Cole JC. Interplay Between Sensation Seeking and Parental Rules in the Emergence of Heavy Episodic Drinking. *J Res Adolesc.* 2019;29(4):814-21. doi: 10.1111/jora.12435.
9. Bonar EE, Souweidane MA, Blow FC, Bohnert AS, Bauermeister JA, Young SD, et al. High-intensity drinking among adolescent and emerging adult risky drinkers. *Subst Abuse.* 2022;43(1):713-21. doi: 10.1080/08897077.2021.2007513.
10. Creamer MR, Portillo GV, Clendennen SL, Perry CL. Is Adolescent Poly-tobacco Use Associated with Alcohol and Other Drug Use? *Am J Health Behav.* 2016;40(1):117-22. doi: 10.5993/ajhb.40.1.13.
11. Vijapur SM, Levy NS, Martins SS. Cannabis use outcomes by past-month binge drinking status in the general United States population. *Drug Alcohol Depend.* 2021;228:108997. doi: 10.1016/j.drugalcdep.2021.108997.
12. Gunn RL, Sokolovsky AW, Drohan MM, Boyle HK, Stevens AK, White HR, et al. The role of alcohol and cannabis co-use in drinking rate and its impact on consequences. *Alcohol Clin Exp Res.* 2022;46(11):2110-20. doi: 10.1111/acer.14946.
13. McGuire WJ. Resistance to persuasion conferred by active and passive prior refutation of

the same and alternative counterarguments. *J Abnorm Soc Psycho*. 1961;63(2):326-32. doi: 10.1037/h0048344.

14. Bandura A. *Social learning theory*. Englewood Cliffs, N.J: Prentice-Hall; 1977. ISBN: 0-13-816751-6.

15. Botvin GJ, Griffin KW. *Life Skills Training: A competence enhancement approach to tobacco, alcohol, and drug abuse prevention*. In: Scheier L, editor. *Handbook of adolescent drug use prevention: Research, intervention strategies, and practice*: American Psychological Association; 2015.

16. Shin Y. Narrative Engagement and Interpersonal Communication about Substance Use on Adolescent Substance Use Behaviors: A Case Study of Keepin' it REAL. *West J Commun*. 2021;85(3):427-50. doi: 10.1080/10570314.2020.1846074.

17. Wynn SR, Schulenberg J, Kloska DD, Laetz VB. The Mediating Influence of Refusal Skills in Preventing Adolescent Alcohol Misuse. *J Sch Health*. 1997 1997/11/01;67(9):390-5. doi: 10.1111/j.1746-1561.1997.tb07183.x.

18. Donohue B, Van Hasselt VB, Hersen M, Perrin S. Substance refusal skills in a population of adolescents diagnosed with conduct disorder and substance abuse. *Addict Behav*. 1999;24(1):37-46. doi: 10.1016/S0306-4603(98)00035-5.

19. Shope JT, Copeland LA, Maharg R, Dielman TE, Butchart AT. Assessment of adolescent refusal skills in an Alcohol Misuse Prevention Study. *Health Educ Q*. 1993;20(3):373-90. doi: 10.1177/109019819302000308.

20. Teesson M, Newton NC, Slade T, Chapman C, Birrell L, Mewton L, et al. Combined prevention for substance use, depression, and anxiety in adolescence: a cluster-randomised controlled trial of a digital online intervention. *The Lancet Digital Health*. 2020;2(2):e74-e84. doi: 10.1016/S2589-7500(19)30213-4.

21. Hasking P, Schofield L. Examining alcohol consumption with the theory of planned behaviour: Do health and alcohol knowledge play a role? *Psychol Health Med*. 2015;20(7):838-45. doi: 10.1080/13548506.2014.969748.

22. Padget A, Bell ML, Shamblen SR, Ringwalt CL. Does Learning About the Effects of Alcohol on the Developing Brain Affect Children's Alcohol Use? *Prev Sci*. 2006;7(3):293-302. doi: 10.1007/s11121-006-0030-9.

23. Morina N, Ijntema H, Meyerbröcker K, Emmelkamp PMG. Can virtual reality exposure therapy gains be generalized to real-life? A meta-analysis of studies applying behavioral assessments. *Behav Res Ther*. 2015;74:18-24. doi: 10.1016/j.brat.2015.08.010.

24. Ratan R, Beyea D, Li BJ, Graciano L. Avatar characteristics induce users' behavioral conformity with small-to-medium effect sizes: a meta-analysis of the proteus effect. *Media Psychol*. 2020;23(5):651-75. doi: 10.1080/15213269.2019.1623698.

25. Bordnick PS, Carter BL, Traylor AC. What virtual reality research in addictions can tell us about the future of obesity assessment and treatment. *J Diabetes Sci Technol*. 2011;5(2):265-71. doi: 10.1177/193229681100500210.

26. Slater M, Banakou D, Beacco A, Gallego J, Macia-Varela F, Oliva R. A separate reality: An update on place illusion and plausibility in virtual reality. *Front Virtual Real*. 2022;3:914392. doi: doi.org/10.3389/frvir.2022.914392.

27. Alcañiz M, Rey B, Tembl J, Parkhutik V. A Neuroscience Approach to Virtual Reality Experience Using Transcranial Doppler Monitoring. *Presence (Camb)*. 2009;18(2):97-111. doi: 10.1162/pres.18.2.97.

28. Pan X, Hamilton A. Why and how to use virtual reality to study human social interaction: The challenges of exploring a new research landscape. *Br J Psychol*. 2018;109(3):395-417. doi: 10.1111/bjop.12290.

29. Blascovich J, Loomis J, Beall AC, Swinth KR, Hoyt CL, Bailenson JN. Immersive virtual


- environment technology as a methodological tool for social psychology. *Psychol Inq.* 2002;13(2):103-24. doi: 10.1207/S15327965PLI1302_01.
30. Zeuwts LH, Vanhuele R, Vansteenkiste P, Deconinck FJ, Lenoir M. Using an immersive virtual reality bicycle simulator to evaluate hazard detection and anticipation of overt and covert traffic situations in young bicyclists. *Virtual Real.* 2023;27:1507–27 doi: 10.1007/s10055-023-00746-7.
31. Durl J, Dietrich T, Pang B, Potter L-E, Carter L. Utilising virtual reality in alcohol studies: A systematic review. *Health Educ J.* 2018;77(2):212-25. doi: 10.1177/0017896917743534.
32. Prediger C, Helmer SM, Hrynyschyn R, Stock C. Virtual Reality-Based Alcohol Prevention in Adolescents: A Systematic Review. *Adolescents.* 2021;1(2):138-50. PMID: doi:10.3390/adolescents1020011.
33. Vallentin-Holbech L, Guldager JD, Dietrich T, Rundle-Thiele S, Majgaard G, Lyk P, et al. Co-Creating a Virtual Alcohol Prevention Simulation with Young People. *Int J Environ Res Public Health.* 2020;17(3):1097. doi: 10.3390/ijerph17031097.
34. Guldager JD, Kjær SL, Grittner U, Stock C. Efficacy of the Virtual Reality Intervention VR FestLab on Alcohol Refusal Self-Efficacy: A Cluster-Randomized Controlled Trial. *Int J Environ Res Public Health.* 2022;19(6):3293. doi: 10.3390/ijerph19063293.
35. Michie S, Van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci.* 2011;6(1):1-12. doi: 10.1186/1748-5908-6-42.
36. Lyk P, Majgaard G, Vallentin-Holbech L, Dalgaard Guldager J, Dietrich T, Rundle-Thiele S, et al. Co-Designing and Learning in Virtual Reality. Development of Tool for Alcohol Resistance Training. *Electron J e-Learn.* 2020;18(3):213-28. doi: 10.34190/EJEL.20.18.3.002.
37. Becker U, Nielsen AS. Alkohol, fakta om virkning. 2023 [cited 2024 January 7th]; Available from: <https://www.sundhed.dk/borger/patienthaandbogen/psyke/sygdomme/alkohol/alkohol-fakta-om-virkning/>.
38. Currie C, Inchley J, Molcho M, Lenzi M, Veselska Z, Wild F. Health behaviour in school-aged children (HBSC) study protocol: background, methodology and mandatory items for the 2013/14 survey. St Andrews: Child and Adolescent Health Research Unit (CAHRU); 2014.
39. Hoyle RH, Stephenson MT, Palmgreen P, Lorch EP, Donohew RL. Reliability and validity of a brief measure of sensation seeking. *Pers Individ Differ.* 2002;32(3):401-14. doi: 10.1016/S0191-8869(01)00032-0.
40. Young RM, Hasking PA, Oei TP, Loveday W. Validation of the Drinking Refusal Self-Efficacy Questionnaire--Revised in an Adolescent Sample (DRSEQ-RA). *Addict Behav.* 2007;32(4):862-8. doi: 10.1016/j.addbeh.2006.07.001.
41. Herting JR, Costner HL. Another perspective on "the proper number of factors" and the appropriate number of steps. *Struct Equ Modeling.* 2000;7(1):92-110. doi: 10.1207/S15328007SEM0701_05.
42. Rosseel Y. lavaan: An R package for structural equation modeling. *J Stat Softw.* 2012;48(2):1-36. doi: 10.18637/jss.v048.i02.
43. Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Modeling.* 1999;6(1):1-55. doi: 10.1080/10705519909540118.
44. Guldager JD, Kjær SL, Lyk P, Dietrich T, Rundle-Thiele S, Majgaard G, et al. User Experiences with a Virtual Alcohol Prevention Simulation for Danish Adolescents. *Int J Environ Res Public Health.* 2020;17(19):6945. doi: 10.3390/ijerph17196945.
45. Hrynyschyn R, Prediger C, Lyk P, Majgaard G, Helmer SM, Stock C. Adolescents' perceptions and user experiences with a virtual reality-based alcohol prevention tool in

- Germany: A focus group study. *Front Public Health*. 2023;11:1054015. doi: 10.3389/fpubh.2023.1054015.
46. Hadley W, Houck C, Brown LK, Spitalnick JS, Ferrer M, Barker D. Moving Beyond Role-Play: Evaluating the Use of Virtual Reality to Teach Emotion Regulation for the Prevention of Adolescent Risk Behavior Within a Randomized Pilot Trial. *J Pediatr Psychol*. 2018;44(4):425-35. doi: 10.1093/jpepsy/jsy092.
47. Zuckerman M, Kolin EA, Price L, Zoob I. Development of a sensation-seeking scale. *J Consult Psychol*. 1964;28(6):477-82. doi: 10.1037/h0040995.
48. Ravert RD, Donnellan MB. Impulsivity and Sensation Seeking: Differing Associations with Psychological Well-Being. *Appl Res Qual Life*. 2021;16(4):1503-15. doi: 10.1007/s11482-020-09829-y.
49. Oei TPS, Morawska A. A cognitive model of binge drinking: The influence of alcohol expectancies and drinking refusal self-efficacy. *Addict Behav*. 2004;29(1):159-79. doi: 10.1016/S0306-4603(03)00076-5.
50. Botvin GJ, Griffin KW, Paul E, Macaulay AP. Preventing tobacco and alcohol use among elementary school students through life skills training. *J Child Adolesc Subst Abuse*. 2003;12(4):1-17. doi: 10.1300/J029v12n04_01.
51. Islam MM, Hoffmann MW, Rahman MB. Knowledge of low-risk drinking and its relationship with a reduction in alcohol consumption: Results from six waves of an Australian national survey. *Addict Behav*. 2019;95:172-7. doi: 10.1016/j.addbeh.2019.03.016.
52. Weser VU, Duncan LR, Sands BE, Schartmann A, Jacobo S, François B, et al. Evaluation of a virtual reality E-cigarette prevention game for adolescents. *Addict Behav*. 2021;122:107027. doi: 10.1016/j.addbeh.2021.107027.
53. Wang L, Christensen JL, Jeong DC, Miller LC. Virtual prognostication: When virtual alcohol choices predict change in alcohol consumption over 6-months. *Comput Hum Behav*. 2019;90:388-96. doi: 10.1016/j.chb.2018.08.025.
54. Real FJ, Whitehead M, Ollberding NJ, Rosen BL, Meisman A, Crosby LE, et al. A virtual reality curriculum to enhance residents' behavioral health anticipatory guidance skills: a pilot trial. *Acad Pediatr*. 2023;23(1):185-92. doi: 10.1016/j.acap.2022.07.011.
55. Thomas JG, Goldstein CM, Bond DS, Hadley W, Tuerk PW. Web-based virtual reality to enhance behavioural skills training and weight loss in a commercial online weight management programme: The Experience Success randomized trial. *Obesity Science & Practice*. 2020;6(6):587-95. doi: 10.1002/osp4.451.
56. Buttazzoni A, Parker A, Minaker L. Investigating the mental health implications of urban environments with neuroscientific methods and mobile technologies: A systematic literature review. *Health Place*. 2021;70:102597. doi: 10.1016/j.healthplace.2021.102597.
57. Davis R, Campbell R, Hildon Z, Hobbs L, Michie S. Theories of behaviour and behaviour change across the social and behavioural sciences: a scoping review. *Health Psychol Rev*. 2015;9(3):323-44. doi: 10.1080/17437199.2014.941722.
58. Yee N, Bailenson J. The Proteus effect: The effect of transformed self-representation on behavior. *Hum Commun Res*. 2007;33(3):271-90. doi: 10.1111/j.1468-2958.2007.00299.x.
59. Rosenman R, Tennekoon V, Hill LG. Measuring bias in self-reported data. *Int J Behav Healthc Res*. 2011;2(4):320-32. doi: 10.1504/ijbhr.2011.043414.


Supplementary Files

Figures

Screenshot of one scene from the VR simulation FestLab.



Path model of the relation of drinking-refusal skills – social pressure, alcohol consumption, drug consumption, sensation seeking, knowledge/awareness of blood alcohol concentration and refusal communication skills with virtual risk decisions.



Multimedia Appendixes

Investigator-developed questionnaire.

URL: <http://asset.jmir.pub/assets/e960274c1ba655f11dd12d20812f93c7.pdf>

Bivariate correlations of all variables.

URL: <http://asset.jmir.pub/assets/549f1d0598ade5733a474bfbbf26c16c.pdf>

Results of the moderation analyses.

URL: <http://asset.jmir.pub/assets/9e75861d601c0158558b0e0a985b9db3.pdf>



TOC/Feature image for homepages

Image for the homepage.

