

# **Predictors of Healthcare Practitioners' Intention to Use AI-Enabled Clinical Decision Support Systems (AI-CDSSs): A Meta-Analysis Based on the Unified Theory of Acceptance and Use of Technology (UTAUT)**

Julius Dingel, Anne-Kathrin Kleine, Julia Cecil, Anna Sigl, Eva Lermer, Susanne Gaube

Submitted to: Journal of Medical Internet Research  
on: February 15, 2024

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

## ***Table of Contents***

---

<b>Original Manuscript.....</b>	<b>5</b>
---------------------------------	----------

Preprint  
JMIR Publications

# Predictors of Healthcare Practitioners' Intention to Use AI-Enabled Clinical Decision Support Systems (AI-CDSSs): A Meta-Analysis Based on the Unified Theory of Acceptance and Use of Technology (UTAUT)

Julius Dingel<sup>1\*</sup>; Anne-Kathrin Kleine<sup>1\*</sup>; Julia Cecil<sup>1</sup>; Anna Sigl<sup>2</sup>; Eva Lermer<sup>1, 2</sup>; Susanne Gaube<sup>3</sup>

<sup>1</sup>LMU Munich Munich DE

<sup>2</sup>Technical University of Applied Sciences Augsburg Augsburg DE

<sup>3</sup>University College London London GB

\*these authors contributed equally

## Corresponding Author:

Anne-Kathrin Kleine

LMU Munich

Geschwister-Scholl-Platz 1

Munich

DE

## Abstract

**Background:** Artificial Intelligence-enabled Clinical Decision Support Systems (AI-CDSSs) offer potential for improving healthcare outcomes, but their adoption among healthcare practitioners remains limited.

**Objective:** The meta-analysis identifies predictors influencing healthcare practitioners' intention to use AI-CDSSs based on the Unified Theory of Acceptance and Use of Technology (UTAUT) and additional literature.

**Methods:** The literature search using electronic databases, forward searches, conference programs, and personal correspondence yielded 7,731 results, of which 17 studies met the inclusion criteria. Random-effects meta-analysis, relative weights analyses, and meta-analytic moderation and mediation analyses were used to examine the relationships of relevant predictor variables with the intention to use AI-CDSSs.

**Results:** The meta-analysis results supported the application of the UTAUT to the context of the intention to use AI-CDSSs. The results show that performance expectancy ( $rc = .66$ ), effort expectancy ( $rc = .55$ ), social influence ( $rc = .66$ ), and facilitating conditions ( $rc = .66$ ) were positively associated with the intention to use AI-CDSSs, in line with the predictions of the UTAUT. The meta-analysis further identified positive attitude ( $rc = .63$ ), trust ( $rc = .73$ ), anxiety ( $rc = -.41$ ), perceived risk ( $rc = -.21$ ), and innovativeness ( $rc = .54$ ) as relevant additional predictors. Trust emerged as the most influential predictor overall. The results of moderation analyses show that the relationship between social influence and use intention becomes weaker with increasing age. In addition, the relationship between effort expectancy and use intention was stronger for diagnostic AI-CDSSs compared to devices that combined diagnostic and treatment recommendations. Finally, the relationships between facilitating conditions and use intention was mediated through performance and effort expectancy.

**Conclusions:** The meta-analysis contributes to the understanding of the predictors of the intention to use AI-CDSSs based on an extended UTAUT model. More research is needed to substantiate the identified relationships and to explain the observed variations in effect sizes by identifying relevant moderating factors. The research findings bear important implications for the design and implementation of training programs for healthcare practitioners to ease the adoption of AI-CDSSs into their practice. Clinical Trial: <https://osf.io/b4j3t>

(JMIR Preprints 15/02/2024:57224)

DOI: <https://doi.org/10.2196/preprints.57224>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/>



## Original Manuscript

# Predictors of Healthcare Practitioners' Intention to Use AI-Enabled Clinical Decision Support Systems (AI-CDSSs): A Meta-Analysis Based on the Unified Theory of Acceptance and Use of Technology (UTAUT)

## Abstract

**Background:** Artificial Intelligence-enabled Clinical Decision Support Systems (AI-CDSSs) offer potential for improving healthcare outcomes, but their adoption among healthcare practitioners remains limited.

**Objective:** The meta-analysis identifies predictors influencing healthcare practitioners' intention to use AI-CDSSs based on the Unified Theory of Acceptance and Use of Technology (UTAUT). Additional predictors are examined based on existing empirical evidence.

**Methods:** The literature search using electronic databases, forward searches, conference programs, and personal correspondence yielded 7,731 results, of which 17 studies met the inclusion criteria. Random-effects meta-analysis, relative weights analyses, and meta-analytic moderation and mediation analyses were used to examine the relationships of relevant predictor variables with the intention to use AI-CDSSs.

**Results:** The meta-analysis results supported the application of the UTAUT to the context of the intention to use AI-CDSSs. The results show that performance expectancy ( $r_c = .66$ ), effort expectancy ( $r_c = .55$ ), social influence ( $r_c = .66$ ), and facilitating conditions ( $r_c = .66$ ) were positively associated with the intention to use AI-CDSSs, in line with the predictions of the UTAUT. The meta-analysis further identified positive attitude ( $r_c = .63$ ), trust ( $r_c = .73$ ), anxiety ( $r_c = -.41$ ), perceived risk ( $r_c = -.21$ ), and innovativeness ( $r_c = .54$ ) as relevant additional predictors. Trust emerged as the most influential predictor overall. The results of the moderation analyses show that

the relationship between social influence and use intention becomes weaker with increasing age. In addition, the relationship between effort expectancy and use intention was stronger for diagnostic AI-CDSSs compared to devices that combined diagnostic and treatment recommendations. Finally, the relationships between facilitating conditions and use intention was mediated through performance and effort expectancy.

**Conclusions:** The meta-analysis contributes to the understanding of the predictors of the intention to use AI-CDSSs based on an extended UTAUT model. More research is needed to substantiate the identified relationships and to explain the observed variations in effect sizes by identifying relevant moderating factors. The research findings bear important implications for the design and implementation of training programs for healthcare practitioners to ease the adoption of AI-CDSSs into their practice.

**Keywords:** UTAUT; AI-CDSSs; meta-analysis; healthcare practitioners

## 1. Introduction

The past decade has witnessed major advancements in the field of healthcare, particularly through the integration of artificial intelligence (AI). AI may be described as machines that mimic cognitive functions associated with the human mind, such as learning and problem solving [1]. One area of progress involves the development of AI-enabled Clinical Decision Support Systems (AI-CDSS; [2–4]). AI-CDSSs use machine learning algorithms to process vast amounts of data and provide case-specific advice to healthcare practitioners to aid clinical decision-making [5–7]. AI-CDSSs use clinical data both from structured (e.g., lab results) and unstructured (e.g., clinician notes or imaging) sources. The interpretation of text-based data can be performed using Natural Language Processing (NLP) to transform text into usable data for clinical predictions [8]. Additionally, deep learning models, including neural networks, can be employed to generate recommendations based on image data, for example, in the detection of pneumonia from chest radiographs [9]. AI-CDSSs may improve the accuracy and efficiency of medical decision-making in several ways.

First, AI-CDSSs may offer structured rationales underpinning clinical decisions that can complement traditional care methods. This structured approach paves the way for clearer understanding, improved communication, and better tracking of the decision-making process in clinical settings [10,11]. Second, AI-CDSSs can integrate data from various sources to provide a comprehensive and personalized recommendation for every patient case [7]. Finally, AI-CDSSs promote the consistency of medical decisions. The use of AI algorithms may ensure that the same set of facts will consistently produce the same recommendations, thus minimizing harmful consequences due to human error [9].

Despite these advantages, the implementation of AI-CDSSs into clinical practice must still overcome numerous barriers. A major challenge in the deployment of AI-CDSSs is



the variability in performance. This can occur when the data used to develop the AI models does not adequately represent the population for which the tool is intended. Another issue is when AI-CDSSs are not used as designed, which can be due to a range of factors including user interface problems, lack of integration into clinical workflows, or insufficient training of healthcare professionals on how to use the system [7,12–14]. The resulting low performance casts doubt on the value of AI-CDSSs in assisting with clinical decision-making [12,15]. Additionally, the lack of understanding how AI recommendations are derived heightens clinicians' reservations about using these systems [16–18]. There are also challenges relating to the alignment of AI-CDSSs with existing workflows that can cause additional workload when new AI systems are incorporated into clinical procedures [7,19–21].

As the development of high-performing AI-CDSSs proceeds, understanding the factors that influence healthcare practitioners' intention to use these systems becomes increasingly relevant. One of the most comprehensive theories to explain individual technology adoption is the Unified Theory of Acceptance and Use of Technology (UTAUT; [22]). The UTAUT proposes that a person's intention to use a technology is determined by their beliefs and attitudes towards that technology, such as the perception of its performance or the perceived effort it would require to use it. The UTAUT's comprehensive nature and its ability to account for various determinants of technology acceptance make it an appropriate model for examining the predictors of healthcare practitioners' intention to use AI-CDSSs.

Research to identify predictors of the intention to use AI-CDSSs has accumulated over the past years [4,23–25]. However, the existing literature remains scattered and in need of systematic synthesis. The overarching goal of this article, therefore, is to quantitatively integrate existing studies on the predictors of healthcare practitioners'

intention to use AI-CDSSs. The proposed hypotheses are based on the UTAUT model and existing empirical evidence. With the current meta-analysis, we make four major contributions to theory and practice. First, we use meta-analytic techniques to estimate the relationship between the predictors of the UTAUT and the intention to use AI-CDSSs, thus providing insight into the applicability of the UTAUT to the context of AI-CDSSs. Second, we identify additional predictors based on the existing literature and examine the relative contribution of the UTAUT and additional predictors in explaining the intention to use AI-CDSSs. With this approach, we contribute to a theoretical refinement and potential extension of the UTAUT model to the context of AI-CDSSs. Third, based on the UTAUT, we examine the role of contextual factors as moderators of the relationships between relevant predictors and use intention, thus shedding light on the conditions that influence the strength of these relationships. Finally, in line with the UTAUT model, this is the first meta-analysis that examines the role of mediators, thus allowing for a better understanding of the complex mechanisms through which use intention may be explained. The study protocol, including all hypotheses and research questions, has been pre-registered through the Open Science Framework (<https://osf.io/b4j3t>).

## **2. Theory and Hypothesis Development**

### **2.1 The Unified Theory of Acceptance and Use of Technology (UTAUT) and the Intention to Use AI-CDSSs**

The UTAUT integrates eight former technology use theories and has become one of the most prominent technology use models [22,26]. The UTAUT has been applied to investigate factors influencing the acceptance and use of technology in different contexts, including healthcare [27–29]. The primary outcome measure considered in the UTAUT, alongside actual use, is the intention to use a technology [22,30,31]. Intentions are indicators of motivation and reflect the level of determination individuals have to actually

perform a certain behavior [32]. The successful deployment of any technology depends largely on the user's intention to use it [33]. Accordingly, understanding the predictors of the intention to use AI-CDSSs may help overcome individual-level impediments thwarting the adoption of AI-CDSSs in healthcare.

The UTAUT consists of four core predictors of individual use intention: Performance expectancy, effort expectancy, social influence, and facilitating conditions [22]. The relationships between these variables and use intention are proposed to be moderated by gender, age, experience, and voluntariness of use [22]. The UTAUT model is shown in Figure 1. All relationships included in the UTAUT are proposed as hypotheses, while all additional relationships and moderators that are derived based on empirical findings and other theories are proposed as research questions.




Figure 1. *The UTAUT model*. Adapted from Venkatesh et al. (2003), User Acceptance of Information Technology: Toward a Unified View, *MIS Quarterly*, 27(3), p. 447.

## 2.2 Predictors of the Intention to Use AI-CDSSs Based on the UTAUT

Performance expectancy refers to the extent to which individuals believe that using a

technology will improve their job performance. AI-CDSSs have the potential to enhance job performance by aiding clinicians in deriving diagnoses or making treatment decisions [34]. If clinicians perceive their decisions to be improved by using AI-CDSSs, then performance expectancy will be high [35,36].

*Hypothesis 1:* Performance expectancy is positively related to the intention to use AI-CDSSs.

Effort expectancy concerns the perceived ease of use of a technology. It is suggested that a system that is perceived to be easy to use is more likely to be accepted than one that is perceived to be complicated to use [22]. If, for example, the perceived effort of using an AI-CDSS in one's existing clinical workflows is perceived to be high, healthcare practitioners may be less willing to use it [3,21,23].

*Hypothesis 2:* Effort expectancy is positively related to the intention to use AI-CDSSs.

Social influence refers to the impact of social factors, such as the expectations and influence of peers, on an individual's intention to use a technology. The positive relationship between social influence and the intention to use AI-CDSSs has consistently been supported in empirical studies [23,36]. For example, it was found that medical professionals holding the belief that their colleagues, top management, and professional bodies endorse the use of AI-CDSSs in clinical settings are more willing to adopt them [36].

*Hypothesis 3:* Social influence is positively related to the intention to use AI-CDSSs.

Facilitating conditions represent the organizational and technical infrastructure necessary for technology adoption [22]. It has been argued that if users believe that the resources and support are in place to facilitate the use of AI-CDSSs, they are more likely to intend to use them [4,22,37].

*Hypothesis 4a:* Facilitating conditions are positively related to the intention to use AI-

CDSSs.

In addition, according to the UTAUT, there is a direct relationship between facilitating conditions and actual technology use [22]. Facilitating conditions refer to the resources and support available to use a technology, including the access to the necessary tools and knowledge. This practical aspect makes their influence on usage immediate, as users are more likely to use technology when they perceive a supportive environment and available resources. Unlike other predictors in the UTAUT, facilitating conditions are proposed as direct antecedents of actual use [22].

*Hypothesis 4b:* Facilitating conditions are positively related to actual use of AI-CDSSs.

### **2.3 Additional Predictors of the Intention to Use AI-CDSSs**

The UTAUT has been modified and additional predictors have been added over time to account for various settings and technologies [30,38–40]. However, a meta-analytic review is limited to the relationships that have been studied in a literature. Following previous research and methodological best practices, we include additional predictors beyond the UTAUT in the meta-analysis that have been examined in at least three independent samples [41,42]. Following this criterion, we identified attitude, trust, perceived risk, AI anxiety, and personal innovativeness as additional predictors of the intention to use AI-CDSSs.

Individual behavior is driven by intention, which is in turn a function of an individual's attitude toward the behavior and subjective norms [30,43]. Indeed, a positive attitude towards AI-CDSSs has been identified as a relevant predictor of the intention to use AI-CDSSs [44–46]. Because the relationship between positive attitude and use intention is not included in the UTAUT, we propose the following research question to explore the relationship between positive attitude and the intention to use AI-CDSSs [22].

*Research Question 1:* Is there a positive relationship between a positive attitude towards AI-CDSSs and the intention to use AI-CDSSs?

Trust becomes relevant if the outcome of a situation is uncertain or the possibility of undesirable outcomes exists [46]. Trust has been argued to be a particularly relevant predictor of the intention to use AI-CDSSs due to a lack of transparency of how recommendations are derived and the high stakes of erroneous decisions in healthcare [23,36]. Generally, we may differentiate between initial trust as the judgement of the trustor before being exposed to the trustee and knowledge-based trust that may be established after the trustor has interacted with the trustee [46]. In the context of AI-CDSSs, some studies refer to initial trust in terms of beliefs in the reliability and safety in AI-CDSSs before the user has been exposed to or actively used the system [2,35,47]. An example item for initial trust is “I believe AI could provide completely accurate diagnosis assistant service” [3]. Another aspect of trust that has been explored in empirical studies is trust in different attributes of the technology, namely its functionality as being able to do a required task, its helpfulness or benevolence as being able to provide effective help when needed or act in the best interest of the user, and its integrity as operating reliably or consistently without failing [46,48,49]. One example item for trust in the system's integrity regarding data security is “I trust that recommendations from the AI-powered care pathway are reliable” [50]. Because trust is not included in the UTAUT model, we propose a research question to explore whether there is a positive relationship between trust and the intention to use AI-CDSSs [22].

*Research Question 2:* Is there a positive relationship between trust and the intention to use AI-CDSSs?

Perceived risk is determined by the unpredictability and perceived intensity of outcomes [50]. In the context of AI-CDSSs, perceived risk refers to the perceived potential

negative consequences associated with their use, including performance failure and data insecurity [4]. An example item for perceived risk of a performance failure is “There is a possibility of malfunction and performance failure, so the system might fail to deliver accurate contouring areas and could mislead my work with inaccurate contouring” [4]. Healthcare professionals may be reluctant to engage with new services fearing that their perceived risk may result in negative user experience or even harm for them or their patients [51]. Different forms of perceived risk have been found to be negatively associated with the intention to use AI-CDSSs [3,35,51,52]. For example, it has been found that performance and legal risk associated with an AI-CDSSs were negatively related to the intention to use AI-CDSSs [36]. Because perceived risk is not included in the UTAUT model, we propose the following research question to investigate whether perceived risk is negatively associated with the intention to use AI-CDSSs [22].

*Research Question 3:* Is there a negative relationship between perceived risk and the intention to use AI-CDSSs?

AI anxiety encompasses general fears and insecurities regarding AI technology. It represents an intuitive, negative affective reaction to AI technologies, for example, based on the fear of making mistakes [53,54]. AI anxiety is often measured with the AI anxiety scale [22]. An example item is “I feel apprehensive about using the system”. If healthcare professionals experience anxiety in using AI-CDSSs, their intention to use them is presumably low. Indeed, AI anxiety has been identified as a negative predictor of the intention to use AI in healthcare [24]. However, because AI anxiety is not included as a predictor of use intention in the UTAUT, we propose the following research question to explore whether AI anxiety is negatively associated with the intention to use AI-CDSSs [22].

*Research Question 4:* Is there a negative relationship between AI anxiety and the intention to use AI-CDSSs?

Personal innovativeness describes an individual's readiness to experiment with and embrace a new technology [55]. Those demonstrating a high degree of personal innovativeness have greater capabilities and, therefore, demonstrate greater readiness to use a new technology [56,57]. Indeed, there exists empirical evidence for a positive link between personal innovativeness and the intention to use AI-CDSSs [3,35].

*Research Question 5:* Is there a positive relationship between personal innovativeness and the intention to use AI-CDSSs?

## **2.4 The Relationship Between AI-CDSSs Use Intention and Actual Use**

The UTAUT proposes that an individual's intention to use a technology is the main predictor of its actual use [22]. However, this relationship has not yet been extensively researched in the context of AI-CDSSs. The limited investigation of actual use may be attributed to the restricted number of AI-CDSSs implemented in clinical practice [58]. Nonetheless, some evidence indicates that use intention predicts the actual use of AI-CDSSs [4,46].

*Research Question 6:* What is the relationship between the intention to use AI-CDSSs and their actual use?

## **2.5 The Relative Contribution of the UTAUT Predictors and Additional Predictors in Explaining AI-CDSSs Use Intention**

Existing empirical research has explored the extent to which the UTAUT predictors account for variance in technology use intention [59]. For example, performance expectancy has often emerged as the strongest predictor of use intention [60–62]. Other research has found that trust has a stronger effect on the intention to use AI-CDSSs than performance expectancy [36]. As the roles of UTAUT and additional predictors in explaining the intention to use AI-CDSSs remain unclear, we propose the following research question:

*Research Question 7:* What is the relative contribution of the UTAUT predictors and



additional predictors in explaining the intention to use AI-CDSSs?

## **2.6 Moderators of the Relationships Between UTAUT Predictors and the Intention to Use AI-CDSSs**

The relationships between UTAUT predictors and use intention are proposed to be moderated by age, gender, user experience with the system, and voluntariness of using the system [22]. First, it has been suggested that younger workers prioritize extrinsic rewards such as improved job performance, thus exhibiting a stronger relationship between performance expectancy and technology use intention [22]. In contrast, it is suggested that older workers generally face greater software challenges and are more likely to place increased relevance on social influences. Accordingly, they may rely more on effort expectancy and social influence when deciding to use a technology [22].

*Hypothesis 5:* The relationship of (a) performance expectancy with the intention to use AI-CDSSs becomes weaker and the relationships of (b) effort expectancy and (c) social influence with the intention to use AI-CDSSs become stronger with increasing age.

Second, the impact of performance expectancy on use intention is expected to be stronger among men, while the relationship of effort expectancy and social influence with use intention would be more pronounced among women [22].

*Hypothesis 6:* The relationship of (a) performance expectancy with the intention to use AI-CDSSs is stronger for men, and the relationships of (b) effort expectancy and (c) social influence with the intention to use AI-CDSSs is stronger for women.

Third, according to the UTAUT, limited experience increases the strength of the relationship between effort expectancy and social influence with use intention because individuals with limited experience tend to overestimate the challenges associated with using a new technology and their opinions are more susceptible to social influence [22]. In contrast, as experience increases, facilitating conditions have been proposed to exhibit a

greater impact on actual technology use, as more experienced users know better how to take advantage of facilitating conditions when using the system [22].

*Hypothesis 7:* The relationships of (a) effort expectancy and (b) social influence with intention to use AI-CDSSs become weaker with increasing experience and the relationship of (c) facilitating conditions with actual use of AI-CDSSs becomes stronger with increasing experience.

Finally, the UTAUT distinguishes between voluntary (i.e., individuals decide themselves whether to use a technology) and mandatory (e.g., the use of a technology is mandated by the supervisor) adoption settings [22]. It has been suggested that social influence affects use intention in mandatory situations more because relevant others have the capacity to either incentivize desired actions or penalize noncompliance [22].

*Hypothesis 8:* The relationship of social influence with the intention to use AI-CDSSs is stronger in mandatory adoption settings.

In addition to the UTAUT moderators, we investigate the influence of additional contextual moderators that are studied in the literature, namely occupation, type of AI-CDSS, and culture. All additional moderators were selected based on a comprehensive preliminary review of the literature. First, healthcare practitioners may work in different contexts requiring them to complete different tasks. These differences may influence their perceptions, beliefs, and attitudes towards AI-CDSSs [23,51]. For instance, one study found differences in the relationship of social influence and perceived risk with use intention between clinicians (e.g., surgery, orthopedics) and non-clinicians (e.g., radiologists, pathologists). Specifically, for non-clinicians, social influence positively predicted the intention to use AI-CDSSs, while perceived risk did not emerge as a significant predictor. In contrast, among clinicians, the reverse pattern was observed [52]. Second, the type of AI-CDSS likely influences practitioners' use intention. Specifically, healthcare practitioners may

place greater emphasis on the effectiveness and safety of treatment AI-CDSSs compared to diagnostic AI-CDSSs as an erroneous treatment decision is associated with more severe consequences [24]. Finally, cultural differences may influence the intention to use AI-CDSSs in healthcare [63,64]. For example, one study found perceived ease of use to be a more relevant predictor of the intention to use information technology among Taiwanese compared to US physicians [64]. Accordingly, we propose the following research question:

*Research Question 8:* Do (a) the practitioner's occupation, (b) the type of AI-CDSS, and (c) cultural background moderate the relationship between UTAUT predictors and the intention to use AI-CDSSs?

Finally, we investigate the influence of methodological moderators such as publication year and the scale used to measure AI-CDSSs use intention. In a meta-analysis based on the UTAUT, it was found that some effect sizes were stronger in more recent studies [59]. Moreover, while most studies use the intention to use scale introduced by Venkatesh et al. (2003), some studies employ self-developed scales to measure use intention [25,35].

*Research Question 9:* Do (a) publication year and (b) the use intention scale employed moderate the relationship between UTAUT predictors and the intention to use AI-CDSSs?

## **2.7 Performance and Effort Expectancy as Mediators of the Relationship Between Facilitating Conditions and the Intention to Use AI-CDSSs**


According to the UTAUT, the effect of facilitating conditions on use intention may be explained through performance and effort expectancy [21,65]. That is, if the required support infrastructure is provided, a person would perceive the system to be both high-performing and easy to use, which, in turn, positively influences their intention to use it. Indeed, effort expectancy has been found to fully mediate the relationship between

facilitating conditions and use intention [65]. Accordingly, we propose the following research question to investigate the mediating role of performance and effort expectancy:

*Research Question 10:* Is the relationship between facilitating conditions and the intention to use AI-CDSSs mediated through (a) performance and (b) effort expectancy?

## 2.8 Overview of the Hypotheses and Research Questions

Figure 2 displays all hypotheses and research questions. We omitted the relationship between facilitating conditions and actual use of AI-CDSSs (Hypothesis 4b) as well as the moderators experience (Hypothesis 7), voluntariness (Hypothesis 8), and occupation (Research Question 8a) from the analyses (see dashed lines in Figure 2) due to the limited number of available independent samples ( $k < 3$ ). All deviations from the pre-registration are presented in Table S1 in the Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16))



**Figure 2.** *The proposed research model.* Dashed lines represent pre-registered hypotheses and research questions that could not be investigated due to the limited number of available independent samples ( $k < 3$ ). Research Question 7 is omitted from the Figure as it refers to the relative weights analysis.

### 3. Methods

#### 3.1 Inclusion and Exclusion Criteria

To qualify for inclusion, the following criteria had to be met: First, studies had to be published in English. Second, studies had to include AI-CDSSs. The second inclusion criterion is fulfilled if a) one of the terms ‘artificial intelligence’, ‘AI’, ‘machine learning’, ‘deep learning’, or ‘deep neural networks’ was used to describe the technology [66] and b) the technology was referred to as a CDSS or it was described as providing recommendations regarding the diagnosis, treatment, or prognosis of health issues [6]. We included studies if AI-CDSSs were mentioned alongside other AI-enabled functionalities [67]. This led to the exclusion of studies that investigated the use intention of other healthcare technologies, such as telemedicine [68] or the medical internet of things [69]. Notably, one study examined the intention to use explainable and non-explainable AI-CDSSs in the same sample [37]. Because only one other study examined explainable AI [44], we included only the data for the non-explainable AI-CDSSs. Third, studies had to include a measure of the intention to use AI-CDSSs as defined in the UTAUT [22], including self-developed scales based on the UTAUT scale. Fourth, studies had to be empirical. This led to the exclusion of non-empirical studies, such as reviews or case studies [70]. Fifth, studies had to measure at least one predictor of the intention to use AI-CDSSs. Sixth, studies had to measure use intention among a sample of healthcare practitioners or medical students based on the list of health professionals by the World Health Organization [71]. For a detailed overview of the inclusion criteria per included study, please refer to Table S2 in the Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)).

#### 3.2 Search Strategy and Data Extraction

This meta-analysis was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines to ensure comprehensive and transparent reporting [72]. We used five steps to search for relevant data. First, relevant scientific articles, dissertations, and theses were searched using the electronic databases Embase, Medline, ProQuest, PsycINFO, and Web of Science between October 15th, 2022, and January 5th, 2023. Two follow-up searches were conducted on May 2nd, 2023 and November 7th, 2023. The search string was developed based on the participants, intervention, comparators, and outcome framework (PICO) [73]. The framework was adapted to fit the research purpose resulting in a three-tiered search term including the population (healthcare professionals), the technology (AI-CDSSs), and the outcome (use intention) of interest. An overview of the search terms is presented in Table S3 the Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)). We used the search term to search in titles, abstracts, and keywords. We conducted follow-up searches in Google Scholar using the following search string: ['healthcare'], AND ['Artificial Intelligence'] AND ['UTAUT']. Second, we conducted forward-searching of studies citing Venkatesh et al.'s (2003) seminal article via Google Scholar and backward searches in review articles [74–78]. Third, abstracts of relevant conference proceedings including the *Conference on Computer Supported Cooperative Work and Social Computing*, the *Conference on Human Factors in Computing Systems*, and the *Institute of Electrical and Electronics Engineers* were searched. Fourth, we sent requests for unpublished articles and data using the mailing list of the German Psychology Association (DGPs). Finally, authors of articles included in the meta-analysis were contacted and asked for unpublished data sets. No additional unpublished data was obtained.

We reached out to authors when critical information needed to decide the inclusion of a study or details essential for the meta-analytic synthesis, such as a correlation table,

were missing. From the 24 authors contacted to procure missing information, we successfully obtained six data sets. These data sets were used to derive the missing information, for instance, to calculate missing correlations between variables of interest.

Figure 3 shows the PRISMA diagram with the number of studies identified, included, and excluded, along with reasons for exclusion. The studies from the literature search were assessed following a three-stage approach. First, titles were screened to identify relevant articles. Second, the abstracts of the remaining articles were reviewed. Third, full article texts were reviewed. As a result of a review of 107 full texts, 17 studies met the inclusion criteria ( $K = 18$  independent samples,  $N = 3,871$ ).

Following the approach of previous meta-analyses, we only included relationships that were identified in a minimum of three separate samples [41,79]. We grouped overlapping variables into construct categories (see Table S4 in the Online Appendix for definitions of superordinate constructs and subconstructs, [https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)). Studies from both the primary and the follow-up literature search were coded by two researchers each (AK and SG for the primary search; and JD and AK for the follow-up search). Any conflicts in the coding were resolved in weekly consensus meetings. Additionally, in line with approaches to ensure accuracy in coding established in previous meta-analyses [80], a random sample of ten of the 18 independent samples (56%) was re-coded by JC and AS. We included agreement on correlations, reliabilities, and moderator categories into the assessment of interrater agreement. Overall interrater agreement was high (94.7%). Notably, no disagreements were observed regarding correlations. Some mistakes in the coding of reliabilities occurred during the re-coding due to referencing an incorrect line from the source document. The final code sheet used for the analyses is published in the OSF Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)).




Figure 3. PRISMA flow chart of the study selection process



### 3.3 Meta-analytic procedures

All analyses were conducted with RStudio [81] using the R packages 'psychmeta' [82] and 'metaSEM' [83].

#### 3.3.1 Bivariate Relationships

To examine the bivariate relationship of the four core constructs of the UTAUT (H1-H4) and the additional predictors (RQ1-RQ5) with the intention to use AI-CDSSs, random-effects meta-analysis was conducted [84]. Effect sizes were based on Pearson product-moment correlations. Composites were calculated if multiple measures of the same construct were reported for the same sample [84]. Specifically, a variance-weighted composite (across measures of the same construct) was calculated for each independent sample to combine multiple measures of the same construct into a single effect size per independent sample [84]. Sampling errors were corrected using sample size weighted correlations. Measurement errors were corrected based on Cronbach's alpha [84]. In addition to the sample size-weighted correlation ( $r$ ) and sample size-weighted and reliability-corrected correlation ( $r_c$ ), the 95% confidence interval (95% CI) and 80% credibility interval (80% CR) for ( $r_c$ ) are reported. Finally, we report the correlation between observed effects and the influence of the study design artifacts ( $\text{cor}(r, a)$ ).

#### 3.3.2 Relative Weights Analysis

We conducted relative weights analyses to capture the contribution of the correlated predictors [85]. Specifically, we calculated multivariate meta-analytic regression models based on the pooled correlation matrices to explore the incremental value of the UTAUT predictors and additional predictor variables in explaining the intention to use AI-CDSSs. We used the harmonic mean of the sample size across the correlations considered as the sample size for the estimated regression models [86]. In relative weights analysis, raw relative weights are calculated to reflect the proportion of variance explained in the outcome

that is attributed to each of the predictors, while rescaled relative weights reflect the percentage of variance that is explained by each predictor variable [87,88].

### 3.3.3 Moderation Analyses

Moderator analyses were carried out for constructs that were represented in a minimum of  $k = 10$  independent samples to ensure adequate coverage of moderator categories [89]. Five constructs met this minimum cutoff and were considered for the moderation analyses (i.e., performance expectancy, effort expectancy, social influence, trust, and perceived risk). We interpreted categorical moderator effects if each of the levels included  $k \geq 3$  independent samples. Age was coded as the mean age of study participants, and gender as the percentage of females in the sample. For the type of AI-CDSS, three categories were initially identified: diagnostic decision support systems, treatment decision support systems, and systems that combined both diagnostic and treatment decision support. However, the category treatment decision support systems had to be excluded from the moderator analysis because of the low number of independent samples focusing on this kind of AI-CDSS ( $k = 2$ ). Culture was operationalized based on the individualism versus collectivism dimension of the Hofstede country comparison tool [90,91]. A higher score denotes stronger individualism. The publication year was coded chronologically. Finally, the scale used to measure the intention to use AI-CDSSs was coded as a categorical moderator. We differentiated between studies using the Venkatesh scale and studies using self-developed scales. We conducted moderation analyses that were not pre-registered as part of exploratory analyses.

### 3.3.4 Mediation Analysis

To test Research Question 6, correlation-based meta-analytic structural equation modeling (MASEM) [92] based on the two-stage structural equation modeling approach (TSSEM) [93,94] was performed. In the first step, the sample-size weighted and reliability-

corrected bivariate correlation matrices for each independent sample were pooled together. In TSSEM, the total sample size is used for the estimation of the MASEM model [94]. In the second step, a path model was fitted to the pooled correlation matrix.

## 4. Results

### 4.1 Study Characteristics

The overall mean age was 36.2 years ( $SD = 13.5$  years, range = 21-53 years) and 48.7% were female. Seven out of the 17 studies focused on diagnostic AI-CDSSs, two on treatment AI-CDSSs, four on treatment and diagnostic AI-CDSSs, and four on unspecific AI-CDSSs. Eleven studies were conducted in Asia (six in China), three in Europe, one in the U.S., and two globally in English-speaking countries.

### 4.2 Meta-Analytic Results

In the following, we report sample size-weighted and reliability-corrected correlations ( $r_c$ ) for the relationships of relevant antecedent variables with AI-CDSSs use intention. In line with Cohen (1988), we classify our reported effects as weak ( $r_c = .1$ ), moderate ( $r_c = .3$ ), and strong ( $r_c = .5$ ).

#### 4.2.1 Bivariate Relationships

The results of bivariate meta-analytic analyses are shown in Table 1. The UTAUT predictors performance expectancy ( $r_c = .66$ , 95% CI [.59, .73]), effort expectancy ( $r_c = .55$ , 95% CI [.43, .67]), social influence ( $r_c = .66$ , 95% CI [.59, .72]), and facilitating conditions ( $r_c = .66$ , 95% CI [.42, .90]) exhibited a strong positive relationship with the intention to use AI-CDSSs. The findings support Hypotheses 1 through 3 and 4a. The relationship between facilitating conditions and actual use was not investigated in a sufficient number of independent samples ( $k > 2$ ). Accordingly, we could not address Hypothesis 4b. Regarding the additional predictors beyond the UTAUT, attitude ( $r_c = .63$ , 95% CI [.52, .73]), trust ( $r_c = .73$ , 95% CI [.63, .82]) and innovativeness ( $r_c = .54$ , 95% CI [.43, .64]) exhibited strong

positive relationships, confirming Research Questions 1, 2, and 5. Perceived risk ( $r_c = -.21$ , 95% CI  $[-.35, -.08]$ ) was weakly negatively related to use intention, supporting Research Question 3. Although the estimate for AI anxiety was strong and negative ( $r_c = -.41$ ), the 95% CI included zero (95% CI  $[-.98, .15]$ ). Accordingly, we cannot conclude that AI anxiety is related to use intention, thus not supporting Research Question 4. The 80% credibility interval for effort expectancy (.27 to .83), facilitating conditions (.33 to .99) and AI anxiety (-.81, -.01) were wide, suggesting the presence of moderators [84,95]. Finally, the intention to use AI-CDSSs was strongly positively related to the actual use of AI-CDSSs, confirming Research Question 6 ( $k = 3$ ,  $N = 478$ ,  $r = .75$ ,  $r_c = .85$ ,  $SD_c = 0.09$ , 95% CI  $[.63, 1.00]$ , 80% CR  $[.70, 1.00]$ ,  $cor(r,a) = .44$ ).

**Table 1**

*Bivariate relationships between predictor variables and AI-CDSS use intention*

Predictor Variable	$k$	$N$	$r$	$r_c$	$SD_c$	95% CI	80% CR	$cor(r,a)$
Performance expectancy	16	3,295	.59	.66	0.13	.59, .73	.50, .82	.39
Effort expectancy	15	3,058	.49	.55	0.22	.43, .67	.27, .83	.28
Social influence	15	3,058	.57	.66	0.12	.59, .72	.52, .80	.46
Facilitating conditions	6	1,048	.57	.66	0.23	.42, .90	.33, .99	.25
Attitude	9	2,048	.51	.63	0.14	.52, .73	.45, .80	.43
Trust	10	1,840	.66	.73	0.13	.63, .82	.55, .90	.35
Perceived risk	10	2,428	-.19	-.21	0.18	-.35, -.08	-.45, .02	.39
Anxiety	3	391	-.37	-.41	0.23	-.98, -.15	-.81, -.01	.38
Innovativeness	5	843	.47	.54	0.09	.43, .64	.46, .61	.81

Note.  $k$  = number of independent samples;  $N$  = cumulative sample size;  $r$  = sample size-weighted correlation;  $r_c$  = sample size-weighted and reliability-corrected correlation;  $SD_c$  = standard deviation of  $r_c$ ; CI = confidence interval for  $r_c$ ; CR = credibility interval;  $cor(r, a)$  = correlation between  $r$  and statistical artifacts ( $a$ ).

#### 4.2.2 Relative Weights Analysis

It was not possible to explore all nine predictors in a single relative weights analysis because they were not investigated together in a sufficient number of independent samples (see Table S5 in the Online Appendix, <https://osf.io/djwkz/?>

view\_only=7d6046aac9424a3c8f2facab08ac1e16). Accordingly, to answer Research Question 7, we analyzed one model with only the UTAUT predictors (Table 2) and four separate extension models, consisting of five to six predictors (Table 3). In the initial model with only the UTAUT predictors, the combined effects of performance expectancy, effort expectancy, social influence, and facilitating conditions explained 50% of the total variance in the intention to use AI-CDSSs. Performance expectancy was the dominant predictor accounting for 31% of the total variance explained, followed by social influence (28%), facilitating conditions (26%), and effort expectancy (15%). In the extension models, trust emerged as the most influential overall predictor of use intention (between 29% and 35% of the total variance explained). In all three models including trust, performance expectancy was the second most influential predictor (between 19% and 24% of the total variance explained). Facilitating conditions (between 20% and 25%) and social influence (between 14% and 21%) consistently explained additional variance in all extension models. In the extension models including trust and perceived risk, and trust and anxiety, the regression estimate of effort expectancy became negative. Finally, AI anxiety and perceived risk negatively predicted use intention and accounted for 10% (AI anxiety) and 2% (perceived risk) of the total variance explained.

**Table 2**

*Multiple regression models and relative weights for the UTAUT predictors*

Predictor	<i>B</i>	<i>SE</i>	<i>t</i> -value	<i>P</i>	Raw RW	RS RW
Performance expectancy	.31	0.02	13.97	<.001	.16	31.19%
Effort expectancy	.08	0.02	3.56	<.001	.08	15.20%
Social influence	.27	0.02	12.29	<.001	.14	27.91%
Facilitating conditions	.21	0.02	9.33	<.001	.13	25.70%

*Note.* *B* = regression estimate; *SE* = standard error of *B*; RW = relative weight; RS = rescaled.

$F = 429.28$  ( $p < .001$ ),  $R^2 = .498$ .

**Table 3**

*Multiple regression models and relative weights for the UTAUT and additional predictors*

Predictor	<i>B</i>	<i>SE</i>	<i>t</i> -value	<i>P</i>	Raw RW	RS RW
UTAUT extension (attitude, perceived risk), $F = 222.31$ ( $p < .001$ ), $R^2 = .509$						
Performance expectancy	.25	0.03	9.36	<.001	.12	24.00%
Effort expectancy	.05	0.03	2.04	.04	.06	12.14%
Social influence	.17	0.03	6.40	<.001	.10	20.54%
Facilitating conditions	.28	0.03	10.82	<.001	.13	25.31%
Attitude	.13	0.03	5.02	<.001	.08	15.91%
Perceived risk	-.04	0.02	-2.20	.03	.01	2.09%
UTAUT extension (trust, innovativeness), $F = 308.50$ ( $p < .001$ ), $R^2 = .542$						
Performance expectancy	.22	0.03	8.77	<.001	.12	22.72%
Effort expectancy	.05	0.02	2.14	.03	.06	11.57%
Social influence	.19	0.03	7.62	<.001	.11	20.40%
Trust	.39	0.03	15.40	<.001	.19	35.04%
Innovativeness	.04	0.02	1.56	.12	.06	10.26%
UTAUT extension (trust, perceived risk), $F = 389.61$ ( $p < .001$ ), $R^2 = .600$						
Performance expectancy	.18	0.02	8.40	<.001	.11	18.76%
Effort expectancy	-.06	0.02	-2.65	.01	.05	8.81%
Social influence	.09	0.02	3.87	<.001	.09	15.66%
Facilitating conditions	.32	0.02	14.99	<.001	.13	22.03%
Trust	.42	0.02	19.79	<.001	.20	33.00%
Perceived risk	-.05	0.02	-2.80	.01	.01	1.74%
UTAUT extension (trust, anxiety), $F = 241.15$ ( $p < .001$ ), $R^2 = .632$						
Performance expectancy	.23	0.03	8.15	<.001	.12	19.25%
Effort expectancy	-.11	0.03	-3.92	<.001	.05	7.24%
Social influence	.07	0.03	2.44	.02	.09	14.13%
Facilitating conditions	.31	0.03	11.43	<.001	.13	20.44%
Trust	.38	0.03	13.48	<.001	.18	28.68%
Anxiety	-.20	0.02	-8.73	<.001	.06	10.26%

Note. *B* = regression estimate; *SE* = standard error of *B*; RW = relative weight; RS = rescaled.

### 4.2.3 Moderation Analyses

Table 4 shows the results of the meta-regression for continuous moderators. Regarding age, older participants revealed a weaker relationship between social influence and use intention ( $B = -.01$ , 95% CI  $[-.01, <-.01]$ ), thus *contradicting* Hypothesis 5c

according to which this effect would become stronger with increasing age. The moderation effect is displayed in Figure S1 in the Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)). Age did not moderate any other relationship, thus not confirming Hypotheses 5a and 5b. Gender did not moderate any of the relationships, thus not confirming Hypothesis 6a, 6b, and 6c. Experience and voluntariness of use were not investigated in a sufficient number of independent samples. Accordingly, we were unable to address Hypotheses 7 and 8. Cultural individualism (Research Question 8c) as a contextual moderator that was measured continuously did not influence any of the relationships. Finally, publication year (Research Question 9a) as a methodological moderator that was measured continuously did not influence any of the relationships.

**Table 4**

*Results of meta-regression*

Predictor variable	Moderator	$k_{mod}$	$B$	$SE$	$P$	95% CI
Performance expectancy	Age	4	<.01	<0.01	.56	-.01, .01
	Gender	16	<.01	<0.01	.88	<-.01, <.01
	Individualism	14	<.01	<0.01	.66	<-.01, <.01
	Publication year	16	.02	0.03	.42	-.03, .07
Effort expectancy	Age	4	<.01	0.01	.97	-.02, <.01
	Gender	15	<.01	<0.01	.63	-.01, <.01
	Individualism	13	<.01	<0.01	.95	<-.01, <.01
	Publication year	15	.03	0.04	.68	-.06, .09
Social influence	Age	4	-.01	<0.01	.03	-.01, <-.01
	Gender	15	<.01	<0.01	.09	<-.01, .01
	Individualism	13	<.01	<0.01	.56	<-.01, <.01
	Publication year	15	.02	0.02	.21	-.02, .07
Trust	Age	3	<.01	0.01	.90	-.02, .02
	Gender	10	<.01	<0.01	.88	<-.01, <.01
	Individualism	9	<.01	<0.01	.91	<-.01, <.01
	Publication year	10	-.04	0.03	.24	-.10, .03
Perceived risk	Gender	10	<.01	<0.01	.64	-.01, .01
	Individualism	9	<.01	<0.01	.21	<-.01, <.01

Publication year	8	-.02	0.04	.60	-.11, .06
------------------	---	------	------	-----	-----------

Note. Moderator analysis for predictor variables with  $k \geq 10$  independent samples.  $k_{mod}$  = number of independent samples per moderator;  $B$  = regression estimate;  $SE$  = standard error; CI = confidence interval; Age = chronological age in years; Gender = Percentage female.

The Wald-type pairwise comparisons for each level of categorical moderators are presented in Table 5. We could not investigate Research Question 8a because information about occupations was not provided in a sufficient number of independent samples. Regarding Research Question 8b, there were no differences for the type of AI-CDSS (diagnostic AI-CDSSs vs. diagnostic and treatment AI-CDSSs) on the relationships of performance expectancy and social influence. However, the positive relationship of effort expectancy on use intention was stronger for diagnostic AI-CDSSs compared to AI-CDSSs that combined diagnostic and treatment recommendations (Diff = -.31, 95% CI [-.58, -.04]). Finally, regarding Research Question 9b, we observed no differences between studies using the Venkatesh et al. (2003) scale and those employing other measures.



**Table 5***Wald-type pairwise comparisons of categorical moderators*

Predictor variable	$k_1$	$k_2$	$F$	$r_{c1}$	$r_{c2}$	Diff.	95% CI
AI-CDSSs type: <i>Diagnostic and treatment</i> AI-CDSSs (level 1) compared to <i>diagnostic</i> AI-CDSSs (level 2)							
Performance expectancy	4	7	1.48	.62	.71	-.09	-.31, .13
Effort expectancy	4	6	7.15	.41	.72	-.31	-.58, -.04
Social influence	4	6	6.97	.59	.72	-.14	-.28, <.01
Trust	3	4	0.09	.70	.72	-.02	-.28, .24
Use intention scale: <i>Other scales</i> (level 1) compared to <i>Venkatesh et al. (2003)</i> (level 2)							
Performance expectancy	10	6	0.07	.66	.68	-.02	-.18, .15
Effort expectancy	9	6	0.48	.59	.51	.08	-.18, .35
Social influence	9	6	0.82	.64	.70	-.06	-.21, .09
Trust	5	5	0.01	.73	.72	.01	-.19, .20
Perceived risk	7	3	2.41	-.16	-.37	.21	-.10, .53

Note. Moderator analysis for constructs with at least  $k = 10$  independent samples.  $k_1$  and  $k_2$  = number of independent samples for moderator level 1 and 2;  $r_{c1}$  and  $r_{c2}$  = sample size-weighted and reliability-corrected correlation for moderator level 1 and 2; Diff. = Mean difference, CI = confidence interval.

#### 4.2.4 The Mediating Role of Performance and Effort Expectancy in the Relationship Between Facilitating Conditions and AI-CDSSs Use Intention

The role of performance and effort expectancy as mediators of the relationship between facilitating conditions and the intention to use AI-CDSSs (Research Question 10a and 10b) were analyzed by fitting two separate mediation models. The results are displayed in Table 6. Performance expectancy mediated the relationship between facilitating conditions and the intention to use AI-CDSSs (indirect effect:  $B = .20$ , 95% CI [.12, .34]). In the second mediation model with effort expectancy as mediator, the direct path from facilitating conditions to use intention did not reach statistical significance, while the indirect effect was positive ( $B = .21$ , 95% CI [.09, .37]). Accordingly, the relationship between facilitating conditions and use intention was fully mediated through effort expectancy.

**Table 6***Mediation models with performance and effort expectancy as mediators*

Path	<i>B</i>	95% CI
Mediator: Performance expectancy		
Direct effects		
Facilitating conditions → Performance expectancy	.38	NA, .54
Performance expectancy → Use intention	.53	.36, .70
Facilitating conditions → Use intention	.29	-.01, .57
Indirect effect		
Facilitating conditions → Performance expectancy → Use intention	.20	.12, .34
Mediator: Effort expectancy		
Direct effects		
Facilitating conditions → Effort expectancy	.48	.35, .62
Effort expectancy → Use intention	.43	.17, .68
Facilitating conditions → Use intention	.29	-.04, .61
Indirect effect		
Facilitating conditions → Effort expectancy → Use intention	.21	.09, .37

Note. *B* = regression estimate; CI = confidence interval.

### 4.3 Sensitivity Analysis

To assess the robustness of the meta-analytic findings, we employed cumulative meta-analysis. This approach involves conducting a sequence of iterative meta-analyses, with each analysis adding an additional effect size for a specific relationship. Effect sizes are added in the order of decreasing precision, meaning the initial effect sizes added represent the most accurate population effect size estimates. If less precise studies tend to skew the meta-analytic estimates, this will be observable as a shift in cumulative results when these studies are included [89]. The results of the cumulative meta-analyses are depicted in Figure S2 in the Online Appendix ([https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)). Five “drifts” were identified and all relationships drifted towards stronger effects as less precise studies were added, indicating an overestimation of the true effect. However meaningful differences were not observed after half the studies were added

compared to after all studies were added, for any of the relationships (Table S6 in the Online Appendix, [https://osf.io/djwkz/?view\\_only=7d6046aac9424a3c8f2facab08ac1e16](https://osf.io/djwkz/?view_only=7d6046aac9424a3c8f2facab08ac1e16)). Accordingly, we conclude that none of the drifts influenced the meta-analytic conclusions.

## 5. Discussion

### 5.1 Summary of Findings and Implications for Future Research

The primary goal of the meta-analysis was to gain a better understanding of the predictors of the intention to use AI-CDSSs among healthcare practitioners based on the UTAUT and its extensions. The results of the meta-analysis provide empirical support for the applicability of the UTAUT to the context of AI-CDSSs. As predicted, performance expectancy, effort expectancy, social influence, and facilitating conditions were positively related to the intention to use AI-CDSSs. These findings are largely in line with the findings of UTAUT meta-analyses in other fields [25,30,38,59]. We provide a summary of the main findings for our hypotheses and research questions in Tables 7 and 8.

**Table 7***The results of the investigation of hypotheses*

Hypothesis	<i>k</i>	Effect size [95% CI]	Result	Main findings
H1	16	$r_c = .66$ [.59, .73]	Supported	Performance expectancy is positively related to the intention to use AI-CDSSs.
H2	15	$r_c = .55$ [.43, .67]	Supported	Effort expectancy is positively related to the intention to use AI-CDSSs.
H3	15	$r_c = .66$ [.59, .72]	Supported	Social influence is positively related to the intention to use AI-CDSSs.
H4a	6	$r_c = .66$ [.42, .90]	Supported	Facilitating conditions are positively related to the intention to use AI-CDSSs.
H4b	<3	n/a	N/A	Not enough independent samples ( $k < 3$ ) to examine the relationship between facilitating conditions and actual use of AI-CDSSs.
H5a	4	$B = <.01$ [-.01, .01]	Not supported	The relationship of performance expectancy with the intention to use AI-CDSSs does not become weaker with increasing age.
H5b	4	$B = <.01$ [-.02, <.01]	Not supported	The relationship of effort expectancy with the intention to use AI-CDSSs does not become stronger with increasing age.
H5c	4	$B = -.01$ [-.01, <-.01]	Not supported	The relationship of social influence with the intention to use AI-CDSSs becomes weaker with increasing age.
H6a	16	$B = <.01$ [<-.01, <.01]	Not supported	The relationship of performance expectancy with the intention to use AI-CDSSs is not stronger for men.
H6b	15	$B = <.01$ [-.01, <.01]	Not supported	The relationship of effort expectancy with the intention to use AI-CDSSs is not stronger for women.
H6c	15	$B = <.01$ [<-.01, .01]	Not supported	The relationship of social influence with the intention to use AI-CDSSs is not stronger for women.
H7a-c	<3	n/a	N/A	Not enough independent samples ( $k < 3$ ) to examine experience as a moderator.
H8	<3	n/a	N/A	Not enough independent samples ( $k < 3$ ) to examine mandatory versus voluntary adoption setting as a moderator.

Note. *k* = number of independent samples; CI = confidence interval;  $r_c$  = sample size-weighted and reliability-corrected correlation; *B* = regression estimate.

**Table 8***The results of the investigation of research questions*

Research question	<i>k</i>	Effect size [95% CI]	Result	Main findings
RQ1	9	$r_c = .63$ [.52, .73]	Answered	Positive attitude towards AI-CDSSs is positively related to the intention to use AI-CDSSs.
RQ2	10	$r_c = .73$ [.63, .82]	Answered	Trust is positively related to the intention to use AI-CDSSs.
RQ3	10	$r_c = -.21$ [-.35, -.08]	Answered	Perceived risk is negatively related to the intention to use AI-CDSSs.
RQ4	3	$r_c = -.41$ [-.98, -.15]	Answered	AI anxiety is negatively related to the intention to use AI-CDSSs.
RQ5	5	$r_c = .54$ [.43, .64]	Answered	Personal innovativeness is positively related to the intention to use AI-CDSSs.
RQ6	3	$r_c = .85$ [.63, 1.00]	Answered	The intention to use AI-CDSSs is positively related to their actual use.
RQ7	N/A	See Table 3	Partially answered	See Table 3
RQ8a	<3	N/A	N/A	Not enough independent samples ( $k < 3$ ) to examine occupation as a moderator.
RQ8b	$k_1 = 4$ ; Diff. = -.31 $k_2 = 6$ [-.58, -.04]		Answered	The positive relationship of effort expectancy on use intention was weaker for diagnostic and treatment AI-CDSSs ( $k_1$ ) compared to diagnostic AI-CDSSs ( $k_2$ ).
RQ8c	13-14	See Table 4	Answered	Cultural background (individualism) does not moderate the relationship of performance expectancy ( $k = 14$ ), effort expectancy ( $k = 13$ ) and social influence ( $k = 13$ ) with the intention to use AI-CDSSs.
RQ9a	15-16	See Table 4	Answered	Publication year does not moderate the relationship of performance expectancy ( $k = 16$ ), effort expectancy ( $k = 15$ ) and social influence ( $k = 15$ ) with the intention to use AI-CDSSs.
RQ9b	6-10	See Table 5	Answered	There are no differences in the relationships of performance expectancy, effort expectancy and social influence with the intention to use AI-CDSSs between samples that used the Venkatesh scale to measure use intention ( $k_2$ ) versus those that used other scales ( $k_1$ ).
RQ10a	16	$B$ (indirect effect) = .20 [.12, .34]	Answered	Performance expectancy mediates the relationship between facilitating conditions and the intention to use AI-CDSSs.
RQ10b	16	$B$ (indirect effect) = .21 [.09, .37]	Answered	Effort expectancy mediates the relationship between facilitating conditions and the intention to use AI-CDSSs.

*Note.*  $k$  = number of independent samples; CI = confidence interval;  $r_c$  = sample size-weighted and reliability-corrected correlation;  $k_1$  and  $k_2$  = number of independent samples for moderator level 1 and 2; Diff. = Mean difference between sample size-weighted and reliability-corrected correlation for moderator level 1 and 2;  $B$  = regression estimate.

The results of relative weights analyses showed that all four UTAUT predictors together explain 50% of the variance in use intention among healthcare practitioners, reaffirming the relevance of the UTAUT predictors in the context of AI-CDSSs. Among the UTAUT predictors, performance expectancy emerged as the most relevant predictor, accounting for 31% of the total explained variance, followed by social influence (28%), facilitating conditions (26%), and effort expectancy (15%). In most UTAUT research, performance expectancy is more relevant than effort expectancy, possibly because performance expectancy is inherently connected to the primary motives behind technology use [31,39]. That is, it directly relates to the perceived benefits that users expect to gain from using a technology [31,39]. Effort expectancy refers to the expected ease of using a technology [22]. While important, the ease of use may become a secondary consideration if the technology does not meet the primary performance-related objectives. In other words, users might be willing to overcome a steeper learning curve if they believe the payoff in performance is worthwhile [62]. This could explain why performance expectancy accounts for a higher percentage of the variance in technology acceptance and use intentions compared to effort expectancy. Overall, the findings of the current meta-analysis reflect a common finding in technology acceptance research where the anticipated improvement in performance is often found to be a stronger driver of user acceptance than the anticipated effort to learn and use the technology [31,39,60,62].

Among the UTAUT predictors, effort expectancy and facilitating conditions had the widest credibility intervals (.56 and .66, respectively), suggesting the presence of moderating influences [84,95]. For example, previous research suggests that radiologists, accustomed to complex machines and heavy workloads, may be willing to invest effort into learning how to use new technology if it reduced their workload,

indicating a moderating influence of occupation on the relationship between effort expectancy and use intention [4,36]. In addition, the strength of the relationship between effort expectancy and use intention has been shown to differ between AI-CDSS for feedback versus decision support [24].

In addition to the core UTAUT variables, we identified attitude, trust, perceived risk, AI anxiety, and personal innovativeness as predictors of the intention to use AI-CDSSs. Although all included studies reported a negative relationship between AI anxiety and use intention, the confidence interval for AI anxiety included zero. This lack of an observed relationship may be due to the low sample sizes (total sample size was  $n = 391$ ) and the resulting high uncertainty in the true effect. Interestingly, in the relative weights analyses, trust proved to be a more relevant factor than performance expectancy in explaining variance in the intention to use AI-CDSSs. The relevance of trust may be explained by the lack of transparency in how AI recommendations are generated, coupled with the high stakes associated with clinical decision-making [96]. Indeed, research has suggested that even highly efficient AI-CDSSs may face resistance in clinical applications if healthcare practitioners do not trust in the system's safety [97–99]. The findings of the current meta-analysis align with research advocating for the inclusion of trust into the UTAUT model [97].

Furthermore, the current meta-analysis emphasizes the need to consider both drivers and inhibitors of the intention to use AI-CDSSs for a more comprehensive understanding of the adoption process [100]. The relative weights analyses demonstrate that AI anxiety explained about ten percent in the intention to use AI-CDSSs, after trust (29%), facilitating conditions (20%), performance expectancy (19%), social influence (14%), and before effort expectancy (7%). The relevance of perceived risk as a predictor of use intention was small (about 2% after all other

predictors). Risk perception is a cognitive assessment of the potential losses and gains from using AI-CDSSs, which is based on logical evaluation and can be mitigated by providing relevant information [101]. In contrast, AI anxiety is an emotional response that encompasses fears and insecurities about AI technology [102]. Accordingly, AI anxiety is less rational and more difficult to alleviate because it can be deeply rooted in concerns about AI's impact on job security, professional autonomy, and the quality of patient care [23,103,104].

Even for relationships with a substantial number of independent samples, such as performance expectancy, effort expectancy, attitude, trust and perceived risk, the credibility intervals are wide ( $> .34$ ), suggesting the presence of moderators [84,95]. This observation is supported by the modest amount of variance accounted for by statistical artifacts, indicating that there may be other reasons for substantial variance between individual studies [84,95]. While we considered multiple moderators suggested by the UTAUT and additional contextual and methodological moderators, we only found two moderation effects:

First, age moderated the relationship between social influence and use intention, with older healthcare practitioners experiencing a weaker relationship between social influence and use intention. This finding does not align with the UTAUT proposing that older individuals place more relevance on the opinion of relevant others when intending to use a new technology [22]. One explanation for this discrepancy may be that practitioners value their professional independence increasingly more with age, thus relying less on the opinion of others regarding technology use as they get older. While differences in professional values and behaviors have been shown to exist between younger and older healthcare practitioners [106], a systematic examination of the effect of age on the relationship between social influence and the intention to use AI-CDSSs is lacking. It has to be



noted that the observed moderation effect is based on only four independent samples, underscoring the need to systematically study the influence of age on the relationship between social influence and use intention.

Second, the relationship between effort expectancy and use intention was stronger for diagnostic AI-CDSSs compared to devices that combined diagnostic and treatment recommendations. When clinicians assess a tool solely for diagnostic purposes, they may find it easier to anticipate the required effort to use it, as the task is less complex and the outcome more direct. This clear understanding may strengthen the relationship between effort expectancy and use intention. Indeed, the perceived risk associated with smart devices has been found to negatively influence the relationship between effort expectancy and use intention [107]. The multifaceted nature of combined tools may make it more challenging for clinicians to evaluate the effort needed to understand and use them. This uncertainty possibly leads to a weaker relationship between effort expectancy and use intention, as clinicians may not be able to adequately assess the effort required, thus not being able to use it as a source of information when it comes to indicating their intention to use it. Future research is needed to investigate the moderating influence of the device type.

The results of the mediation analyses indicate that the relationship between facilitating conditions and use intention is fully explained through effort expectancy and partially explained through performance expectancy. This finding aligns with the UTAUT proposing that when performance and effort expectancy are considered, facilitating conditions lose their importance in predicting use intention [22]. One explanation for the relevance of effort expectancy as a mediator may be that issues related to the support infrastructure, a critical aspect of facilitating conditions, are also conceptually addressed by effort expectancy [22]. That is, if healthcare organizations establish the appropriate support infrastructure, the effort required to use AI-CDSSs

becomes lower [21,52]. Similarly, if a user perceives that the technology is supported by adequate facilitating conditions, they may be more likely to believe in the performance benefits when using the system, explaining the mediating role of performance expectancy.

## 5.2 Practical Implications

Performance expectancy and trust emerged as the two most relevant predictors of AI-CDSSs use intention, suggesting that measures targeted towards healthcare practitioners' beliefs in the performance and trustworthiness of AI-CDSSs may be effective in enhancing their intention to use them. However, the consistently positive link between performance expectancy and use intention also suggests that healthcare institutions need to take measures to deter the perception of low-performing systems as high-performing, which could potentially cause more harm than benefit [105]. Healthcare practitioners require transparent communication regarding the performance and limitations of AI-CDSSs, alongside adequate training to ensure their correct use. In addition, regulatory bodies like the FDA need to ensure that available AI-CDSSs meet certain safety and performance standards [66,106,107]. Adequate policies and oversight in these contexts may ensure a balance between the adoption and safe application of AI-CDSSs in healthcare decision-making.

Trust in technology is a multifaceted construct including users' perceptions of a system's functionality, helpfulness or benevolence, and integrity [46,47]. Consequently, actions taken to enhance performance expectancy may not be sufficient for building trust in a system [97]. If organizations aim to improve healthcare practitioners' trust in AI-CDSSs, they need to address the various facets relevant to trust in technology. This includes dealing with ethical issues related to data privacy and the potential misuse of AI-CDSS as well as addressing the lack of transparency

and explainability in AI-generated recommendations [108,109]. For example, trust has been associated with the system's capability to explain its decision-making process, emphasizing the role of explainable AI as a path to building trust in AI-CDSSs [97,110]. Additionally, regulatory strategies should be designed to promote and maintain trust in AI-CDSS, along with safe patient outcomes. This might include the use of post-market surveillance systems to monitor the performance of deployed AI-CDSSs over time, which has been suggested as a method for identifying and mitigating issues of utility and safety in real-world clinical settings [114,115]. Developers can integrate user-centered design principles to tailor AI-CDSSs to the needs and workflows of specific clinical specialties and roles. The early inclusion of user feedback may facilitate the development of user-friendly AI-CDSSs and increase trust in these systems [116,117]. Healthcare administrators may foster practitioners' trust by providing training programs to increase familiarity with the technology and by designing evaluation metrics that can monitor system performance and user satisfaction [118]

Social influence has been demonstrated to be a relevant predictor of healthcare practitioners' intentions to use AI-CDSSs, particularly among younger professionals. Institutions aiming to adopt AI-CDSSs can leverage the important role of social influence by establishing a culture that values technological advancements and by engaging key opinion leaders to advocate and exemplify the use of these systems. Additionally, trainings can be structured not only to educate but also to establish a shared understanding and a community of practice that positively reinforces the application of AI-CDSSs [111,112]. By addressing the social aspects of technology acceptance, healthcare institutions can ensure that their investment in AI is met with a user base that is both competent and willing to integrate these tools into their daily practice.

The importance of facilitating conditions underscores the need for healthcare organizations to provide a supportive infrastructure that simplifies the integration of AI-CDSSs into existing workflows. For instance, the provision of training programs, allowing healthcare practitioners to gain first-hand experience, and setting up accessible support teams ready to address system-related issues can considerably boost healthcare practitioners' intention to use such systems [22,113].

AI anxiety has emerged as a barrier to the intention to use AI-CDSSs in the relative weights analysis. Therefore, hospitals and other healthcare institutions should consider measures to counteract any irrational negative emotional reactions to AI before and during the integration of AI-CDSSs into clinical workflows. One potential method to mitigate AI anxiety involves increasing medical staff involvement in the development process [104] or providing more training opportunities to increase their exposure to AI-enabled devices, thus reducing irrational fears [111]

### **5.3 Limitations and Implications for Future Research**

The current meta-analysis is not without limitations. First, the present study offers insight into the predictors of use intention as the key determining factor of actual use. However, some healthcare practitioners may express intentions to use AI-CDSSs but are hesitant when it comes to their actual implementation. Few studies included in the meta-analysis examined the predictors of actual use, underscoring the need for additional research on predictors of the actual use of AI-CDSSs [4,46].

Second, we were unable to explain the considerable variation in some of the effects based on moderator analyses. We could not evaluate three UTAUT moderators—experience with AI-CDSSs, voluntariness of use, and occupation—owing to insufficient samples incorporating these variables. Additionally, although all studies including AI anxiety reported negative relationships with use intention, the confidence interval of the meta-analytic estimate included zero due to the low sample

size and the associated high uncertainty in the estimate. More studies on the relationship of AI anxiety with the intention to use AI-CDSSs are needed. The large credibility intervals and the low correlations between estimates and statistical artifacts suggest the existence of moderating factors not included in the meta-analysis [84,95]. Future research should explore moderating effects, such as differences in the observed relationships among healthcare practitioners working in different fields or roles, to better understand the boundary conditions that affect the relationships between predictors and the intention to use AI-CDSSs.

Third, the nine relevant predictors could not be examined in a single relative weight analysis. The use of multiple models with subsets of predictors is a pragmatic approach to addressing data sparsity. However, the selected approach hinders definitive conclusions regarding the importance of all considered predictors. Furthermore, innovativeness could not be assessed in the relative weights analysis due to a lack of available samples assessing this predictor. The compromises that had to be made in the relative weights analyses highlight the need for an updated meta-analysis that includes complete predictor sets.

Fourth, the insight derived from the meta-analysis are primarily confined to unspecific AI-CDSSs. Given that AI-CDSS adoption is still limited, only a handful of studies have delved into exploring predictors of specific AI-CDSSs with distinctive features [4,24,37]. The results of these studies show that the attitude towards AI-CDSSs may vary depending on use cases and system features. Future research should examine the adoption of individual systems and variations in effects across different types of AI-CDSSs.

Fifth, the existing body of research on AI-CDSS adoption primarily relies on cross-sectional observational studies, with questionnaires as the main method of data collection. These studies inherently limit the establishment of causal

relationships, thus underscoring the need for future research to include longitudinal or experimental designs for more robust evidence of causality. Longitudinal studies may also be used to shed light on the development of use intention and the relevance of relevant predictors over time. For example, it is possible that initial trust plays a crucial role during the implementation phase, but becomes less relevant once a system has been successfully implemented.

Sixth, we selected the UTAUT as a general theoretical framework to examine the predictors of the intention to use AI-CDSSs. However, there has been some criticism of the UTAUT [62,114]. For example, the UTAUT may not answer questions related to the determinants and processes involved in *value-adding* technology use [62,115,116]. We found support for the prediction that beliefs about the performance and ease of use of AI-CDSSs lead to a higher intention to use these systems. However, based on the UTAUT, it may not be resolved whether these beliefs are well-founded, i.e., whether positive expectations actually lead to beneficial use because the system is indeed high-performing and easily implementable. Another criticism pertains to the UTAUT's narrow viewpoint on individual use. Other models, such as the non-adoption, abandonment, scale-up, spread, and sustainability (NASSS) framework [117] adopt a system perspective. This approach enables the examination of predictors on micro (individual technology users), meso (organizational processes and systems), and macro (national policy and wider context) levels, thereby more accurately representing the complex processes involved in technology adoption [117,118]. Additionally, the UTAUT focuses on an individual's intention to use a technology and does not fundamentally consider how well the technology fits the task it is being utilized for. Theories such as the task-technology fit (TTF) model examine the interconnectedness between task and technological characteristics. The model delves into how features of both the assigned task and the technology at hand shape

the task-technology fit, thereby influencing the overall performance and use intentions [119].

Finally, some of the predictors included in the current meta-analysis, particularly the additional predictors beyond the UTAUT, may not be adequately represented using standard measurement instruments. For instance, AI anxiety has a multitude of dimensions, such as privacy violation anxiety, bias behavior anxiety, job replacement anxiety, learning anxiety, or ethics violation anxiety [120]. The current meta-analysis does not distinguish between these different aspects of AI anxiety as separate predictors of AI-CDSS use intentions. Similarly, trust in AI is a multifaceted construct that includes perceptions of the system's benevolence, competence, and integrity [46,47]. Moreover, trust may refer to different aspects of an AI-CDSS, such as trust in the reliability of its predictions when being applied to different contexts, trust in legal protection if harm to patients occurs from using the AI-CDSS, and trust in data privacy [3,45,121]. More research is needed that explores the relevance of different elements of trust (i.e., benevolence, competence, integrity) and elements of the AI-CDSS that may be trusted to different degrees (e.g., reliability, legal and liability issues, privacy concerns) for the intention to use AI-CDSSs in clinical decision-making [121].

## 6. Conclusion

The meta-analysis underscores the relevance of the UTAUT to examine the predictors of the intention to use AI-CDSSs in healthcare. The results indicate that performance expectancy, effort expectancy, social influence, and facilitating conditions are positively related to the intention to use AI-CDSSs among healthcare practitioners. The analyses further reveal the relevance of the additional predictors attitude, trust, personal innovativeness, AI anxiety, and perceived risk. The results of mediation analyses show that expectancy and performance expectancy explain the

relationship between facilitating conditions and use intention. Despite identifying age and AI-CDSSs type as moderating influences, there is scope for future research to investigate other possible moderators to explain the variability in the observed effects. While the UTAUT model provides a theoretical framework for studying healthcare practitioners' intention to use AI-CDSSs, it remains relatively silent on the predictors of value-adding use of AI-CDSSs. Future research could investigate the conditions that encourage value-adding usage by applying comprehensive frameworks that consider both individual and broader organizational processes (for example, clinic systems and administrative hurdles). Finally, the findings of this meta-analysis provide starting points for the development and integration of AI-CDSSs that are likely to be adopted by healthcare practitioners as end users.



**Conflicts of Interest**

The authors have no conflicts of interest to declare.



## Acknowledgements

The research was funded by the Volkswagen Foundation (Grant number: 98 525).



## References

1. Russell SJ, Norvig P. Artificial intelligence: a modern approach. Pearson; 2016. Available from: <https://thuvienso.hoasen.edu.vn/handle/123456789/8967> [accessed Apr 1, 2024] ISBN:978-1-292-15396-4
2. Henry K, Kornfield R, Sridharan A, Linton R, Groh C, Wang T, Wu A, Mutlu B, Saria S. Human-machine teaming is key to AI adoption: clinicians' experiences with a deployed machine learning system. *Npj Digit Med* 2022 Jul 21;5(97):1–6. doi: 10.1038/s41746-022-00597-7
3. Tran A, Nguyen L, Nguyen H, Nguyen C, Vu L, Zhang M, Vu T, Nguyen S, Tran B, Latkin C, Ho R, Ho C. Determinants of intention to use artificial intelligence-based diagnosis support system among prospective physicians. *Front Public Health* 2021 Nov 26;9:1–9. doi: 10.3389/fpubh.2021.755644
4. Zhai H, Yang X, Xue J, Lavender C, Ye T, Li J-B, Xu L, Lin L, Cao W, Sun Y. Radiation oncologists' perceptions of adopting an artificial intelligence-assisted contouring technology: Model development and questionnaire study. *J Med Internet Res* 2021 Sep 30;23(9):1–16. doi: 10.2196/27122
5. Berner ES, La Lande TJ. Overview of clinical decision support systems. *Clin Decis Support Syst Theory Pract* 2016;1–17. doi: 10.1007/978-3-319-31913-1\_1
6. Knop M, Weber S, Mueller M, Niehaves B. Human factors and technological characteristics influencing the interaction of medical professionals with artificial intelligence-enabled clinical decision support systems: Literature review. *JMIR Hum Factors* 2022 Mar 24;9(1):1–12. doi: 10.2196/28639
7. Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of clinical decision support systems: benefits, risks, and strategies for success. *Npj Digit Med* Nature Publishing Group; 2020;3(1):1–10. doi: 10.1038/s41746-020-0221-y
8. Juhn Y, Liu H. Artificial intelligence approaches using natural language processing to advance EHR-based clinical research. *J Allergy Clin Immunol* 2020;145(2):463–469. doi: 10.1016/j.jaci.2019.12.897
9. Ramgopal S, Sanchez-Pinto LN, Horvat CM, Carroll MS, Luo Y, Florin TA. Artificial intelligence-based clinical decision support in pediatrics. *Pediatr Res* 2023;93(2):Article 2. doi: 10.1038/s41390-022-02226-1
10. Harada T, Miyagami T, Kunitomo K, Shimizu T. Clinical decision support systems for diagnosis in primary care: A scoping review. *Int J Environ Res Public Health* 2021 Aug 10;18(16):1–14. doi: 10.3390/ijerph18168435
11. Musen MA, Middleton B, Greenes RA. Clinical Decision-Support Systems. In: Shortliffe EH, Cimino JJ, editors. *Biomed Inform Comput Appl Health Care* Biomed Cham: Springer International Publishing; 2021. p. 795–840. doi: 10.1007/978-3-030-58721-5\_24
12. Mucha H, Robert S, Breitschwerdt R, Fellmann M. Usability of clinical decision support systems. *Z Für Arbeitswissenschaft* 2023 Mar 1;77(1):92–101. doi:

10.1007/s41449-022-00324-8

13. Vasey B, Nagendran M, Campbell B, Clifton DA, Collins GS, Denaxas S, Denniston AK, Faes L, Geerts B, Ibrahim M, Liu X, Mateen BA, Mathur P, McCradden MD, Morgan L, Ordish J, Rogers C, Saria S, Ting DSW, Watkinson P, Weber W, Wheatstone P, McCulloch P. Reporting guideline for the early-stage clinical evaluation of decision support systems driven by artificial intelligence: DECIDE-AI. *Nat Med Nature Publishing Group*; 2022 May;28(5):924–933. doi: 10.1038/s41591-022-01772-9
14. Zhang R, Zhang Z, Wang D, Liu Z. Editorial: Responsible AI in healthcare: Opportunities, challenges, and best practices. *Front Comput Sci* 2023;5. doi: 10.3389/fcomp.2023.1265902
15. Vasey B, Ursprung S, Beddoe B, Taylor EH, Marlow N, Bilbro N, Watkinson P, McCulloch P. Association of clinician diagnostic performance with machine learning–based decision support systems: A systematic review. *JAMA Netw Open* 2021 Mar 11;4(3):1–15. doi: 10.1001/jamanetworkopen.2021.1276
16. Choi D-J, Park JJ, Ali T, Lee S. Artificial intelligence for the diagnosis of heart failure. *NPJ Digit Med* 2020;3(1):1–6. doi: 10.1038/s41746-020-0261-3
17. Duran J, Jongsma K. Who is afraid of black box algorithms? On the epistemological and ethical basis of trust in medical AI. *J Med Ethics* 2021;47(5):329–335. doi: 10.1136/medethics-2020-106820
18. Wang L, Chen X, Zhang L, Li L, Huang Y, Sun Y, Yuan X. Artificial intelligence in clinical decision support systems for oncology. *Int J Med Sci* 2023;20(1):79. doi: 10.7150/ijms.77205
19. Hummelsberger P, Koch TK, Rauh S, Dorn J, Lerner E, Raue M, Hudecek MFC, Schicho A, Colak E, Ghassemi M, Gaube S. Insights on the current state and future outlook of AI in health care: Expert interview study. *JMIR AI* 2023;2(1):1–17. doi: 10.2196/47353
20. Lorenzini G., Arbelaes Ossa L., Shaw D.M., Elger B.S. Artificial intelligence and the doctor-patient relationship expanding the paradigm of shared decision making. *Bioethics United Kingdom: NLM (Medline)*; 2023;((Lorenzini, Arbelaes Ossa, Shaw, Elger) Institute for Biomedical Ethics, University of Basel, Basel, Switzerland). doi: 10.1111/bioe.13158
21. Van Cauwenberge D., Van Biesen W., Decruyenaere J., Leune T., Sterckx S. “Many roads lead to Rome and the Artificial Intelligence only shows me one road”: an interview study on physician attitudes regarding the implementation of computerised clinical decision support systems. *BMC Med Ethics United Kingdom: NLM (Medline)*; 2022;23(1):50. doi: 10.1186/s12910-022-00787-8
22. Venkatesh V, Morris MG, Davis GB, Davis FD. User acceptance of information technology: Toward a unified view. *MIS Q* 2003;27(3):425–478. doi: 10.2307/30036540
23. Cheng M, Li X, Xu J. Promoting healthcare workers’ adoption intention of artificial-intelligence-assisted diagnosis and treatment: The chain mediation of social influence and human-computer trust. *Int J Environ Res Public Health* 2022;19(20):1–19. doi:

10.3390/ijerph192013311

24. Kleine A-K, Kokje E, Lerner E, Gaube S. Attitudes toward the adoption of 2 artificial intelligence-enabled mental health tools among prospective psychotherapists: Cross-sectional study. *JMIR Hum Factors* 2023;10:1–15. doi: <https://doi.org/10.2196/46859>
25. Tamori H, Yamashina H, Mukai M, Morii Y, Suzuki T, Ogasawara K. Acceptance of the use of artificial intelligence in medicine among japan's doctors and the public: A questionnaire survey. *JMIR Hum Factors* 2022 Mar 16;9(1):1–10. doi: 10.2196/24680
26. Dwivedi YK, Rana NP, Chen H, Williams MD. A meta-analysis of the unified theory of acceptance and use of technology (UTAUT). *Gov Sustain Inf Syst - Manag Transf Diffus IT Hamburg, Germany, September 22-24: Springer; 2011. p. 155–170. doi: 10.1007/978-3-642-24148-2\_10*
27. Fujimori R, Liu K, Soeno S, Naraba H, Ogura K, Hara K, Sonoo T, Ogura T, Nakamura K, Goto T. Acceptance, barriers, and facilitators to implementing artificial intelligence-based decision support systems in emergency departments: Quantitative and qualitative evaluation. *JMIR Form Res* 2022 Jun 13;6(6):1–12. doi: 10.2196/36501
28. Laka M, Milazzo A, Merlin T. Factors that impact the adoption of clinical decision support systems (CDSS) for antibiotic management. *Int J Environ Res Public Health* 2021 Jan;18(4):1–14. doi: 10.3390/ijerph18041901
29. Seliaman ME, Albahly MS. The reasons for physicians and pharmacists' acceptance of clinical support systems in Saudi Arabia. *Int J Environ Res Public Health Multidisciplinary Digital Publishing Institute; 2023 Jan;20(4):1–19. doi: 10.3390/ijerph20043132*
30. Dwivedi YK, Rana NP, Jeyaraj A, Clement M, Williams MD. Re-examining the unified theory of acceptance and use of technology (UTAUT): Towards a revised theoretical model. *Inf Syst Front* 2019;21:719–734. doi: 10.1007/s10796-017-9774-y
31. Khechine H, Lakhal S, Ndjambou P. A meta-analysis of the UTAUT model: Eleven years later. *Can J Adm Sci Can Sci Adm* 2016;33(2):138–152. doi: 10.1002/cjas.1381
32. Ajzen I. The theory of planned behavior. *Organ Behav Hum Decis Process* 1991;50(2):179–211. doi: 10.1016/0749-5978(91)90020-T
33. Davis FD. Perceived usefulness, perceived ease of use, and user acceptance of information technology. *MIS Q* 1989;13(3):319–340. doi: 10.2307/249008
34. Duan Y, Edwards JS, Dwivedi YK. Artificial intelligence for decision making in the era of Big Data—evolution, challenges and research agenda. *Int J Inf Manag* 2019;48:63–71. doi: 10.1016/j.ijinfomgt.2019.01.021
35. Fan W, Liu J, Zhu S, Pardalos P. Investigating the impacting factors for the healthcare professionals to adopt artificial intelligence-based medical diagnosis support system (AIMDSS). *Ann Oper Res* 2020 Nov;294(1–2):567–592. doi: 10.1007/s10479-018-2818-y
36. Prakash AV, Das S. Medical practitioner's adoption of intelligent clinical diagnostic decision support systems: A mixed-methods study. *Inf Manage* 2021;58(7). doi:

10.1016/j.im.2021.103524

37. Panigutti C, Beretta A, Giannotti F, Pedreschi D. Understanding the impact of explanations on advice-taking: A user study for AI-Based clinical decision support systems. *Proc 2022 CHI Conf Hum Factors Comput Syst New York, NY, USA: Association for Computing Machinery*; 2022. p. 1–9. doi: 10.1145/3491102.3502104
38. Akinnuwesi BA, Uzoka F-ME, Fashoto SG, Mbunge E, Odumabo A, Amusa OO, Okpeku M, Owolabi O. A modified UTAUT model for the acceptance and use of digital technology for tackling COVID-19. *Sustain Oper Comput* 2022 Jan 1;3:118–135. doi: 10.1016/j.susoc.2021.12.001
39. Dwivedi Y, Rana N, Tamilmani K, Raman R. A meta-analysis based modified unified theory of acceptance and use of technology (meta-UTAUT): A review of emerging literature. *Curr Opin Psychol* 2020 Apr 1;36:13–18. doi: 10.1016/j.copsyc.2020.03.008
40. Venkatesh V, Thong JYL, Xu X. Consumer acceptance and use of information technology: Extending the unified theory of acceptance and use of technology. *MIS Q Management Information Systems Research Center, University of Minnesota*; 2012;36(1):157–178. doi: 10.2307/41410412
41. Rudolph CW, Katz IM, Lavigne KN, Zacher H. Job crafting: A meta-analysis of relationships with individual differences, job characteristics, and work outcomes. *J Vocat Behav* 2017;102:112–138.
42. Kleine A-K, Rudolph CW, Zacher H. Thriving at work: A meta-analysis. *J Organ Behav* 2019;40(9–10):973–999. doi: 10.1002/job.2375
43. Fishbein M, Ajzen I. Belief, attitude, intention, and behavior: An introduction to theory and research. *Addison-Wesley Publication Company*; 1975.
44. Dalvi-Esfahani M, Mosharaf-Dehkordi M, Leong LW, Ramayah T, Jamal Kanaan-Jebna AM. Exploring the drivers of XAI-enhanced clinical decision support systems adoption: Insights from a stimulus-organism-response perspective. *Technol Forecast Soc Change* 2023 Oct 1;195. doi: 10.1016/j.techfore.2023.122768
45. Kohnke A, Cole ML, Bush R. Incorporating UTAUT predictors for understanding home care patients' and clinician's acceptance of healthcare telemedicine equipment. *J Technol Manag Innov* 2014;9(2):29–41. doi: 10.4067/S0718-27242014000200003
46. Ritter C. User-based barriers to the adoption of artificial intelligence in healthcare [PhD Thesis]. *Capella University*; 2019.
47. McKnight DH, Carter M, Thatcher JB, Clay PF. Trust in a specific technology: An investigation of its components and measures. *ACM Trans Manag Inf Syst* 2011 Jul 1;2(2):1–25. doi: 10.1145/1985347.1985353
48. McKnight DH. Trust in information technology. *Blackwell Encycl Manag* 2005. p. 329–331.
49. Thiebes S, Lins S, Sunyaev A. Trustworthy artificial intelligence. *Electron Mark* 2021 Jun 1;31(2):447–464. doi: 10.1007/s12525-020-00441-4

50. Cornelissen L, Egger C, van Beek V, Williamson L, Hommes D. The drivers of acceptance of artificial intelligence-powered care pathways among medical professionals: Web-based survey study. *JMIR Form Res* 2022;6(6):1–12. doi: 10.2196/33368
51. Bauer RA. Consumer behavior as risk taking. *Mark Crit Perspect Bus Manag* New York, NY: Routledge; 2001. p. 568.
52. Pan J, Ding S, Wu D, Yang S, Yang J. Exploring behavioural intentions toward smart healthcare services among medical practitioners: a technology transfer perspective. *Int J Prod Res* 2019 Sep 17;57(18):5801–5820. doi: 10.1080/00207543.2018.1550272
53. Calisto FM, Nunes N, Nascimento JC. Modeling adoption of intelligent agents in medical imaging. *Int J Hum-Comput Stud* 2022 Dec;168:1–15. doi: 10.1016/j.ijhcs.2022.102922
54. Johnson D, Verdicchio M. AI anxiety. *J Assoc Inf Sci Technol* 2017 Jun 22;68. doi: 10.1002/asi.23867
55. Kim J, Kadkol S, Solomon I, Yeh H, Soh J, Nguyen T, Choi J, Lee S, Srivatsa A, Nahass G, Ajilore O. AI anxiety: A comprehensive analysis of psychological factors and interventions. *SSRN Electron J* 2023 Sep 15;29. doi: 10.2139/ssrn.4573394
56. Agarwal R, Prasad J. A conceptual and operational definition of personal innovativeness in the domain of information technology. *Inf Syst Res INFORMS*; 1998 Jun;9(2):204–215. doi: 10.1287/isre.9.2.204
57. Ciftci O, Berezina K, Kang M. Effect of personal innovativeness on technology adoption in hospitality and tourism: Meta-analysis. In: Wörndl W, Koo C, Stienmetz JL, editors. *Inf Commun Technol Tour 2021 Cham: Springer International Publishing*; 2021. p. 162–174. doi: 10.1007/978-3-030-65785-7\_14
58. Lu J, Yao JE, Yu C-S. Personal innovativeness, social influences and adoption of wireless Internet services via mobile technology. *J Strateg Inf Syst* 2005 Sep 1;14(3):245–268. doi: 10.1016/j.jsis.2005.07.003
59. Smith H, Downer J, Ives J. Clinicians and AI use: where is the professional guidance? *J Med Ethics Institute of Medical Ethics*; 2023 Aug 22;1–5. PMID:37607805
60. Blut M, Chong A, Tsigna Z, Venkatesh V. Meta-analysis of the unified theory of acceptance and use of technology (UTAUT): Challenging its validity and charting a research agenda in the Red Ocean. *J Assoc Inf Syst* 2022 Jan 1;23:13–95. doi: 10.17705/1jais.00719
61. Holden RJ, Karsh B-T. The technology acceptance model: its past and its future in health care. *J Biomed Inform* 2010;43(1):159–172. doi: 10.1016/j.jbi.2009.07.002
62. Liu L, Miguel Cruz A, Rios Rincon A, Buttar V, Ranson Q, Goertzen D. What factors determine therapists' acceptance of new technologies for rehabilitation—a study using the Unified Theory of Acceptance and Use of Technology (UTAUT). *Disabil Rehabil* 2015;37(5):447–455. doi: 10.3109/09638288.2014.923529
63. Shachak A, Kuziemyky C, Petersen C. Beyond TAM and UTAUT: Future directions for

- HIT implementation research. *J Biomed Inform* 2019 Dec 1;100:1–5. doi: 10.1016/j.jbi.2019.103315
64. Huang K-Y, Choi N, Chengalur-Smith I. Cultural dimensions as moderators of the UTAUT model: A research proposal in a healthcare context. *Proc Sixt Am Conf Inf Syst Lima, Peru, August 12-15, 2010*; 2010. p. 1–10. Available from: <https://aisel.aisnet.org/amcis2010/188>
65. Lin H-C. An investigation of the effects of cultural differences on physicians' perceptions of information technology acceptance as they relate to knowledge management systems. *Comput Hum Behav* 2014 Sep;38:368–380. doi: 10.1016/j.chb.2014.05.001
66. Venkatesh V. Determinants of perceived ease of use: Integrating control, intrinsic motivation, and emotion into the technology acceptance model. *Inf Syst Res* 2000 Dec;11(4):342–365. doi: 10.1287/isre.11.4.342.11872
67. Benjamens S, Dhunoo P, Meskó B. The state of artificial intelligence-based FDA-approved medical devices and algorithms: an online database. *Npj Digit Med* 2020;3(1):118. doi: 10.1038/s41746-020-00324-0
68. Wang D, Wang L, Zhang Z, Wang D, Zhu H, Gao Y, Fan X, Tian F. “Brilliant AI doctor” in rural clinics: Challenges in AI-powered clinical decision support system deployment. *Proc 2021 CHI Conf Hum Factors Comput Syst New York, NY, USA: Association for Computing Machinery*; 2021. p. 1–18. doi: 10.1145/3411764.3445432
69. Pikkemaat M, Thulesius H, Nymberg V. Swedish primary care physicians' intentions to use telemedicine: a survey using a new questionnaire - physician attitudes and intentions to use telemedicine (PAIT). *Int J Gen Med* 2021;14:3445–3455. doi: 10.2147/IJGM.S319497
70. Alomari A, Soh B. Determinants of medical internet of things adoption in healthcare and the role of demographic factors incorporating modified UTAUT. *Int J Adv Comput Sci Appl Science and Information (SAI) Organization Limited*; 2023;14(7):17–31. doi: 10.14569/IJACSA.2023.0140703
71. Wichmann JL, Willemink MJ, De Cecco CN. Artificial intelligence and machine learning in radiology: Current state and considerations for routine clinical implementation. *Invest Radiol* 2020;55(9):619–627. doi: 10.1097/RLI.0000000000000673
72. Definition and list of health professionals. *Transform Scaling Health Prof Educ Train World Health Organ Guidel 2013 World Health Organization*; 2013. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK298950/> [accessed Jan 27, 2024]
73. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151(4):264–269. doi: 10.7326/0003-4819-151-4-200908180-00135
74. Richardson WS, Wilson MC, Nishikawa J, Hayward RS. The well-built clinical question: a key to evidence-based decisions. *ACP J Club* 1995;123(3). doi: 10.7326/ACPJC-1995-123-3-A12



75. Alhashmi SFS, Alshurideh M, Al Kurdi B, Salloum SA. A systematic review of the factors affecting the artificial intelligence implementation in the health care sector. In: Hassanien A-E, Azar AT, Gaber T, Oliva D, Tolba FM, editors. *Proc Int Conf Artif Intell Comput Vis AICV2020 Cham*: Springer International Publishing; 2020. p. 37–49. doi: 10.1007/978-3-030-44289-7\_4
76. AlQudah AA, Al-Emran M, Shaalan K. Technology acceptance in healthcare: A systematic review. *Appl Sci* 2021;11(22):1–40. doi: 10.3390/app112210537
77. Chen M, Zhang B, Cai Z, Seery S, Gonzalez MJ, Ali NM, Ren R, Qiao Y, Xue P, Jiang Y. Acceptance of clinical artificial intelligence among physicians and medical students: A systematic review with cross-sectional survey. *Front Med* 2022;9. doi: 10.3389/fmed.2022.990604
78. Chong AYL, Blut M, Zheng S. Factors influencing the acceptance of healthcare information technologies: A meta-analysis. *Inf Manage* 2022 Apr 1;59(3). doi: 10.1016/j.im.2022.103604
79. Santomartino SM, Yi PH. Systematic review of radiologist and medical student attitudes on the role and impact of AI in radiology. *Acad Radiol* 2022 Nov 1;29(11):1748–1756. doi: 10.1016/j.acra.2021.12.032
80. Jiang L, Lavaysse LM. Cognitive and affective job insecurity: A meta-analysis and a primary study. *J Manag SAGE Publications Inc*; 2018 Jul 1;44(6):2307–2342. doi: 10.1177/0149206318773853
81. Hoffman BJ, Woehr DJ. A quantitative review of the relationship between person–organization fit and behavioral outcomes. *J Vocat Behav* 2006 Jun 1;68(3):389–399. doi: 10.1016/j.jvb.2005.08.003
82. Posit team. RStudio: Integrated development environment for R. Boston, MA: Posit Software, PBC; 2023. Available from: <http://www.posit.co/>
83. Dahlke JA, Wiernik BM. psychmeta: An R package for psychometric meta-analysis. *Appl Psychol Meas* 2019;43(5):415–416. doi: 10.1177/0146621618795933
84. Cheung MW-L. metaSEM: An R package for meta-analysis using structural equation modeling. *Front Psychol* 2015;5:7. doi: 10.3389/fpsyg.2014.01521
85. Schmidt FL, Hunter JE. *Methods of meta-analysis: Correcting error and bias in research findings*. Third edition. Sage; 2015. Available from: <https://doi.org/10.4135/9781483398105>
86. Johnson JW. A heuristic method for estimating the relative weight of predictor variables in multiple regression. *Multivar Behav Res* 2000;35(1):1–19. doi: 10.1207/S15327906MBR3501\_1
87. Viswesvaran C, Ones DS. Theory testing: Combining psychometric meta-analysis and structural equations modeling. *Pers Psychol* 1995;48(4):865–885. doi: 10.1111/j.1744-6570.1995.tb01784.x
88. LeBreton JM, Hargis MB, Griepentrog B, Oswald FL, Ployhart RE. A multidimensional approach for evaluating variables in organizational research and

- practice. *Pers Psychol* 2007;60(2):475–498. doi: 10.1111/j.1744-6570.2007.00080.x
89. Tonidandel S, LeBreton JM. Relative importance analysis: A useful supplement to regression analysis. *J Bus Psychol* 2011;26:1–9. doi: 10.1007/s10869-010-9204-3
  90. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. *Introduction to meta-analysis*. 2nd ed. Hoboken, NJ: John Wiley & Sons; 2021.
  91. Hofstede G. *Culture's Consequences: International Differences in Work-Related Values*. SAGE; 1984. ISBN:978-0-8039-1306-6
  92. Minkov M, Kaasa A. Do dimensions of culture exist objectively? A validation of the revised Minkov-Hofstede model of culture with World Values Survey items and scores for 102 countries. *J Int Manag* 2022 Dec 1;28(4). doi: 10.1016/j.intman.2022.100971
  93. Jak S. *Meta-analytic structural equation modelling*. Springer; 2015.
  94. Cheung MW-L. *Meta-analysis: A structural equation modeling approach*. John Wiley & Sons; 2015.
  95. Cheung MW-L, Chan W. Meta-analytic structural equation modeling: a two-stage approach. *Psychol Methods* 2005;10(1):40–64. doi: 10.1037/1082-989X.10.1.40
  96. Whitener EM. Confusion of confidence intervals and credibility intervals in meta-analysis. *J Appl Psychol US: American Psychological Association*; 1990;75(3):315–321. doi: 10.1037/0021-9010.75.3.315
  97. Tucci V, Saary J, Doyle TE. Factors influencing trust in medical artificial intelligence for healthcare professionals: a narrative review. *J Med Artif Intell AME Publishing Company*; 2022 Mar 30;5(0):13. doi: 10.21037/jmai-21-25
  98. Berge GT, Granmo OC, Tveit TO, Munkvold BE, Ruthjersen AL, Sharma J. Machine learning-driven clinical decision support system for concept-based searching: a field trial in a Norwegian hospital. *BMC Med Inform Decis Mak* 2023 Jan 10;23(1):5. doi: 10.1186/s12911-023-02101-x
  99. Evans RP, Bryant LD, Russell G, Absolom K. Trust and acceptability of data-driven clinical recommendations in everyday practice: A scoping review. *Int J Med Inf* 2024 Mar 1;183. doi: 10.1016/j.ijmedinf.2024.105342
  100. Lambert SI, Madi M, Sopka S, Lenes A, Stange H, Buszello C-P, Stephan A. An integrative review on the acceptance of artificial intelligence among healthcare professionals in hospitals. *Npj Digit Med* 2023 Jun 10;6(1):1–14. doi: 10.1038/s41746-023-00852-5
  101. Philippi P, Baumeister H, Apolinário-Hagen J, Ebert DD, Hennemann S, Kott L, Lin J, Messner E-M, Terhorst Y. Acceptance towards digital health interventions – model validation and further development of the Unified Theory of Acceptance and Use of Technology. *Internet Interv* 2021 Dec 1;26:1–9. doi: 10.1016/j.invent.2021.100459
  102. Liu C-F, Chen Z-C, Kuo S-C, Lin T-C. Does AI explainability affect physicians' intention to use AI? *Int J Med Inf* 2022 Dec 1;168. doi: 10.1016/j.ijmedinf.2022.104884

103. Kaya F, Aydin F, Schepman A, Rodway P, Yetişensoy O, Demir Kaya M. The roles of personality traits, AI anxiety, and demographic factors in attitudes toward artificial intelligence. *Int J Human-Computer Interact* Taylor & Francis; 2024 Jan 17;40(2):497–514. doi: 10.1080/10447318.2022.2151730
104. Huo W, Zheng G, Yan J, Sun L, Han L. Interacting with medical artificial intelligence: Integrating self-responsibility attribution, human-computer trust, and personality. *Comput Hum Behav* 2022 Jul 1;132. doi: 10.1016/j.chb.2022.107253
105. Huo W, Yuan X, Li X, Luo W, Xie J, Shi B. Increasing acceptance of medical AI: The role of medical staff participation in AI development. *Int J Med Inf* 2023 Jul 1;175:105073. doi: 10.1016/j.ijmedinf.2023.105073
106. Foster TL, Bowie P. Professional values and behaviours of younger and older general practitioners in Scotland: a cross-sectional survey. *BMJ Lead BMJ Specialist Journals*; 2020 Jun 1;4(2):57–63. doi: 10.1136/leader-2019-000138
107. Jena RK. Exploring Antecedents of Peoples' Intentions to Use Smart Services in a Smart City Environment: An Extended UTAUT Model. *J Inf Syst* 2022 Mar 1;36(1):133–149. doi: 10.2308/ISYS-2020-050
108. Gaube S, Suresh H, Raue M, Merritt A, Berkowitz SJ, Lerner E, Coughlin JF, Guttig JV, Colak E, Ghassemi M. Do as AI say: susceptibility in deployment of clinical decision-aids. *Npj Digit Med* 2021 Feb 19;4(1):1–8. doi: 10.1038/s41746-021-00385-9
109. Kleine A-K, Lerner E, Cecil J, Heinrich A, Gaube S. Advancing mental health care with AI-enabled precision psychiatry tools: A patent review. *Comput Hum Behav Rep* 2023 Dec 1;12:1–11. doi: 10.1016/j.chbr.2023.100322
110. Muehlematter UJ, Daniore P, Vokinger KN. Approval of artificial intelligence and machine learning-based medical devices in the USA and Europe (2015–20): A comparative analysis. *Lancet Digit Health Elsevier B.V.*; 2021;3(3):195–203. doi: 10.1016/S2589-7500(20)30292-2
111. Hlávka JP. Chapter 10 - Security, privacy, and information-sharing aspects of healthcare artificial intelligence. In: Bohr A, Memarzadeh K, editors. *Artif Intell Healthc* Academic Press; 2020. p. 235–270. doi: 10.1016/B978-0-12-818438-7.00010-1
112. Lukyanenko R, Maass W, Storey VC. Trust in artificial intelligence: From a Foundational Trust Framework to emerging research opportunities. *Electron Mark* 2022 Dec 1;32(4):1993–2020. doi: 10.1007/s12525-022-00605-4
113. Yang W, Wei Y, Wei H, Chen Y, Huang G, Li X, Li R, Yao N, Wang X, Gu X, Amin MB, Kang B. Survey on explainable AI: From approaches, limitations and applications aspects. *Hum-Centric Intell Syst* 2023 Sep 1;3(3):161–188. doi: 10.1007/s44230-023-00038-y
114. Price WN, Cohen IG. Privacy in the age of medical big data. *Nat Med Nature Publishing Group*; 2019 Jan;25(1):37–43. doi: 10.1038/s41591-018-0272-7
115. Goodman KE, Morgan DJ, Hoffmann DE. Clinical Algorithms, Antidiscrimination Laws, and Medical Device Regulation. *JAMA* 2023 Jan 24;329(4):285–286. doi: 10.1001/jama.2022.23870

116. Neff MC, Schaaf J, Noll R, Holtz S, Schütze D, Köhler SM, Müller B, Ahmadi N, von Wagner M, Storf H. Initial User-Centred Design of an AI-Based Clinical Decision Support System for Primary Care. *Stud Health Technol Inform* 2024 Jan 25;310:1051–1055. PMID:38269975
117. Beltrão G, Paramonova I, Sousa S. User Interface Design for AI-Based Clinical Decision-Support System: Preliminary Study. 2022. p. 1–4. doi: 10.23919/CISTI54924.2022.9820378
118. Wang W, Wang Y, Chen L, Ma R, Zhang M. Justice at the Forefront: Cultivating felt accountability towards Artificial Intelligence among healthcare professionals. *Soc Sci Med* 2024 Apr 1;347:1–11. doi: 10.1016/j.socscimed.2024.116717
119. Kwak Y, Ahn J-W, Seo YH. Influence of AI ethics awareness, attitude, anxiety, and self-efficacy on nursing students' behavioral intentions. *BMC Nurs* 2022 Sep 30;21(1):267. doi: 10.1186/s12912-022-01048-0
120. Vallo Hult H, Hansson A, Gellerstedt M. Digitalization and physician learning: Individual practice, organizational context, and social norm. *J Contin Educ Health Prof* 2020;40(4):220–227. PMID:33284172
121. Thakkar B, Bharathi V. Medical specialists' perception about adoption of artificial intelligence in the healthcare sector. *Cardiometry* 2022 Dec;(25):426–434. doi: 10.18137/cardiometry.2022.25.426434
122. Bayaga A, du Plessis A. Ramifications of the unified theory of acceptance and use of technology (UTAUT) among developing countries' higher education staffs. *Educ Inf Technol* 2023 Sep 19; doi: 10.1007/s10639-023-12194-6
123. Novak LL, Anders S, Gadd CS, Lorenzi NM. Mediation of adoption and use: a key strategy for mitigating unintended consequences of health IT implementation. *J Am Med Inform Assoc* 2012 Nov 1;19(6):1043–1049. doi: 10.1136/amiajnl-2011-000575
124. Shachak A, Montgomery C, Dow R, Barnsley J, Tu K, Jadad AR, Lemieux-Charles L. End-user support for primary care electronic medical records: a qualitative case study of users' needs, expectations, and realities. *Health Syst* 2013 Nov 1;2(3):198–212. doi: 10.1057/hs.2013.6
125. Greenhalgh T, Wherton J, Papoutsis C, Lynch J, Hughes G, A'Court C, Hinder S, Fahy N, Procter R, Shaw S. Beyond adoption: A new framework for theorizing and evaluating nonadoption, abandonment, and challenges to the scale-up, spread, and sustainability of health and care technologies. *J Med Internet Res* 2017 Nov 1;19(11):1–21. doi: 10.2196/jmir.8775
126. Greenhalgh T, Wherton J, Papoutsis C, Lynch J, Hughes G, A'Court C, Hinder S, Procter R, Shaw S. Analysing the role of complexity in explaining the fortunes of technology programmes: empirical application of the NASSS framework. *BMC Med* 2018 May 14;16(1):66. doi: 10.1186/s12916-018-1050-6
127. Abdekhoda M, Dehnad A, Zarei J. Factors influencing adoption of e-learning in healthcare: integration of UTAUT and TTF model. *BMC Med Inform Decis Mak* 2022 Dec 9;22:327. PMID:36494800

128. Li J, Huang J-S. Dimensions of artificial intelligence anxiety based on the integrated fear acquisition theory. *Technol Soc* 2020 Nov 1;63. doi: 10.1016/j.techsoc.2020.101410
129. Jones C, Thornton J, Wyatt JC. Artificial intelligence and clinical decision support: clinicians' perspectives on trust, trustworthiness, and liability. *Med Law Rev* 2023 May 22;31(4):501–520. PMID:37218368