

# Measuring Engagement in Provider-guided Digital Health Interventions: Development of a Conceptual and Analytical Framework Using Nurse WRITE as an Exemplar

Yan Wang, Annette DeVito Dabbs, Teresa Hagan Thomas, Grace Campbell, Heidi Ann Scharf Donovan

Submitted to: JMIR Formative Research  
on: February 19, 2024

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

*Table of Contents*

**Original Manuscript..... 5**  
**Supplementary Files..... 20**  
    Figures ..... 21  
        Figure 1..... 22

# Measuring Engagement in Provider-guided Digital Health Interventions: Development of a Conceptual and Analytical Framework Using Nurse WRITE as an Exemplar

Yan Wang<sup>1</sup> PhD; Annette DeVito Dabbs<sup>2</sup> PhD; Teresa Hagan Thomas<sup>3</sup> PhD; Grace Campbell<sup>4</sup> PhD, MSW; Heidi Ann Scharf Donovan<sup>5</sup> PhD

<sup>1</sup>Department of Health & Community Systems, School of Nursing University of Pittsburgh Pittsburgh US

<sup>2</sup>Department of Acute & Tertiary Care, School of Nursing University of Pittsburgh Pittsburgh US

<sup>3</sup>Department of Health Promotion & Development, School of Nursing University of Pittsburgh Pittsburgh US

<sup>4</sup>School of Nursing Duquesne University Pittsburgh US

<sup>5</sup>Health & Community Systems, School of Nursing University of Pittsburgh Pittsburgh US

## Corresponding Author:

Yan Wang PhD

Department of Health & Community Systems, School of Nursing  
University of Pittsburgh  
3500 Victoria Street, Victoria Building  
Pittsburgh  
US

## Abstract

**Background:** Limited guidance exists for analyzing participant engagement in provider-guided digital health interventions (DHIs). System usage is commonly used, with acknowledged limitations in measuring socio-affective and cognitive aspects of engagement. Nurse WRITE, an online nurse-guided DHI for managing symptoms among women with recurrent ovarian cancer, offers an opportunity to develop a framework for assessing multi-dimensional engagement.

**Objective:** This study aims to develop a conceptual and analytic framework to measure socio-affective, cognitive, and behavioral engagement with provider-guided DHIs. We then illustrate the framework's ability to describe and categorize engagement using Nurse WRITE as an example.

**Methods:** A sample of 68 participants from Nurse WRITE who posted on the message boards were included. We adapted a prior framework for conceptualizing and operationalizing engagement across three dimensions and finalized a set of six distinct measures. Utilizing patients' posts, we created two socio-affective engagement measures—total count of socio-affective engagement classes (e.g., sharing personal experience) and total word count—and two cognitive engagement measures—total count of cognitive engagement classes (e.g., asking information-seeking questions) and average response rate. Additionally, we devised behavioral engagement measures using website data—the total count of symptom care plans and plan reviews. K-means clustering categorized the participants into distinct groups based on levels of engagement across three dimensions. Descriptive statistics and narratives were used to describe engagement in three dimensions.

**Results:** On average, participants displayed socio-affective engagement 34.7 times, writing 14,851 words. They showed cognitive engagement 19.4 times with a 78.3% response rate to nurses' inquiries. Participants also submitted an average of 1.6 symptom care plans and 0.7 plan reviews. Participants were clustered into high (n = 13), moderate (n = 17), and low engagers (n = 38) based on the six measures. High-engagers wrote a median of 36,956 words. They demonstrated socio-affective engagement approximately 81 times and cognitive engagement around 46 times, approximately six times that of the low engagers and twice that of the moderate engagers. High engagers had a median 91.74% response rate to the nurses' queries, whereas moderate engagers had 86.36%, and low engagers had 68.32%. High engagers completed a median of three symptom care plans and two reviews, while moderate engagers completed two plans and one review. Low-engagers completed a median of one plan and no reviews.

**Conclusions:** This study developed and reported an engagement framework to guide behavioral intervention scientists to understand and analyze participants' engagement with provider-guided DHIs. Significant variations in engagement levels across three dimensions highlight the importance of measuring engagement with provider-guided DHIs in socio-affective, cognitive,

and behavioral dimensions. Future studies should validate the framework with other DHIs, explore the influence of patient and provider factors on engagement, and investigate how engagement influences intervention efficacy.

(JMIR Preprints 19/02/2024:57529)

DOI: <https://doi.org/10.2196/preprints.57529>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#)

## Original Manuscript

# Measuring Engagement in Provider-guided Digital Health Interventions: Development of a Conceptual and Analytical Framework Using Nurse WRITE as an Exemplar

Yan Wang<sup>1</sup>, Annette DeVito Dabbs<sup>2</sup>, Teresa H Thomas<sup>3</sup>, Grace Campbell<sup>4</sup>, Heidi AS Donovan<sup>5</sup>

<sup>1</sup> PhD, RN

Department of Health and Community Systems, School of Nursing, University of Pittsburgh, 3500 Victoria Street

<sup>2</sup> PhD, RN, ACNS-BC, FAAN

Department of Acute and Tertiary, School of Nursing, University of Pittsburgh, 3500 Victoria Street

<sup>3</sup> PhD, RN

Department of Health Promotion and Development, School of Nursing, University of Pittsburgh, 3500 Victoria Street

<sup>4</sup> PhD, MSW, BSN, RN, CNL, CRRN, FARN

School of Nursing, Duquesne University, Fisher Hall 517, Pittsburgh, PA; School of Medicine, Department of Gynecology, Oncology, and Reproductive Sciences, University of Pittsburgh, 3550 Terrace Street, Pittsburgh, PA

<sup>5</sup> PhD, RN

Department of Health and Community Systems, School of Nursing, University of Pittsburgh, 3500 Victoria Street

## Abstract

**Background:** Limited guidance exists for analyzing participant engagement in provider-guided digital health interventions (DHIs). System usage is commonly assessed, with acknowledged limitations in measuring socio-affective and cognitive aspects of engagement. Nurse WRITE, an 8-week online nurse-guided DHI for managing symptoms among women with recurrent ovarian cancer, offers an opportunity to develop a framework for assessing multi-dimensional engagement.

**Objective:** This study aims to develop a conceptual and analytic framework to measure socio-affective, cognitive, and behavioral engagement with provider-guided DHIs. We then illustrate the framework's ability to describe and categorize engagement using Nurse WRITE as an example.

**Methods:** A sample of 68 participants from Nurse WRITE who posted on the message boards were included. We adapted a prior framework conceptualizing and operationalizing engagement across three dimensions and finalized a set of six distinct measures. Utilizing patients' posts, we created two socio-affective engagement measures – total count of socio-affective engagement classes (e.g., sharing personal experience) and total word count –, and two cognitive engagement measures – total count of cognitive engagement classes (e.g., asking information-seeking question) and average question completion percentage. Additionally, we devised behavioral engagement measures using website data – total count of symptom care plans and plan reviews. K-means clustering categorized the participants into distinct groups based on levels of engagement across three dimensions. Descriptive statistics and narratives were used to describe engagement in three dimensions.

**Results:** On average, participants displayed socio-affective engagement 34.7 times, writing 14,851 words. They showed cognitive engagement 19.4 times with an average of 78.3% completion to nurses' inquiries. Participants also submitted an average of 1.6 symptom care plans and 0.7 plan reviews. Participants were clustered into high ( $n = 13$ ), moderate ( $n = 17$ ), and low engagers ( $n = 38$ ) based on the six measures. High engagers wrote a median of 36,956 words. They demonstrated socio-affective engagement approximately 81 times and cognitive engagement around 46 times, approximately six times that of the low engagers and twice that of moderate engagers. High engagers had a median of 91.74% completion of the nurses' queries, whereas moderate engagers had 86.36%, and low engagers had 68.32%. High engagers completed a median of three symptom care plans and two reviews, while moderate engagers completed two plans and one review. Low engagers completed a median of one plan with no reviews.

**Conclusions:** This study developed and reported an engagement framework to guide behavioral intervention scientists in understanding and analyzing participants' engagement with provider-guided DHIs. Significant variations in engagement levels across three dimensions highlight the importance of measuring engagement with provider-guided DHIs in socio-affective, cognitive, and behavioral dimensions. Future studies should validate the framework with other DHIs, explore the influence of patient and provider factors on engagement, and investigate how engagement influences intervention efficacy.

**Keywords:** Engagement; Digital health intervention; Framework; Symptom management; eHealth; Gynecological cancer

## Introduction

Active participant engagement is pivotal to maximizing intervention benefits and enhancing health outcomes [1]. Therefore, measuring and evaluating engagement levels is essential for valid conclusions about intervention efficacy. In recent decades, digital technologies, including mHealth and eHealth, have been extensively integrated into interventions targeting symptom management [2,3] and behavior change promotion [4-6]. Studies indicate that cancer survivors' use of technology and engagement with digital health interventions (DHIs) tends to decline over time [7,8], possibly due to participant intervention fatigue, cognitive overload, habituation to frequent contact, and negative emotions [9]. While human support is often considered a potential remedy for this issue in provider-guided DHIs [10], previous research has not adequately accounted for the complexity of patient engagement with provider-guided DHIs.

Although there is no consensus on the definition of engagement, it is predominantly seen as a multidimensional construct [11], including socio-affective, cognitive, and behavioral dimensions. Extant literature heavily relies on assessing behavioral engagement through system usage patterns, such as website visits, app usage, or posting frequency [12-20], which has been criticized in recent

systematic reviews [21]. Research indicates that solely measuring system usage does not guarantee that participants are receiving the intended dosage (the theoretically or empirically determined amount of exposure) of an intervention or that their interactions with the intervention are meaningful and effective. [22,23]. Attempts have been made to gauge participants' cognitive and affective engagement through subjective measures such as self-report questionnaires [24], ecological momentary assessments (EMAs) [25], and qualitative methods like interviews and think-aloud activities conducted during or after the intervention [26]. However, these measures have notable limitations. Exit interviews, for instance, fail to capture participants' real-time experiences during the intervention and may be compromised by recall bias. This issue was highlighted in a previous study on cancer survivors' engagement with an app where participants struggled to recall app functions or usage details [26]. Complementing this line of research, many researchers emphasize the importance of employing multiple methods and integrating diverse datasets that encompass cognitive and socio-affective engagement alongside usage data to unravel the complexities inherent in the concept and measurement of engagement [21, 27]. Building on these studies, we sought to develop a comprehensive conceptual and analytic framework to describe and assess patient engagement with a provider-guided DHI based on a more conceptual understanding of dimensions of engagement, including socio-affectivity [28], cognition [28] and behavior.

The WRITE Symptoms Study (NRG Oncology's GOG-259; NR010735) was a three-arm online symptom management intervention designed for women with recurrent cancer [29]. The Nurse WRITE intervention arm was a nurse-guided DHI conducted on online asynchronous message boards. A study nurse guided participants using the Representational Approach to patient education [30, 31]. This disease-agnostic approach thoroughly assesses participants' symptom beliefs before offering personalized symptom management recommendations and individualized problem-solving support. The other intervention arm adopted a similar process, replacing nurse interaction with self-directed (SD) online modules for participants. Both Nurse and SD WRITE demonstrated efficacy in improving symptom control, as measured by the Symptom Representation Questionnaire [31] compared to the control arm. However, there was a notable high variation in patient completion of the intervention. Importantly, all interactions between the nurses and participants are documented verbatim on the message board, providing an excellent opportunity to examine the complexity of the engagement.

In a previous study, we constructed the "MedNgage Dataset" [28] that covers coded cognitive and socio-affective engagement from the library of nurse-participant posts on asynchronous message boards in Nurse WRITE. Drawing from the social presence model [32], cognitive science of grounding in communication [33], and linguistic discourse theories [34], we collaborated with linguists and used iterative coding to develop the socio-affective and cognitive engagement (SACe) conceptual framework to capture these dimensions of engagement. First, we specifically characterized socio-affective engagement in provider-guided DHIs as the participants' endeavors to establish emotional connections with providers during activities like communication and collaboration. Second, cognitive engagement is defined as the participant's coordination of intervention content and process with providers (e.g., patient understands the content provided and effectively follows the intervention activities, such as answering protocolized questions). In a complex provider-guided DHI like Nurse WRITE, building emotional connections (socio-affective engagement) and participants' understanding of the intervention (cognitive engagement) can empower patients to explore and adopt the strategies best suited to managing their symptoms. This is the third dimension, behavioral engagement, and by taking action, participants' comprehension of skills and processes is enhanced, further reinforcing their connection with the provider.

Expanding on this foundation, the current study has two primary objectives. First, it aims to finalize a conceptual and analytic framework, building on the SACe, that quantifies engagement levels with provider-guided DHIs across three dimensions: socio-affective, cognitive, and behavioral. Secondly, it applies the framework to measure and categorize patients' engagement patterns with the Nurse



Write intervention as an exemplar. This provides guidance and insights for future evaluations of engagement in provider-guided DHIs.

## Methods

### Overview

This study was an ancillary analysis of 68 patients with recurrent ovarian cancer from the Nurse WRITE arm of the WRITE Symptoms Study. The study included qualitative data from patients' asynchronous message boards and quantitative website tracking data.

### Ethical Considerations

The WRITE symptoms interventions were approved by the University of Pittsburgh Human Research Protection Office Institutional Review Board (IRB) and each of the 34 participating clinical sites (IRB number: PRO09090033), allowing for de-identified data sharing [29]. All activities and questionnaires were conducted using a secure, password-protected website developed at the University of Pittsburgh, ensuring accurate and Health Insurance Portability and Accountability Act (HIPAA)-compliant data collection. This secondary data analysis used the de-identified dataset, posing no risks for participants.

### Setting and Sample

The parent trial enrolled 497 women with recurrent ovarian cancer from various Gynecologic Oncology Group (GOG)/NRG Oncology-affiliated sites [29]. To be eligible for the parent study, participants had to be 18 years or older, diagnosed with recurrent or persistent ovarian, fallopian, or primary peritoneal cancer, and possess a GOG performance status of less than 3, indicating that they are ambulatory and capable of self-care but unable to perform work activities, with more than 50% of waking hours spent up and about [35, 36]. Additionally, they were required to experience at least three symptoms associated with cancer or its treatment (e.g., pain, fatigue, neuropathy) and be able to read and write in English. Details of the study design, protocol for Nurse WRITE (intervention group), and results have been described elsewhere [29]. Among the 166 participants randomized to Nurse WRITE, 141 women met the study criteria and posted at least once on the message board. For this study, we selected a convenience sample of the first 68 participants randomized to Nurse WRITE, representing 50% of the total number of message board posts from the pool of 141 participants. This selection aimed to ensure alignment with the sample in the previous "MedNgage" study [28] and to manage the qualitative analysis workload.

### Asynchronous Message Boards

Nurse WRITE was delivered by nurses highly trained to the protocol that delivered the intervention with high fidelity, offering information and support tailored to each participant's needs [37]. Through open-ended prompts and protocolized questions, participants engaged in dynamic interactions with nurse interventionists on a private asynchronous message board on the study website, including expressing symptom perceptions, reflecting, discussing coping efforts, and collaborating with nurses in developing personalized symptom care plans for three participant-identified target symptoms during the 8-week intervention period. To create a symptom care plan, participants needed to go through six out of seven key intervention elements (representational assessment; identifying and exploring gaps, errors, and confusions; creating conditions for conceptual change; introducing replacement information; summary; goal setting and planning). The 7<sup>th</sup> element is reviewing and revising the symptom care plan after two weeks. This involved assessing goal achievement, strategy implementation, and any encountered difficulties. Participants were then instructed to try the revised plan and review the results following the same approach as needed on their own.

## Data Collection and Analysis

### *Participant Data*

Patient demographic characteristics were assessed with the Center for Research in Chronic Disorders Socio-Demographic Survey [38] for age, education, race, marital status, and employment.

### *Aim 1: Finalize a conceptual and analytic framework with engagement measures*

In the previous study [28], We applied the SACe framework to identify linguistic signals and categorize engagement within patient-nurse communication on Nurse WRITE's asynchronous message boards. This analysis revealed eight different socio-affective engagement classes (e.g., expressing positive sentiment, sharing cancer-related experiences, showing interest in continued communication) – behaviors representing participants' efforts to build emotional connections with nurses. Additionally, we identified seven cognitive engagement classes (e.g., answering intervention questions, agreeing with the nurse's suggestions, asking information-seeking questions) – behaviors that reflected participants' collaborative content and process coordination with nurses through written communication during the intervention. Figure 1 illustrates how engagement is conceptualized and operationalized in socio-affective, cognitive, and behavioral dimensions. The diagram progresses from left to right, beginning with conceptualizing engagement into three dimensions, each with its accompanying definition. Subsequently, socio-affective engagement is operationalized into eight distinct classes of behaviors; then, cognitive engagement is similarly operationalized into seven classes, each with a corresponding description. The third dimension, behavioral engagement, is measured by two intervention milestones: symptom care plan creation and review. This process culminated in the development of six measures to assess engagement within these three dimensions quantitatively.




Figure 1. Conceptualizing and operationalizing engagement with provider-guided DHIs across socio-affective, cognitive, and behavioral dimensions.

### Socio-affective dimension

In our prior study [28], we identified eight classes of behaviors in the socio-affective dimension, representing participants' emotional connection efforts with the nurse. To quantify engagement in this dimension, we developed two measures. First, the total count of socio-affective engagement classes was computed by summing the number of instances of each class. Further descriptions of socio-affective engagement classes in participant posts can be found in Figure 1. Additionally, the total word count served as a proxy for overall participant effort in communication and building connections with nurses [39], constituting the second socio-affective engagement measure.

### Cognitive dimension

In our prior study [28], we also identified seven classes in the cognitive dimension, demonstrating participants' efforts in coordinating intervention content and process with nurses. Engagement was quantified by summing the number of instances of each class, yielding the first cognitive measure: the total count of cognitive engagement classes. Full descriptions of each cognitive engagement class can be found in Figure 1. To obtain the second cognitive engagement measure, in this study, two trained raters independently analyzed the patients' posts to determine the question completion rate—reflecting cognitive engagement—by calculating the percentage of nurses' intervention-related questions and requests addressed by each participant on message boards. For instance, if a nurse poses seven protocolized questions about symptom representation in her post (e.g., 'How severe is your symptom?' or 'What have you tried to manage the symptoms?'), and the patient responds to

three of these questions, their question completion rate would be 3 out of 7. This metric serves as a vital indicator of patient cognitive engagement, indicating their level of understanding and coordination of intervention activities with the nurse, regardless of participant verbosity. Inter-rater reliability of participant question completion percentage, assessed on 178 posts, yielded a Cohen Kappa score of 0.84.

### Behavioral dimension

Website data was extracted to determine behavioral engagement measures: total count of symptom care plans and total count of care plan reviews and revisions.

### *Aim 2: Apply the framework to categorize patients' engagement patterns*

We conducted a descriptive analysis of patient characteristics (e.g., age, education). Following that, we quantified participants' engagement with Nurse WRITE using the six measures across socio-affective, cognitive, and behavioral dimensions: total count of socio-affective engagement classes, total word count, total count of cognitive classes, average question completion percentage, total count of care plans and plan reviews. Next, we used k-means clustering to categorize participants into distinctive groups based on these six measures. K-means clustering is an unsupervised machine learning algorithm that aims to group data based on feature similarity, with the number of groups represented by K [40]. Additionally, we conducted a narrative analysis [41] of participants' posts to elucidate further engagement patterns within these distinctive groups, including understanding participants' experiences and motivations.

## Results

### Sample characteristics

The majority of participants included in this analysis were white (63 out of 68; 93%), married/cohabitating (51 out of 68; 75%), middle-aged (mean= 59.7, SD = 9.5, range= 24-83), and had a bachelor's degree (median = 16, IQR = 4, range = 11-22 years of formal education). More than half of the participants were not working (37 out of 68; 54%), including individuals who were disabled/unable to work, retired, and unemployed. On average, participants reported moderate symptom severity (5.33 out of 10) and symptom distress (2.16 out of 4). Over half of the sample (38 out of 68; 56%) had at least one comorbidity.

### **Aim 1: Finalize a conceptual and analytic framework with engagement measures**

On average, participants exhibited socio-affective engagement classes of behaviors, such as expressing positive sentiment and appreciation, 34.7 times (SD = 28.8). The participants also contributed substantially to the message boards, with an average word count of 14,851 (SD = 14,064). Participants demonstrated dimensions of cognitive engagement behaviors such as answering protocolized questions, seeking information through inquiries, and agreeing with the nurse's opinions, on average, 19.4 times (SD = 16.3). Furthermore, they maintained an average completion of 78.3% (SD = 14.9 %) when addressing nurses' questions or fulfilling intervention tasks. Regarding behavioral engagement, on average, participants completed 1.6 symptom care plans (SD = 0.2) and 0.7 plan reviews (SD = 0.9).

### **Aim 2: Apply the framework to categorize patients' engagement patterns**

K-means clustering analysis revealed the formation of three main groups characterized by their level of engagement in the three dimensions (i.e., total count of socio-affective engagement classes, total word count, total count of cognitive engagement classes, question completion percentage, total count of symptom care plans and care plan reviews completed). Given the non-normal distribution of each engagement measure, we used the median and IQR to describe the engagement levels within each

main group. Table 1 presents the median and IQR values for measures in the three engagement dimensions across each main group.

Table 1. Median and IQR of six engagement measures across three dimensions for high (n = 13), moderate (n = 17), and low engagers (n = 38).

Engagement measures	High engagers	Moderate engagers	Low engagers
<b>Socio-affective dimension, Mdn (IQR)</b>			
Total count of socio-affective classes	81 (19)	39 (11)	13.5 (16)
Total word count	36,956 (20,066)	18,034 (5,787)	6315.5 (4,722)
<b>Cognitive dimension, Mdn (IQR)</b>			
Total count of cognitive classes	46 (12)	25 (8)	7.5 (8.3)
Average question completion percentage	91.7 (11.6)	86.4 (16.4)	68.3 (19.5)
<b>Behavioral dimension, Mdn (IQR)</b>			
Total count of symptom care plans	3 (0)	2 (1)	1 (1)
Total count of plan reviews	2 (1)	1 (0)	0 (0)

Abbreviations: Mdn = median, IQR = interquartile range.

There were 13 high engagers, 17 moderate engagers, and 38 low engagers. Regarding socio-affective engagement classes and the total word count, high engagers showed approximately six times higher engagement than low engagers and twice that of moderate engagers. More specifically, high engagers established rapport by addressing nurses by name, expressing sincere appreciation, sharing personal experiences, displaying vulnerability, and expressing positive emotions towards the intervention, resulting in a high median total count of socio-affective engagement behaviors of 81 times (IQR = 19). In contrast, moderate (Mdn = 39, IQR = 11) and low engagers (Mdn = 13.5, IQR = 16) showed less interest in building personal connections with the nurse. Regarding the total word count, highly engaged participants wrote a median of 36,956 words (IQR = 20,066) to their nurse, compared to moderate engagers (Mdn = 18,034, IQR = 5,787) and low engagers (Mdn = 6,315.5, IQR = 4,722).

Cognitively, high engagers actively participated and demonstrated a strong commitment to learning symptom management techniques. They more consistently completed nurses' questions and requests (Mdn = 91.7%, IQR = 11.6%) compared to moderate engagers (Mdn = 86.4%, IQR = 16.4%) and low engagers (Mdn = 68.3%, IQR = 19.5%); sought information, acknowledged nurses' posts, agreed with suggestions, showing a median frequency of cognitive engagement of 46 (IQR = 12), which is 6.1 times more often than low engagers and 1.8 times that of moderate engagers. Some high engagers even independently initiated second and third symptom care plans, sharing effective strategies and leading the intervention without prompts from the nurse. For example, one participant shared with the nurse the significance of prayer (which was not formally integrated as a strategy within the intervention), emphasizing its role in symptom management and coping with cancer. Compared to high engagers, the moderate engagers primarily focused on answering part of the intervention protocolized questions and agreeing with the nurse without gaining the confidence to lead themselves through the intervention.

Regarding behavioral engagement, high engagers completed a median of three symptom care plans and two revisions. Some of them exceeded intervention targets by working on the fourth or fifth symptoms. In contrast, moderate engagers completed a median of two symptom care plans and one review, whereas low engagers only completed one symptom care plan and no reviews. Highly

engaged participants, who formed an early rapport with nurses, often shared their insight and understanding following learning intervention materials, such as symptom care guides and common concerns for women with gynecological cancer. One patient wrote, “It was the first time I actually felt normal. When going to the doc, these symptoms have never even come up. I felt like I belonged! And really wasn't off my rocker.” This newfound perspective motivated her to pursue improved symptom management actively.

## Discussion

### Principal Results

Significant variations existed among engagers in socio-affective, cognitive, and behavioral dimensions, providing insights into how a provider-guided DHI operates. High engagers demonstrated approximately sixfold higher total word counts and socio-affective and cognitive engagement behaviors than low engagers and twice as much as moderate engagers. A positive feedback loop may exist between participant engagement and behavior change and symptom management [1, 42, 43]. Enhanced emotional connections with nurses and increased written communication (socio-affective engagement) appear to contribute to a deeper understanding of intervention content and improved problem-solving skills (cognitive engagement). This synergy of socio-affective and cognitive engagement streamlines the development of highly personalized symptom management care plans (behavioral engagement) and the implementation of various strategies to adopt healthy behaviors and manage symptoms. Consequently, these factors motivate participants to engage further with the nurse and the intervention, as the literature suggests [42]. Prior reviews also underscored the significance of connecting engagement with a DHI to desired behavior change [1,21]. Using a combination of measures, we have taken the first steps in measuring this meaningful engagement within a provider-guided DHI. However, further research is warranted to investigate and validate the relationship between engagement across these three dimensions and patient symptom outcomes, including patient-perceived symptom control.

The observed variations in participants' word counts may be related to the Nurse WRITE intervention's writing-intensive nature and differences in education levels among the groups. Limited computer literacy and busy work schedules may also have constrained participants' ability and available time for message board contributions, aligning with prior research showing that education level, computer literacy, and employment can influence participants' engagement with DHIs [42, 44]. While significant differences in participants' average question completion percentages were found among high, moderate, and low engagers, these differences were not as substantial as observed in other cognitive and socio-affective engagement measures. This is likely because some low and moderate engagers answered all the questions in their posts but logged on to the message board less frequently (and therefore had fewer posts), which artificially inflated their average completion percentage.

Regarding behavioral engagement, variations in the number of plans and plan reviews may be related to participants' perception of symptom severity and burden. Those who perceive more severe symptoms may be more likely to seek assistance from nurse interventionists to manage symptoms. In contrast, those who perceive milder symptoms may have prioritized work and family responsibilities or considered the intervention unnecessary. Further research is necessary to investigate and validate the impact of patient factors such as education, employment, and symptom perception on engagement levels to facilitate intervention tailoring.

Applying the DHI engagement framework to Nurse WRITE also provided valuable insights regarding the optimal duration and timing of provider-guided DHI for symptom management among highly disease-burdened individuals. Although the intervention aimed to address three target symptoms, 50% of high engagers couldn't complete the intervention goal (three “target” care plan reviews), indicating a need for a more extended intervention period. Moderate and low engagers took

longer to respond than high engagers, suggesting their potential to achieve the intervention goals with more time. These findings resonate with prior research on web-based distress management programs for cardiovascular patients, emphasizing the importance of investigating intervention duration and timing [45]. Our findings suggest that future digital symptom management interventions for advanced cancer patients should extend beyond eight weeks to ensure maximum benefit.

## Limitations

This study adopted an exploratory, iterative approach to assessing patient engagement patterns owing to the limited convenient sample size of 68 participants. Qualitative analysis unveiled subtleties in engagement patterns that extend beyond the three main groups identified by K-means clustering. Certain participants exhibited high levels of socio-affective engagement but demonstrated lower to moderate levels of cognitive engagement. Their emphasis leaned towards expressing emotions rather than active involvement with the intervention content aimed at care plan development, albeit such occurrences were infrequent. Additionally, the duration of Nurse WRITE was limited to 8 weeks, potentially insufficient for participants to fully achieve intervention goals. Future investigations with larger sample sizes and extended intervention periods are imperative to elucidate and build upon these findings comprehensively.

An alternative analysis of participants' communication and connection-building efforts could involve examining word count per post as a socio-affective engagement measure. However, the need for supporting linguistic literature for this approach presents a limitation. Furthermore, implementing such a measure may be challenging due to the diverse posting styles observed among Nurse WRITE participants. Some individuals favor numerous shorter posts, while others consolidate their responses into longer messages, potentially affecting the accuracy of the analysis.

This paper identifies and explains engagement patterns, proposing hypotheses regarding potential engagement influencers (e.g., employment and symptom severity) for future investigation. Building upon these hypotheses, our future work will focus on elucidating group characteristics, examining engagement-influencing factors, and providing insights for tailoring interventions, such as adjusting intervention dosage.

## Comparison with Prior Work

Compared to traditional measures of cognitive and socio-affective engagement dimensions, such as exit interview interventions [26] or the number of postings [20], our engagement framework offers a more nuanced perspective. Socio-affective and cognitive engagement classes of behaviors provide real-time insights into participants' focus, interests, emotions, and actions on message boards, allowing us to quantify the intrinsic aspects of two-way communication. These measures, along with the other socio-affective (total word count) and cognitive (average question completion percentage), guide behavioral intervention scientists in assessing participants' efforts and the outcomes of two-way interactions in a complex provider-guided DHI.

Expanding on this research direction, it's crucial to advance tools (e.g., built upon MedNgage models [28]) to efficiently capture various engagement metrics in near real-time and provide suggestions for intervention tailoring. This advancement targets the reduction of labor-intensive retrospective qualitative analysis and addresses methodological challenges outlined in the recent review [21]. One potential strategy is to incorporate a large language model (like ChatGPT or locally accessible alternatives) into the analysis of qualitative transcripts while taking measures to ensure patient data confidentiality. Such an approach could include summarizing patient-provider conversations and leveraging the outputs of MedNgage models [28] to produce detailed, easily comprehensible qualitative reports for nurse interventionists.



## Conclusions

Expanding upon prior research, we have developed a comprehensive framework for behavioral intervention scientists to analyze patient engagement in provider-guided DHIs. Employing this framework on Nurse WRITE—a provider-guided DHI—we classified participants as high, moderate, and low engagers across the three dimensions of engagement (socio-affective, cognitive, and behavioral). This provides insights into the operational intricacies of a successful provider-guided DHI across various levels of participant engagement. Further research is essential to validate this framework with other provider-guided DHIs, explore the impact of patient factors (e.g., education, employment, and symptom perception) on engagement, and assess how engagement influences the efficacy of the intervention. In the context of DHIs for symptom management among advanced cancer survivors, extending the intervention period beyond eight weeks should be considered so that participants have more opportunities to engage and obtain the full benefits of the intervention.

## Acknowledgments

The RCT (WRITE Symptoms intervention) was supported by a grant from the National Institutes of Health, National Institute of Nursing Research NIH-NINR R01NR010735-NRG Oncology GOG-259 (H.S.D., PI) as well as NIH-NCI grants to NRG Oncology (U10CA180822), NRG Operations (U10CA180868), and UG1CA189867 (NCORP). This research was supported by the Eta Chapter Sigma Theta Tau International Honor Society of Nursing Research Award (Y.W.) and the University of Pittsburgh School of Nursing Margaret E. Wilkes Scholarship (Y.W.). We want to express our sincere gratitude to Hannah Bergquist for her valuable contributions to the qualitative analysis of the asynchronous message boards. Lastly, we extend our immense appreciation to all the women who participated in this trial.

## Conflicts of Interest

The authors have no conflict of interest to disclose.

## Abbreviations

DHI: digital health intervention

GOG/NRG Oncology: Gynecologic Oncology Group/NRG Oncology

RCT: randomized controlled trial

IQR: interquartile range

Mdn: median

## References

1. Yardley L, Spring BJ, Riper H, Morrison LG, Crane DH, Curtis K, et al. Understanding and promoting effective engagement with digital behavior change interventions. *Am J Prev Med*. 2016 Nov; 51(5): 833–842. <https://doi.org/10.1016/j.amepre.2016.06.015>.
2. Agboola S, Kamdar M, Flanagan C, Searl M, Traeger L, Kvedar J, Jethwani K. Pain management in cancer patients using a mobile app: study design of a randomized controlled trial. *JMIR Res Protoc*. 2014 Dec 12; 3(4):e76. <https://doi.org/10.2196/resprot.3957>.
3. Parker S, Prince A, Thomas L, Song H, Milosevic D, Harris MF, et al. Electronic, mobile and telehealth tools for vulnerable patients with chronic disease: a systematic review and realist synthesis. *BMJ Open*. 2018 Aug 29; 8(8):e019192. <https://doi.org/10.1136/bmjopen-2017-019192>.
4. Hong YA, Goldberg D, Ory MG, Towne SD Jr., Forjuoh SN, Kellstedt D, et al. Efficacy of a mobile-enabled web app (iCanFit) in promoting physical activity among older cancer survivors: A pilot study; *JMIR Cancer*. 2015 Jun; 1(1): e7.



- <https://doi.org/10.2196/cancer.4389>.
5. Sun L, Wang Y, Greene B, Xiao Q, Jiao C, Ji M, et al. Facilitators and barriers to using physical activity smartphone apps among Chinese patients with chronic diseases; *BMC Med Inform and Decis Mak*. 2017 Apr; 17(1): 44. <https://doi.org/10.1186/s12911-017-0446-0>
  6. Wang Y, Sun L, Xu Y, Xiao Q, Chang P, & Wu Y. Comparison and analysis of top-10 exercise Android apps in mainland China. *Stud Health Technol Inform*. 2015; 216:1024.
  7. Demark-Wahnefried W, Schmitz KH, Alfano CM, Bail JR, Goodwin PJ, Thomson CA, et al. Weight management and physical activity throughout the cancer care continuum. *CA Cancer J Clin*. 2018 Jan; 68(1): 64–89. <https://doi.org/10.3322/caac.21441>
  8. Viola A, Panigrahi G, Devine KA. Digital interventions for adolescent and young adult cancer survivors. *Curr Opin Support Palliat Care*. 2020 Mar;14(1):51-59. <https://doi.org/10.1097/SPC.0000000000000480>.
  9. Nahum-Shan I, Smith SN, Spring BJ, Collins LM, Witkiewitz K, Tewari A, et al. Just-in-Time Adaptive Interventions (JITAI) in mobile health: key components and design principles for ongoing health behavior support. *Ann Behav Med*. 2018 May; 52(6): 446–462. <https://doi.org/10.1007/s12160-016-9830-8>
  10. Renfrew ME, Morton DP, Morton JK, & Przybylko G. The Influence of Human Support on the Effectiveness of Digital Mental Health Promotion Interventions for the General Population. *Front Psychol*. 2021 Aug; 12: 716106. <https://doi.org/10.3389/fpsyg.2021.716106>
  11. Kelders SM, van Zyl LE, & Ludden GDS. The Concept and Components of Engagement in Different Domains Applied to eHealth: A Systematic Scoping Review. *Front Psychol*. 2020; 11: 926. <https://doi.org/10.3389/fpsyg.2020.00926>
  12. Donkin L, Christensen H, Naismith SL, Neal B, Hickie IB, & Glozier N. A systematic review of the impact of adherence on the effectiveness of e-therapies. *J Med Internet Res*. 2011 Aug; 13(3): e52. <https://doi.org/10.2196/jmir.1772>
  13. Couper MP, Alexander GL, Zhang N, Little RJ, Maddy N, Nowak MA, et al. Engagement and retention: measuring breadth and depth of participant use of an online intervention. *J Med Internet Res*. 2010 Nov; 12(4): e52. <https://doi.org/10.2196/jmir.1430>
  14. Schubart JR, Stuckey HL, Ganeshamoorthy A, & Sciamanna CN. Chronic health conditions and internet behavioral interventions: a review of factors to enhance user engagement. *Comput Inform Nurs*. 2011 Feb; 29(2): 81-92. <https://doi.org/10.1097/NCN.0b013e3182065eed>
  15. Arden-Close EJ, Smith E, Bradbury K, Morrison L, Dennison L, Michaelides D, et al. A visualization tool to analyse usage of web-based interventions: the example of positive online weight reduction (POWeR). *J Med Internet Res*. 2015 May; 2(1): e8. doi:10.2196/humanfactors.4310.
  16. Chen Z, Koh PW, Ritter PL, Lorig K, Bantum EO, & Saria S. Dissecting an online intervention for cancer survivors: four exploratory analyses of internet engagement and its effects on health status and health behaviors. *Health Educ Behav*. 2015 Feb; 42(1): 32-45. <https://doi.org/10.1177/1090198114550822>
  17. Davies C, Duncan MJ, Vandelanotte C, Hall S, Corry K & Hooker C. Exploring the feasibility of implementing a pedometer-based physical activity program in primary school settings: a case study of 10,000 steps. *Health Promot J Austr*. 2012 Aug; 23(2): 141-144. <https://doi.org/10.1071/he12141>
  18. Morrison C & Doherty G. Analyzing engagement in a web-based intervention platform through visualizing log-data. *J Med Internet Res*. 2014 Nov;16(11): e252. <https://doi.org/10.2196/jmir.3575>
  19. Poirier J & Cobb NK. Social influence as a driver of engagement in a web-based health intervention. *J Med Internet Res*. 2012 Feb; 14(1): e36. <https://doi.org/10.2196/jmir.1957>
  20. Han JY, Kim JH, Yoon HJ, Shim M, McTavish FM, Gustafson DH. Social and psychological

- determinants of levels of engagement with an online breast cancer support group: posters, lurkers, and nonusers. *J Health Commun.* 2012 Nov; 17(3): 356–371. <https://doi.org/10.1080/10810730.2011.585696>
21. Short CE, DeSmet A, Woods C, Williams SL, Maher C, Middelweerd A, Müller AM, Wark PA, Vandelanotte C, Poppe L, Hingle MD, Crutzen R. Measuring Engagement in eHealth and mHealth Behavior Change Interventions: Viewpoint of Methodologies. *J Med Internet Res.* 2018 Nov; 20(11): e292. <https://doi.org/10.2196/jmir.9397>
  22. Paz Castro R, Haug S, Filler A, Kowatsch T, Schaub MP. Engagement Within a Mobile Phone-Based Smoking Cessation Intervention for Adolescents and its Association With Participant Characteristics and Outcomes. *J Med Internet Res.* 2017 Nov; 19(11): e356. <https://doi.org/10.2196/jmir.7928>
  23. Gouveia R, Karapanos E, & Hassenzahl M. How do we engage with activity trackers? a longitudinal study of Habito. In Proceedings of the 2015 ACM International Joint Conference on Pervasive and Ubiquitous Computing (UbiComp '15); Association for Computing Machinery, New York, NY, USA, 1305–1316. <https://doi.org/10.1145/2750858.2804290>
  24. Lefebvre RC, Tada Y, Hilfiker SW, & Baur C. The Assessment of User Engagement with eHealth Content: The eHealth Engagement Scale. *J. of Comput. Mediat. Commun.* 2010; 15: 666-681. <https://doi.org/10.1111/j.1083-6101.2009.01514.x>
  25. Potts C, Bond R, Ryan A, Mulvenna M, McCauley C, Laird E, Goode D. Ecological Momentary Assessment Within a Digital Health Intervention for Reminiscence in Persons With Dementia and Caregivers: User Engagement Study. *JMIR Mhealth Uhealth.* 2020 Jul 6;8(7):e17120. <https://doi.org/10.2196/17120>.
  26. Crafoord MT, Fjell M, Sundberg K, Nilsson M, Langius-Eklöf A. Engagement in an Interactive App for Symptom Self-Management during Treatment in Patients With Breast or Prostate Cancer: Mixed Methods Study. *J Med Internet Res.* 2020 Aug; 22(8): e17058. <https://doi.org/10.2196/17058>
  27. Martey RM, Kensk K, Folkestad J, Feldman L, Gordis E, Shaw A, et al. Measuring Game Engagement: Multiple Methods and Construct Complexity. *Simulation & Gaming*; 2014 Nov; 45(4-5): 528-547. <https://doi.org/10.1177/1046878114553575>
  28. Wang Y, Donovan HS, Hassan S, & Alikhani M. MedNgage: A Dataset for Understanding Engagement in Patient-Nurse Conversations. In Findings of the Association for Computational Linguistics: ACL 2023; 2023; 4613–4630. <https://aclanthology.org/2023.findings-acl.282/>
  29. Donovan HS, Sereika SM, Wenzel LB, Edwards RP, Knapp JE, Hughes SH, et al. Effects of the WRITE Symptoms Interventions on Symptoms and Quality of Life Among Patients With Recurrent Ovarian Cancers: An NRG Oncology/GOG Study (GOG-0259). *J Clin Oncol.* 2022 May; 40(13): 1464-1473. <https://doi.org/10.1200/JCO.21.00656>
  30. Donovan HS, Ward SE, Song MK, Heidrich SM, Gunnarsdottir S, Phillips CM. An update on the representational approach to patient education. *J Nurs Scholarsh.* 2007;39(3):259-65. doi: 10.1111/j.1547-5069.2007.00178.x.
  31. Donovan HS, Ward S, Sherwood P, Serlin RC. Evaluation of the Symptom Representation Questionnaire (SRQ) for assessing cancer-related symptoms. *J Pain Symptom Manage.* 2008 Mar; 35(3): 242-257. <https://doi.org/10.1016/j.jpainsymman.2007.04.017>
  32. Swan K, Garrison DR & Rochardson JC. A Constructivist Approach to Online Learning: The Community of Inquiry Framework. In Information technology and constructivism in higher education: Progressive learning frameworks; 2009. p. 43-57. IGI Global. <https://doi.org/10.4018/978-1-60566-654-9.ch004>
  33. Clark HH & Brennan SE. Grounding in communication. In J. M. L. Lauren B. Resnick, and Stephanie D. Teasley (Ed.), Perspectives on socially shared cognition. American Psychological Association; 1991. p. 127- 149 <https://doi.org/10.1037/10096-006>

34. Asher N, Asher NM, Lascarides A, Bird S, Press, CU, Boguraev, B, et al. Logics of Conversation. Studies in Natural Language Processing. Cambridge University Press. 2003.
35. Oken MM, Creech RH, Tormey DC, Horton J, Davis TE, McFadden ET, Carbone PP. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol*. 1982, Dec; 5(6), 649-655. <https://www.ncbi.nlm.nih.gov/pubmed/7165009>
36. Rubin SC. Chemotherapy of Gynecologic Cancers: Society of Gynecologic Oncologists Handbook. Lippincott Williams & Wilkins; 2004.
37. Dumrongpakapakorn P. The development of a comprehensive fidelity monitoring plan for a web-based psycho-educational intervention: The WRITE symptoms study [Doctoral dissertation]. [Pittsburgh, PA]: University of Pittsburgh; 2011.
38. Sereika S, & Engberg S. Development of Standardized Sociodemographic and Co-Morbidity Questionnaires. In: The 17th International Nursing Research Congress Focusing on Evidence- Based Practice; 2016.
39. Tausczik YR, & Pennebaker JW. The psychological meaning of words: LIWC and computerized text analysis methods. *Journal of Language and Social Psychology*; 2009 Dec; 29(1): 24-54. <https://doi.org/10.1177/0261927X09351676>
40. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel, O, et al. Scikit-learn: Machine Learning in Python. *Journal of Machine Learning Research*; 2011 Oct; 12: 2825-2830. URL: <https://scikit-learn.org/stable/modules/generated/sklearn.cluster.KMeans.html> [accessed 2024-05-10]
41. Stephens, C, & Breheny, M. Narrative Analysis in Psychological Research: An Integrated Approach to Interpreting Stories. *Qualitative Research in Psychology*; 2013; 10(1), 14–27. <https://doi.org/10.1080/14780887.2011.586103>
42. Perski O, Blandford A, West R, Michie S. Conceptualising engagement with digital behaviour change interventions: a systematic review using principles from critical interpretive synthesis. *Transl Behav Med*. 2017 Jun; 7(2): 254–267. <https://doi.org/10.1007/s13142-016-0453>
43. Peeler A, Nelson K, Agrawalla V, Badawi S, Moore R & Li D. Living with multimorbidity: A qualitative exploration of shared experiences of patients, family caregivers, and healthcare professionals in managing symptoms in the United States. *J Adv Nurs*. 2024 Jan; 00:1–15. <https://doi.org/10.1111/jan.15998>
44. Vandelanotte C, Müller AM, Short CE, Hingle M, Nathan N, Williams SL, Lopez ML, Parekh S, Maher CA. Past, Present, and Future of eHealth and mHealth Research to Improve Physical Activity and Dietary Behaviors. *J Nutr Educ Behav*. 2016 Mar; 48(3): 219-228 e211. <https://doi.org/10.1016/j.jneb.2015.12.006>
45. Habibović M, Cuijpers P, Alings M, van der Voort P, Theuns D, Bouwels L, Herrman JP, Valk S, Pedersen S. Attrition and adherence in a WEB-Based Distress Management Program for Implantable Cardioverter defibrillator Patients (WEBCARE): randomized controlled trial. *J Med Internet Res*. 2014 Feb; 16(2): e52. <https://doi.org/10.2196/jmir.2809>

## Supplementary Files

## Figures

Conceptualizing and operationalizing engagement with provider-guided DHIs across socio-affective, cognitive, and behavioral dimensions.

Construct	Dimension and its definition	Class and its description	Measure
Engagement	Socio-affective: participant's emotional connection with interventionists during activities, like communication and collaboration.	Positive sentiment: participant expresses positive emotions/feedback on the intervention and symptom management progress. Negative sentiment: participant expresses negative emotions/feedback on the intervention and symptom management progress. Appreciation: participant appreciates and recognizes the nurse's contribution and support. Cancer-related experience: participant shows their vulnerability (e.g., fear, guilt); discusses with nurse about the cancer-related issues that she finds hard to share with others. Personal experience: participant shares personal information with the nurse (e.g., family, friends, pets, health care team, religion). Interest in communication: participant invites nurse to ask for more information. Vocatives: participant addresses the nurse by the name. Greetings: participant starts posts with a greeting; signs off the email; shows the benevolence and kindness to the nurse.	Total count of socio-affective engagement classes  Total word count in posts
	Cognitive: participant's coordination of content and process with interventionists.	Intervention information: participant responds to the nurse's questions or requests to complete intervention activities, such as protocolized questions. Information-seeking question: participant asks questions to seeking information or suggestions. Clarification question: participant asks nurse questions seeking clarification about something the nurse said. The goal is to reduce misunderstanding. Agreement: participant agrees with the nurse's opinion or suggestion; agrees to the plans or do what the nurse suggests doing. Disagreement: participant disagrees with the nurse's opinion or suggestion; disagrees to the plans or do what the nurse suggests doing. Acknowledgment: participant acknowledges/affirms the nurse suggestions, ideas, summary, questions. Initiative taking: participant anticipates and takes initiative in symptom management; provides suggestions for future study design.	Total count of cognitive engagement classes  Average question completion percentage
	Behavioral: participant's completion of intervention tasks.		Total count of symptom care plans Total count of plans reviewed