

# **Investigation of Reliability and Sensitivity of a Virtual Assessment Developed for Workplace Concussions: Protocol for a Method Comparison Study**

Keely Barnes, Heidi Sveistrup, Mark Bayley, Michel Rathbone, Monica Taljaard, Mary Egan, Martin Bilodeau, Motahareh Karimijashni, Shawn Marshall

Submitted to: JMIR Research Protocols  
on: March 04, 2024

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 26

    Figures ..... 27

        Figure 1..... 28

    Multimedia Appendixes ..... 29

        Multimedia Appendix 1..... 30

# Investigation of Reliability and Sensitivity of a Virtual Assessment Developed for Workplace Concussions: Protocol for a Method Comparison Study

Keely Barnes<sup>1, 2, 3</sup> MHK; Heidi Sveistrup<sup>1, 2, 4, 5</sup> PhD; Mark Bayley<sup>6, 7</sup> MD; Michel Rathbone<sup>8</sup> MD, PhD; Monica Taljaard<sup>3, 9</sup> PhD; Mary Egan<sup>1, 2</sup> PhD; Martin Bilodeau<sup>1, 2, 4</sup> PhD; Motahareh Karimijashni<sup>1, 3</sup> MSc; Shawn Marshall<sup>2, 3, 10</sup> MD, MSc

<sup>1</sup>University of Ottawa, Faculty of Health Sciences, School of Rehabilitation Sciences Ottawa CA

<sup>2</sup>Bruyère Research Institute Ottawa CA

<sup>3</sup>Clinical Epidemiology Program, Ottawa Hospital Research Institute Ottawa CA

<sup>4</sup>University of Ottawa, Faculty of Health Sciences, School of Human Kinetics Ottawa CA

<sup>5</sup>Carleton University, Systems and Computer Engineering Technology Ottawa CA

<sup>6</sup>Kite Research Institute, Toronto Rehabilitation Institute, University Health Network Toronto CA

<sup>7</sup>Division of Physical Medicine and Rehabilitation, Temerty Faculty of Medicine, University of Toronto Toronto CA

<sup>8</sup>McMaster University, Faculty of Health Sciences, Department of Medicine, Division of Neurology Hamilton CA

<sup>9</sup>School of Epidemiology and Public Health, University of Ottawa Ottawa CA

<sup>10</sup>Department of Medicine, University of Ottawa Ottawa CA

## Corresponding Author:

Keely Barnes MHK

University of Ottawa, Faculty of Health Sciences, School of Rehabilitation Sciences

200 Lees Ave.

Ottawa

CA

## Abstract

**Background:** Workplace mild traumatic brain injuries are frequently associated with persistent symptoms leading to a reduction in productivity at work or even disability. People who sustain workplace injuries frequently need rehabilitation and support and the challenges of delivering these services was heightened during the COVID-19 pandemic as injured workers had to be cared for remotely. Currently, clinicians are conducting both in-person and virtual (remote) concussion assessments, however, the measures that are being used to complete these assessments have undocumented psychometric properties. This study will document psychometric properties of the clinical measures that are being used remotely and their ability to produce similar results to in-person assessments.

**Objective:** Through this method comparison study, we aim to: 1) evaluate sensitivity of the measures included in a virtual assessment toolkit when compared to an in-person assessment and 2) determine the inter-rater and intra-rater reliability of the measures included in a virtual assessment toolkit.

**Methods:** Patient-participants (people living with acquired brain injuries) will attend two assessments (in-person and virtual) at the Ottawa Hospital. The two assessments will be identical consisting of the measures included in our virtual concussion assessment toolkit that was previously developed, which includes finger-to-nose testing, the Vestibular/Ocular Motor Screening tool, balance testing, cervical spine range of motion, saccades testing, and evaluation of effort. All virtual assessments will occur using the Microsoft Teams platform and will be audio-video recorded. The clinician-assessor and patient-participant will complete a feedback form following completion of the assessments. A different clinician will also document the findings on observed videos of the virtual assessment shortly after completion of both in person and virtual assessments and approximately one month later. Inter-rater reliability will be assessed by comparing the second clinician's observation with the first clinician's initial virtual assessment. Intra-rater reliability will be evaluated by comparing second clinician's observation with their own assessment, conducted approximately one month later. Sensitivity will be documented by comparing the findings (identification of abnormality) of the in-person assessment completed by the initial clinician-assessor with the findings of the second clinician-assessor on the observation of the recording of the virtual assessment.

**Results:** The study recruitment is expected to be completed by September 2024.

**Conclusions:** Currently, it is unknown if completing concussion assessments virtually produce similar results to the in-person assessment. This work will act as a first step to determining the similarity of the virtual assessment to the matching in-person assessment and will provide information on the reliability of the virtual assessment.

(JMIR Preprints 04/03/2024:57663)

DOI: <https://doi.org/10.2196/preprints.57663>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.  
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org>

## Original Manuscript

## Investigation of Reliability and Sensitivity of a Virtual Assessment Developed for Workplace Concussions: Protocol for a Method Comparison Study

Keely Barnes MHK, PhD (c)<sup>1,2,3</sup>; Heidi Sveistrup PhD<sup>1,2,4,5</sup>; Mark Bayley MD<sup>6,7</sup>; Michel Rathbone MD, PhD<sup>8</sup>; Monica Taljaard PhD<sup>3,9</sup>; Mary Egan PhD<sup>1,2</sup>; Martin Bilodeau PhD<sup>1,2,4</sup>; Motahareh Karimijashni MSc, PhD (c)<sup>1,3</sup>; Shawn Marshall MD, MSc<sup>2,3,10</sup>

1. University of Ottawa, Faculty of Health Sciences, School of Rehabilitation Sciences, Ottawa, Canada
2. Bruyère Research Institute, Ottawa, Canada
3. Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, Canada
4. University of Ottawa, Faculty of Health Sciences, School of Human Kinetics, Ottawa, Canada
5. Carleton University, Systems and Computer Engineering Technology, Ottawa, Canada
6. Kite Research Institute, Toronto Rehabilitation Institute, University Health Network, Toronto, Canada
7. Division of Physical Medicine and Rehabilitation, Temerty Faculty of Medicine, University of Toronto, Toronto, Canada
8. McMaster University, Faculty of Health Sciences, Department of Medicine, Division of Neurology, Hamilton, Canada
9. School of Epidemiology and Public Health, University of Ottawa, Ottawa, Canada
10. Department of Medicine, University of Ottawa, Ottawa, Canada

### Corresponding Author

Keely Barnes  
University of Ottawa  
200 Lees Ave.  
Ottawa, Canada  
K1N 6N5  
Kbarn076@uottawa.ca

Journal: JMIR Research Protocols

## Abstract

**Background:** Workplace concussions are frequently associated with persistent symptoms leading to a reduction in productivity at work or even disability. People who sustain workplace injuries

frequently need rehabilitation and support and the challenges of delivering these services was heightened during the COVID-19 pandemic as injured workers had to be cared for remotely. Currently, clinicians are conducting both in-person and virtual (remote) concussion assessments, however, the measures that are being used to complete these assessments have undocumented psychometric properties. This study will document psychometric properties of the clinical measures that are being used remotely and their ability to produce similar results to in-person assessments.

**Objectives:** Through this method comparison study, we aim to: 1) evaluate sensitivity of the measures included in a virtual assessment toolkit when compared to an in-person assessment and 2) determine the inter-rater and intra-rater reliability of the measures included in a virtual assessment toolkit.

**Methods:** Sixty patient-participants (people living with acquired brain injuries (ABIs)) will be recruited from and attend two assessments (in-person and virtual) at the Ottawa Hospital located in Ontario, Canada. The two assessments will be identical consisting of the measures included in our virtual concussion assessment toolkit that was previously developed, which includes finger-to-nose testing, the Vestibular/Ocular Motor Screening tool, balance testing, cervical spine range of motion, saccades testing, and evaluation of effort. All virtual assessments will occur using the Microsoft Teams platform and will be audio-video recorded. The clinician-assessor and patient-participant will complete a feedback form following completion of the assessments. A different clinician will also document the findings on observed videos of the virtual assessment shortly after completion of both in person and virtual assessments and approximately one month later. Inter-rater reliability (Cohen's Kappa) will be assessed by comparing the second clinician's observation with the first clinician's initial virtual assessment. Intra-rater reliability (Cohen's Kappa) will be evaluated by comparing second clinician's observation with their own assessment, conducted approximately one month later. Sensitivity will be documented by comparing the findings (identification of abnormality) of the in-person assessment completed by the initial clinician-assessor with the findings of the second clinician-assessor on the observation of the recording of the virtual assessment.

**Results:** As of May 2024, we have recruited seven clinician-assessors and completed study assessments with 39 patient-participants. The study recruitment is expected to be completed by September 2024.

**Conclusion:** Currently, it is unknown if completing concussion assessments virtually produce similar results to the in-person assessment. This work will act as a first step to determining the similarity of the virtual assessment to the matching in-person assessment and will provide information on the reliability of the virtual assessment.

**Keywords:** telehealth; virtual care; concussion; mTBI; assessment; examination

## Introduction

A direct or indirect head impact leading to a concussion, a form of mild traumatic brain injury (mTBI) [1], results in an alteration of brain functioning [2,3]. Most individuals who sustain a

concussion experience rapid recovery within the first two weeks to one month following the injury. There are, however, instances, where individuals continue to experience persistent symptoms after a concussion, with a reported occurrence of approximately 15-20% [4]. These symptoms may persist beyond one year following the injury [3,5]. Factors such as pre-morbid characteristics (age, sex, concussion history), psychological, and injury-related (severity of injury) factors are likely contributors to the persistence of the symptoms [6]. Post-concussion symptom presentation can vary widely and could include cognitive deficits, such as memory issues, vestibular deficits, such as balance issues or dizziness, ocular deficits, such as blurred vision, among others [1,7–9]. There is a need for a multifactorial assessment approach due to the diversity in symptom presentation. Specifically, it is recommended to incorporate measures targeting each of the commonly experienced symptoms into a battery of tests [10].

In the workplace context, there is evidence suggesting a growing documented incidence of concussion [11]. Kristman et al. reported that there are approximately 39-58 workplace concussions per 10,000 claims annually in Ontario, Canada [12]. Workplace concussions often lead to greater delays in return to work, compared to injuries sustained outside of the workplace context [5,13]. It is estimated that roughly 60 to 90% of adults return to work within 6 months post-concussion injury, with the remainder continuing to be off work at this timepoint [5]. Compensation associated with a workplace injury may be a contributor to this delay in return to work [5,13].

Several factors can contribute to delays in return to work following concussion including lower level of education, more severe immediate symptoms post-injury, and other injuries sustained [5]. In general, workplace concussions pose significant challenges. These include financial burdens that are felt by the healthcare system, employers, the Workplace Safety and Insurance Board (WSIB), and the workers themselves [14]. Healthcare costs associated with Ontario, Canada WSIB claims related to the diagnostic codes of concussion and intracranial injury averaged \$2,234.81 per claim and non-healthcare claims averaged \$4,676.38 per claim [14]. For the patient, lives are disrupted due



to prolonged recovery times [13] resulting in the need for additional support from the WSIB for individuals presenting for specialized assessments.

The onset of the COVID-19 pandemic affected how specialized assessments post-concussion could be completed and created a need to switch to virtual care [15]. However, it is not known whether the measures being used by clinicians in virtual assessments are the same to those used for in-person assessments [16]. Additionally, the reliability of the virtual concussion assessments being conducted is unknown. Measures used in-person that have shown strong psychometric properties may not produce the same or similar results when administered virtually [17]. Furthermore, the agreement between measures used in the in-person concussion assessment compared to the same measures used in the virtual assessment has not yet been established. Currently, clinicians are continuing to complete both in-person and virtual concussion assessments, however, there is limited information on the measures used in the virtual assessment. It is important to gain an understanding of the properties of the clinical measures and their ability to produce similar results to the in-person assessment to inform effectiveness of use in a virtual context [18].

The preparatory work for this study was initiated by developing a virtual assessment toolkit. Clinician surveys were used to identify relevant measures for assessing the physical domains of concussion [19]. A working group refined the measures feasible for virtual administration based on expert opinion. Focus groups then identified barriers and facilitators for virtual concussion assessments. The resulting virtual assessment toolkit was then used in a feasibility study to evaluate recruitment rates, procedure acceptability, and time to assess (manuscript in preparation of submission).

## **Objectives**

The objectives of this method comparison study are twofold: 1) to evaluate sensitivity of the virtual assessment against the gold standard of the in-person assessment; 2) and to determine inter-rater and intra-rater reliability of the measures in the virtual assessment. This study will further

contribute to the development of a virtual toolkit of assessments across four physical domains along with the evaluation of effort and is specifically designed for the WSIB context.

## Methods

This study will follow a prospective method-comparison study design [17].

### *Ethical Considerations*

Ethics approval was obtained from the Ottawa Health Sciences Network Research Ethics Board (20230311-01H) the Bruyère Research Ethics Board (M16-22-006) and the University of Ottawa Board of Ethics (H-06-23-9348). All potential risks and adverse consequences will be relayed to participants verbally and all participants will provide verbal consent over the telephone or informed consent in-person prior to their scheduled appointment. A wet ink signature will be obtained from all participants on the day of their scheduled assessment. Privacy and confidentiality will be maintained by using de-identified data. Patient-participants will be provided with a \$30 gift card and parking voucher as a token of appreciation for their participation in the study.

### *Participants*

#### *Patient-Participants.*

Eligible participants will be adults aged 18 years or older who have sustained an acquired brain injury (ABI) (concussion or other form of ABI) and are being assessed by a physician, physician assistant, and/or physiotherapist at the Ottawa Hospital located in Ontario, Canada. In addition to people with concussions, people living with ABI including moderate to severe traumatic brain injury (TBI) or hypoxic brain injury will be recruited. In this study, we are assessing the consistency of physical examination findings, and people with other forms of ABI, such as severe TBI, more frequently have evidence of central nervous system deficits, whereas these assessments are often normal during an examination of a person post-concussion [20,21]. This inclusion of other forms of ABI is intended to ensure recruitment of participants with a broad range of identifiable deficits on all components of the assessment. Those unable to complete the study assessments, such

as those unable to verbally communicate symptoms with the clinician, and unable to speak English or French will be excluded from the study. No additional exclusion criteria will be imposed on the participants. Participant information will be documented including demographics, functional status, environmental aids used, socioeconomic and work status, injury information, and information related to technology experience.

#### *Clinician-Participants.*

Clinicians with expertise in conducting assessments for people with ABI will be invited to participate in this project. Demographic information for participating clinicians will be documented, including age, years practicing with ABI patients, professional background (physiotherapist, physician assistant, physiatrist, other) and self-reports of competency with in-person neurological assessments and virtual neurological assessments.

#### ***Sample Size***

The target sample size of the method-comparison study is 60 patient-participants. For the objective of assessing sensitivity of the virtual assessment against the gold standard of in-person assessment, a total of 60 participants, of which 50 (83%) are expected to have an abnormal finding, and with an estimated sensitivity ranging from 77-96% [22], the two-sided 95% confidence interval around the estimated sensitivity will have a total width ranging from 8.7% to 25.3% (or margin of error  $\pm 4.4\%$  to  $12.7\%$ ) [23]. For the objective of assessing inter-rater agreement between two clinicians conducting a virtual assessment using the kappa statistic, a sample size of 60 participants achieves 80% power to detect a true kappa value of 0.89 using a one-sided test at the 5% significance level assuming that 90% of participants are classified as abnormal. Kappa under the null hypothesis was specified as 0.5.

#### ***Recruitment and Consent***

##### *Patient-Participants*

Purposive sampling methods [24] will be used to recruit individuals living with ABI with

impairments spanning all the domains of interest. The participants will be recruited through the Ontario Workers Network and relevant rehabilitation clinics at the Ottawa Hospital. Electronic medical records will be screened to identify eligible patients, who will then be called or approached in-person for consent.

### *Clinician-Participants*

Clinicians will be recruited from the Ontario Workers Network and relevant rehabilitation clinics at the Ottawa Hospital to participate as clinician assessors in the study. Consent will be obtained over the telephone or face-to-face.

### **Outcomes**

The virtual toolkit of assessments will include the following outcome measures: the classification of abnormality (binary variable) on the finger-to-nose test, Vestibular/Ocular Motor Screening (VOMS) tool [25], balance testing, cervical spine range of motion, saccades testing, and evaluation of effort. The procedures outlined below provide details for these outcome measures, including how the sensitivity of the virtual assessment when compared to the in-person assessment and the inter-rater and intra-rater reliability of the virtual assessment will be assessed.

### **Procedures**

#### *Training*

Training of the assessing clinicians on the outcome measures will occur prior to commencement of patient-participant recruitment. A previously developed training manual [26] will be used to complete the training. Training will be uniform with the goal of standardizing clinician completion of the assessments.

#### *Assessments*

Data collection will occur in REDCap (Research Electronic Data Capture), a secure electronic data capture application. An assessing clinician will conduct two assessments: one in-person and one virtual. Both assessments will be identical consisting of all measures from the virtual

assessment toolkit (see Table 1). Both assessments will be completed at the Ottawa Hospital. When possible, assessments will occur on the same day with a brief rest period in between [27,28]. The order of assessment completion (in-person versus virtual) will be randomized and counter-balanced. A random numbers table will be generated and imported into REDCap and the randomization will occur through REDCap. In addition to the assessments identified in Table 1, clinicians will document adverse events (if any) and will record findings from the assessments on a paper copy, which will be directly inputted into REDCap by a research team member.

**Table 1:** Outline of measures included in the virtual assessment toolkit and to be completed in both assessments\*

Domain	Measures	Documentation of Findings by Clinicians	Properties from the Literature
<b>Neurological Examination</b>	Coordination: finger-to-nose	Abnormal (hesitation, tremor, over or undershooting [29])/Normal	Stroke: Sensitivity: 0.71 [30] TBI: Test-retest Kappa reliability: 0.54 [31]
<b>Vestibular</b>	VOMS	Change in symptoms ( $\geq 2$ point increase indicates abnormality [25]) Average distance for three trials of near point convergence in centimeters ( $\geq 5$ cm convergence indicates abnormality [25])	Concussion: Sensitivity: 77-96% [22] Test-retest ICC reliability: 0.60-0.80 [32]
	Balance (feet together, single leg stance, tandem stance) eyes open and eyes closed for 20 seconds each	Abnormal (unable to complete 20-second balance [26])/Normal	TBI: Single leg stance: Sensitivity: 0.45 [33] Test-retest Kappa: 0.43-0.63 [34]
<b>Oculomotor</b>	Saccades	Abnormal (observe saccade initiation, range of motion and conjugacy, speed, accuracy, intrusions or oscillations [35])/Normal	TBI: Sensitivity: 64-77% [36] Pediatric concussion: Test-retest Kappa reliability: 0.7 [37]
<b>Cervical</b>	Range of motion (flexion, extension, right and left lateral flexion, right and left rotation)	Estimated angles for each recorded (abnormality determined by comparison to norms in people aged 20-59 years): flexion= 50-72°; extension=58-77°; lateral flexion=37-47°; rotation=67-81° [38])	Whiplash: Sensitivity: 86% [39] Orthopedic Disorders: Within-tester ICC reliability: 0.78-0.90 Between-tester ICC reliability: 0.54-0.79 [40]
<b>Effort</b>	Rating scale	Full employment of effort documented as Yes/No (clinician discretion) Rating scale documented from 0 (no effort) to 10 (maximum effort) (clinician discretion)	Not reported

ICC: intraclass correlation coefficient; TBI: traumatic brain injury; VOMS: Vestibular/Ocular Motor Screening

\*For a complete presentation of the instructions used to administer the measures, see Johnston et al. [(26)]

For the virtual assessment, the patient-participants will be located in a separate room from the assessing clinicians, within the Ottawa Hospital. All patient-participants will use the same computer and will be provided with technical support from a research team member, as needed. The virtual assessments will be conducted in the clinical environment to standardize as many components of the assessment as possible including computer screen, lighting, image size, room setup, etc. During the virtual assessment, a number of safety measures will be implemented: i) a research team member will be present in the room throughout; and ii) the patient-participant will stand in front of a wall/bed/chair or with their back at a corner during the vestibular assessments in case of loss of balance. Further, the clinician completing the assessment will be able to support the patient-participant if extreme exacerbation of symptoms arises. All the virtual assessments will occur through the Microsoft Teams platform. The study assessments will be audio-video recorded using the record function on Microsoft Teams.

Where possible, both assessments will be completed on the same day. It is estimated that a portion (~5%) of participants may be unable to complete both the in-person and virtual assessment on the same day, either due to lack of time or inability to tolerate a second assessment as a result of symptom aggravation (e.g, headaches, dizziness, nausea, sensitivity to noise and light, etc.). For these participants, the second assessment (virtual or in person) will be scheduled for completion within one week of the initial assessment.

Following completion of the assessments, the patient-participants and clinician-assessors will complete a feedback form. Participants will complete a feedback form (see Appendix A) related to completion of the virtual assessment, the environmental set-up, and confidence in the assessment findings.

#### *Observation and Rating of Audio-Video Recordings*

A second clinician (Rater B: Figure 1) will complete two ratings of the recording of the virtual assessment (in a randomized order for participants) at one-month intervals. Clinicians will record

findings from the assessments on paper or electronic copies, which will be inputted directly into REDCap by a research team member. See Figure 1 for an outline of study assessment procedures.




Figure 1. Assessment procedures modified from Russell et al. [17]

## Analysis

Patient and clinician participant characteristics will be descriptively analyzed.

### Assessments

IBM SPSS (version 28) will be used to calculate sensitivity of virtual assessments versus in-person assessments, as well as kappa (inter-rater and intra-rater reliability) of the virtual assessments (see Figure 1 for comparisons).

### **Method-Comparison (Sensitivity)**

The sensitivity of virtual assessment compared to the reference standard in-person assessment (Rater A in-person and Rater A virtual) will be determined where the assessment of the uncertain classification of the measures used virtually will occur against the reference standard or 'true' classification, which is the in-person assessment in this study. The findings of the initial clinician assessor (Rater A) that completed the in-person and virtual assessments will be compared to

determine sensitivity of the measures administered virtually. Sensitivity (true positives divided by the sum of true positives and false negatives multiplied by 100) [41], in this case, refers to the ability of the virtual measure to identify a deficit or positive finding when the in-person measure identifies the deficit or positive finding. Values closer to 100% indicate greater sensitivity.

### ***Reliability***

To calculate inter-rater reliability of the virtual assessment, the findings of rater A (virtual assessment) will be compared to the findings of rater B<sub>1</sub> (initial observation of virtual assessment recording). To determine intra-rater reliability of the virtual assessment, the findings of rater B<sub>1</sub> (initial observation of virtual assessment) will be compared to the findings of rater B<sub>2</sub> (second observation of virtual assessment approximately one month after initial observation). Estimates of reliability will be calculated using Kappa (classification of abnormal versus normal) [42] for all measures included in the toolkit.

### ***Adverse Events***

Adverse events, type and severity, will be summarized by assessment type (in-person or virtual). Adverse events will be considered a significant worsening of symptoms requiring additional medical attention, such as a visit to the emergency department, and new injuries [43].

## **Results**

As of May 2024, we enrolled 39 patient-participants and seven clinician-assessors. Recruitment and testing are expected to be completed by September 2024.

## **Discussion**

This study will establish the reliability and sensitivity properties associated with five physical components of the virtual concussion assessment. It is anticipated that properties will vary from moderate to strong. There is an identified gap in the current knowledge with limited research conducted on the reliability and comparability of virtual concussion assessments in comparison to in-person assessments. While virtual care has demonstrated to be a valuable approach to concussion



assessment and management [19, 44], there is a need for additional information regarding the psychometric properties of outcome measures. This information is essential for instilling confidence in the administration of these measures in a practical setting. Virtual approaches to care have been used for many years, however, the COVID-19 pandemic increased their use significantly in an effort to reduce hospital visits [15]. While the COVID-19 pandemic initiated this increased use of virtual care, some assessments are continuing to be offered virtually. Using measures with acceptable properties in a virtual context is important in practice to ensure accurate identification of problem areas on virtual assessments so that treatment interventions can be appropriately directed [45]. Despite measures showing adequate properties when administered in-person, they may produce different results when administered in a virtual context [17, 46], providing a rationale to carry out this work. The aim of completing this method comparison study is to obtain preliminary information on the virtual assessment in relation to the WSIB context.

The outcomes of this study may inform future policies regarding virtual care. If the virtual concussion assessment demonstrates effectiveness in identifying deficits compared to in-person assessments and maintains consistency in findings, governments and insurers could support the offer of such services to patients in need, with appropriate compensation for clinicians [47]. Future work should aim to expand on the toolkit explored in this work to include other concussion measures, such as cognitive and emotional measures, and to include other demographics (pediatrics).

### *Strengths and Limitations of Methodological Approach*

The strengths of this method-comparison approach include the utilization of commonly used technology and software for virtual brain injury assessments in a practical setting. Further, including a wide range of ABI severities in our sample ensures inclusion of abnormality spanning all relevant domains. This makes the findings not only applicable to current practice, but also potentially generalizable to different types of ABI.

We plan to assess the virtual evaluation in a controlled hospital setting. Moving forward, it

would be beneficial to evaluate the virtual assessment in environments that are less standardized, such as patients' homes or community clinics where physician-specialists are in a different location than the patient.

Finally, due to constraints in clinic scheduling, a methodological decision was made to have the same clinician (Rater A) conduct both the in-person and virtual assessments with the participants. In attempts to address this concern, the order of study assessments (in-person and virtual) will be randomized and counter-balanced.

## **Conclusion**

The method-comparison study is an important endeavour, considering the limited exploration of virtual concussion assessments in the literature despite their increasing use in practice. The study aims to determine whether virtual and in-person assessments yield similar results in terms of identification of abnormality in common physical domains of a concussion assessment. The study will also assess the inter-rater and intra-rater reliability of the virtual assessment, contributing valuable insights to the field of concussion rehabilitation.

## **Funding acknowledgement**

KB is supported by an admission scholarship from the University of Ottawa, the Ontario Graduate Scholarship, and the EPIC-AT Fellowship through AGE-WELL. This Project is funded by a grant provided by the Workplace Safety and Insurance Board (Ontario). The provision of grant support by WSIB does not in any way infer or imply endorsement of the content by the WSIB. MK is supported by an admission scholarship, international doctoral scholarship, Special Merit Scholarship from the University of Ottawa, Hans K. Uthoff, MD FRCSC Graduate fellowship, Ontario Physiotherapy Association Scholarship, and BMO financial Group Graduate scholarship.

## **Data Availability**

Data sharing is not applicable to this article as no data sets were analyzed during the current study.

All data generated or analyzed during the study will included in a future publication once the study is completed.

## Authors' Contributions

All authors have made substantial contribution to the work presented in this paper.

## Conflicts of Interest

None declared.

## Abbreviations

ABI: acquired brain injury

mTBI: mild traumatic brain injury

TBI: traumatic brain injury

VOMS: Vestibular Ocular Motor Screening

WSIB: Workplace Safety and Insurance Board

## References

1. Scorza KA, Raleigh MF, O'connor FG. Current concepts in concussion: evaluation and management. *Am Fam Physician*. 2012;85(2):123-32.
2. Mullally WJ. Concussion. *Am J Med*. 2017 Aug;130(8):885–92.
3. Ventura RE, Balcer LJ, Galetta SL. The concussion toolbox: the role of vision in the assessment of concussion. *In Seminars in neurology*; 35(05): 599-606.
4. Marshall S., Lithopoulos A., Curran D., Fischer L., Velikonja D., & Bayley, M.. *Living Concussion Guidelines: Guideline for Concussion & Prolonged Symptoms for Adults 18 years of Age or Older*; 2023. <https://concussionsontario.org>

5. Iverson GL, Gardner AJ, Terry DP, Ponsford JL, Sills AK, Broshek DK, et al. Predictors of clinical recovery from concussion: a systematic review. *Br J Sports Med*. 2017;51(12):941–8.
6. Kontos AP, Collins MW. Concussion: A clinical profile approach to assessment and treatment. *Wash DC Am Psychol Assoc*. 2018:23-58.
7. Greenberg MS, Wood NE, Spring JD, Gurvits TV, Nagurney JT, Zafonte RD, et al. Pilot Study of Neurological Soft Signs and Depressive and Postconcussive Symptoms During Recovery From Mild Traumatic Brain Injury (mTBI). *J Neuropsychiatry Clin Neurosci* [Internet]. 2015 Jul [cited 2024 May 14];27(3):199–205. Available from: <http://psychiatryonline.org/doi/10.1176/appi.neuropsych.14050111>
8. Murray NG, Szekely B, Moran R, Ryan G, Powell D, Munkasy BA, et al. Concussion history associated with increased postural control deficits after subsequent injury. *Physiol Meas* [Internet]. 2019 [cited 2024 May 14];40(2):024001. Available from: <https://iopscience.iop.org/article/10.1088/1361-6579/aafcd8/meta>
9. Clugston JR, Houck ZM, Asken BM, Boone JK, Kontos AP, Buckley TA, et al. Relationship between the King-Devick test and commonly used concussion tests at baseline. *J Athl Train* [Internet]. 2019 [cited 2024 Jan 9];54(12):1247–53. Available from: <https://meridian.allenpress.com/jat/article-abstract/54/12/1247/433791>
10. Guskiewicz KM. Balance Assessment in the Management of Sport-Related Concussion. *Clin Sports Med*. 2011;30(1):89–102.
11. Mollayeva T, Colantonio A. Concussion in Ontario Health care workers: implication for prevention. Accessed January 26, 2024. [https://www.researchgate.net/profile/Tatyana-Mollayeva/publication/351334405\\_concUssion\\_in\\_ontario\\_HealtH\\_care\\_WorKers\\_implications\\_for\\_prevention/links/6091ac9f92851c490fb6f4a1/concUssion-in-ontario-HealtH-care-WorKers-implications-for-prevention.pdf](https://www.researchgate.net/profile/Tatyana-Mollayeva/publication/351334405_concUssion_in_ontario_HealtH_care_WorKers_implications_for_prevention/links/6091ac9f92851c490fb6f4a1/concUssion-in-ontario-HealtH-care-WorKers-implications-for-prevention.pdf)
12. Kristman VL, Côté P, Van Eerd D, Vidmar M, Rezai M, Hogg-Johnson S, Wennberg RA, Cassidy JD. Prevalence of lost-time claims for mild traumatic brain injury in the working population: improving estimates using workers compensation databases. *Brain Inj*. 2008;22(1):51-9.
13. Terry DP, Iverson GL, Panenka W, Colantonio A, Silverberg ND. Workplace and non-workplace mild traumatic brain injuries in an outpatient clinic sample: A case-control study. de Castro F, editor. *PLOS ONE*. 2018;13(6):e0198128.
14. Colantonio A, Mroczek D, Patel J, Lewko J, Fergenbaum J, Brison R. Examining occupational traumatic brain injury in Ontario. *Can J Public Health*. 2010;101:S58-62.
15. Womble MN, Reynolds E, Kissinger-Knox A, Collins MW, Kontos AP, West RV, Eagle S, Elbin RJ. The emerging role of telehealth for concussion clinical care during the coronavirus (COVID-19) pandemic. *J Head Trauma Rehabil*. 2022;37(2):E49-54.
16. Montes J, Eichinger KJ, Pasternak A, Yochai C, Krosschell KJ. A post pandemic roadmap toward remote assessment for neuromuscular disorders: limitations and opportunities. *Orphanet J Rare Dis*. 2022;17:1-7.

17. Russell TG, Martin-Khan M, Khan A, Wade V. Method-comparison studies in telehealth: Study design and analysis considerations. *J Telemed Telecare*. 2017;23(9):797–802.
18. McLeod TC, Leach C. Psychometric properties of self-report concussion scales and checklists. *J Athl. Train*. 2012;47(2):221-3.
19. Barnes K, Sveistrup H, Bayley M, Egan M, Rathbone M, Taljaard M, Marshall S. Identification of clinical measures to use in a virtual concussion assessment: protocol for a mixed methods study. *JMIR Res protoc*. 2022;11(12):e40446.
20. Fuller G. Neurological examination made easy. (No Title). 1999.
21. Leddy JJ, Haider MN, Noble JM, et al. Clinical Assessment of Concussion and Persistent Post-Concussive Symptoms for Neurologists. *Curr Neurol Neurosci Rep*. 2021;21:1-14.
22. Büttner F, Howell DR, Doherty C, Blake C, Ryan J, Delahunt E. Clinical detection and recovery of vestibular and oculomotor impairments among amateur athletes following sport-related concussion: a prospective, matched-cohort study. *J Head Trauma Rehabi*. 2021;36(2):87-95.
23. Bujang MA, Adnan TH. Requirements for minimum sample size for sensitivity and specificity analysis. *J Clin Diagn Res: JCDR*. 2016;10(10):YE01.
24. Etikan I, Musa SA, Alkassim RS. Comparison of convenience sampling and purposive sampling. *Am J Theor Appl Stat*. 2016;5(1):1-4.
25. Mucha A, Collins MW, Elbin RJ, Furman JM, Troutman-Enseki C, DeWolf RM, Marchetti G, Kontos AP. A brief vestibular/ocular motor screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med*. 2014;42(10):2479-86.
26. Johnston S, Leddy J, Nick Reed OT, Cortel-LeBlanc A, Hafizi R, Laing S, et al. Virtual Concussion Exam Training Manual. [cited 2024 Jan 15]; Available from: <https://cattonline.com/wp-content/uploads/2022/01/Virtual-Concussion-Exam-Manual.pdf>
27. Mani S, Sharma S, Singh DK. Concurrent validity and reliability of telerehabilitation-based physiotherapy assessment of cervical spine in adults with non-specific neck pain. *J Telemed Telecare*. 2021;27(2):88–97.
28. Stewart GW, McQueen-Borden E, Bell RA, Barr T, Juengling J. Comprehensive assessment and management of athletes with sport concussion. *Int.J.Sports Phys.Ther*. 2012;7(4):433.
29. Nishida K, Usami T, Matsumoto N, Nishikimi M, Takahashi K, Matsui S. The finger-to-nose test improved diagnosis of cerebrovascular events in patients presenting with isolated dizziness in the emergency department. *Nagoya J Med Sci*. 2022;84(3):621.
30. Swaine BR, Sullivan SJ. Reliability of the scores for the finger-to-nose test in adults with traumatic brain injury. *Physical therapy [Internet]*. 1993 [cited 2024 Mar 25];73(2):71–8. Available from: <https://academic.oup.com/ptj/article-abstract/73/2/71/2729045>
31. Rodrigues MR, Slimovitch M, Chilingaryan G, Levin MF. Does the Finger-to-Nose Test measure upper limb coordination in chronic stroke?. *Journal of neuroengineering and rehabilitation*. 2017 Dec;14 (1):6.

32. Kontos AP, Monti K, Eagle SR, Thomasma E, Holland CL, Thomas D, Bitzer HB, Mucha A, Collins MW. Test–retest reliability of the Vestibular Ocular Motor Screening (VOMS) tool and modified Balance Error Scoring System (mBESS) in US military personnel. *J Sci Med Sport*. 2021;24(3):264-8.
33. Baracks J, Casa DJ, Covassin T, Sacko R, Scarneo SE, Schnyer D, Yeargin SW, Neville C. Acute sport-related concussion screening for collegiate athletes using an instrumented balance assessment. *J Athl Train*. 2018;53(6):597-605.
34. Vartiainen MV, Rinne MB, Lehto TM, Pasanen ME, Sarajuuri JM, Alaranta HT. The test–retest reliability of motor performance measures after traumatic brain injury. *Advances in physiotherapy*. 2006 Jan 1;8(2):50-9.
35. Termsarasab P, Thammongkolchai T, Rucker JC, Frucht SJ. The diagnostic value of saccades in movement disorder patients: a practical guide and review. *J Clin Mov Disord*. 2015;2:1-0.
36. Hunfalvay M, Roberts CM, Murray N, Tyagi A, Kelly H, Bolte T. Horizontal and vertical self-paced saccades as a diagnostic marker of traumatic brain injury. *Concussion*. 2019;4(1):CNC60.
37. Corwin DJ, Arbogast KB, Swann C, Haber R, Grady MF, Master CL. Reliability of the visio-vestibular examination for concussion among providers in a pediatric emergency department. *Am J Emerg Med*. 2020;38(9):1847-53.
38. Swinkels RA, Swinkels-Meewisse IE. Normal values for cervical range of motion. *Spine*. 2014 Mar 1;39(5):362-7.
39. Dall’Alba PT, Sterling MM, Treleaven JM, Edwards SL, Jull GA. Cervical range of motion discriminates between asymptomatic persons and those with whiplash. *Spine*. 2001;26(19):2090-4.
40. Youdas JW, Carey JR, Garrett TR. Reliability of measurements of cervical spine range of motion—comparison of three methods. *Phys Ther*. 1991;71(2):98-104.
41. Trevethan R. Sensitivity, specificity, and predictive values: foundations, pliabilities, and pitfalls in research and practice. *Frontiers in public health*. 2017;5:308890.
42. Kirshner B, Guyatt G. A methodological framework for assessing health indices. *J Chronic Dis*. 1985;38(1):27-36.
43. Chan C, Iverson GL, Purtzki J, Wong K, Kwan V, Gagnon I, Silverberg ND. Safety of active rehabilitation for persistent symptoms after pediatric sport-related concussion: a randomized controlled trial. *Arch Phys Med Rehabil*. 2018;99(2):242-9.
44. van Ierssel J, O’Neil J, King J, Zemek R, Sveistrup H. Clinician Perspectives on Providing Concussion Assessment and Management via Telehealth: A Mixed-Methods Study. *J Head Trauma Rehabil* [Internet]. 2023 [cited 2024 Jan 15];38(3):E233–43. Available from: <https://www.ingentaconnect.com/content/wk/htr/2023/00000038/00000003/art00011>
45. Fawcett AL. Principles of assessment and outcome measurement for occupational therapists and physiotherapists: theory, skills and application. John Wiley & Sons; 2013.

46. O'Neil J, Barnes K, Morgan Donnelly E, Sheehy L, Sveistrup H. Identification and description of telerehabilitation assessments for individuals with neurological conditions: A scoping review. *Digit Health*. 2023;9:20552076231183233.
47. Thomas-Jacques T, Jamieson T, Shaw J. Telephone, video, equity and access in virtual care. *NPJ Digital Medicine*. 2021;4(1):159.

## **Appendix A Feedback Forms**

Study ID: \_\_\_\_\_

**Patient-Participant Feedback**

1. Do you have any feedback about the environmental set-up of the virtual assessment (screen size, lighting, space, etc.)?

2. Do you think you your clinician obtained the same information in the virtual assessment when compared to the in-person assessment?

3. I feel confident in my clinicians' findings on the virtual assessment:

- ☐ Strongly Disagree  
☐ Disagree  
☐ Neutral  
☐ Agree  
☐ Strongly Agree

4. I feel confident in my clinicians' findings on the in-person assessment:

- ☐ Strongly Disagree  
☐ Disagree  
☐ Neutral  
☐ Agree  
☐ Strongly Agree

5. Do you have any other feedback regarding the virtual assessment?

Version date: February 7, 2023



Clinician ID: \_\_\_\_\_

Patient ID: \_\_\_\_\_

#### Clinician Feedback

1. Do you have any feedback about the environmental set-up of the virtual assessment (screen size, lighting, space, etc.)?

2. Do you think you obtained the same information in the virtual assessment when compared to the in-person assessment?

3. I feel confident in my findings on the virtual assessment:

- ☐ Strongly Disagree  
☐ Disagree  
☐ Neutral  
☐ Agree  
☐ Strongly Agree

4. I feel confident in my findings on the in-person assessment:

- ☐ Strongly Disagree  
☐ Disagree  
☐ Neutral  
☐ Agree  
☐ Strongly Agree


5. Do you have any other feedback regarding the virtual assessment?

Version date: February 7, 2023

## Supplementary Files

## Figures

Assessment procedures modified from Russell et al. [17].



## **Multimedia Appendixes**

Peer Review Report.

URL: <http://asset.jmir.pub/assets/446efbb81648917dcb4dfb7df40a1aff.pdf>

