

Impact of an Electronic Health Record based Interruptive Alert among Patients with Headaches seen in Primary Care: A Cluster Randomized Controlled Trial

Apoorva Pradhan, Eric A. Wright, Vanessa A. Hayduk, Juliana Berhane, Mallory Sponenberg, Leeann Webster, Hannah Anderson, Siyeon Park, Jove Graham, Scott Friedenberg

Submitted to: JMIR Medical Informatics
on: March 25, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

| | |
|-------------------------------------|-----------|
| Original Manuscript..... | 5 |
| Supplementary Files..... | 33 |
| Figures | 34 |
| Figure 1..... | 35 |
| Multimedia Appendixes | 36 |
| Multimedia Appendix 1..... | 37 |
| Multimedia Appendix 2..... | 37 |
| Multimedia Appendix 3..... | 37 |
| Multimedia Appendix 4..... | 37 |
| Multimedia Appendix 5..... | 37 |
| CONSORT (or other) checklists..... | 38 |
| CONSORT (or other) checklist 0..... | 38 |

Impact of an Electronic Health Record based Interruptive Alert among Patients with Headaches seen in Primary Care: A Cluster Randomized Controlled Trial

Apoorva Pradhan¹ BAMS, MPH; Eric A. Wright^{1,2} PharmD, MPH; Vanessa A. Hayduk¹ BS; Juliana Berhane³ MS; Mallory Sponenberg⁴ RN; Leeann Webster⁵ RPh, MBA, CDE; Hannah Anderson¹ MS; Siyeon Park⁶ PharmD; Jove Graham¹ PhD; Scott Friedenber⁷ MD

¹Center for Pharmacy Innovation and Outcomes Geisinger Danville US

²Department of Bioethics and Decision Sciences Geisinger Danville US

³Pharmacy Support Services Geisinger Danville US

⁴Health Information Technology Geisinger Danville US

⁵Enterprise Pharmacy Geisinger Danville US

⁶Pharmesol Inc. Auburndale US

⁷Department of Neurology Neuroscience Institute Geisinger and Geisinger Commonwealth School of Medicine Danville US

Corresponding Author:

Apoorva Pradhan BAMS, MPH

Center for Pharmacy Innovation and Outcomes

Geisinger

100 North Academy Ave

Danville

US

Abstract

Background: Headaches including migraines are one of the most common causes of disability and account for nearly 20-30% of referrals from primary care to neurology. Electronic health record-based alerts have been used in the past within primary care with varying degree of success to influence provider behaviors in multiple disease states except for headaches.

Objective: The aim of this project was to evaluate the impact of an electronic alert implemented in primary care on patients' overall headache management.

Methods: We conducted a stratified cluster-randomized study across 38 primary care clinic sites between December 2021 to December 2022 at a large integrated healthcare delivery system in the United States. Clinics were stratified into six blocks based on region and patient-to-provider ratios and then 1:1 randomized within each block into either control or intervention. Providers practicing at intervention clinics received an interruptive alert in the electronic health record. The primary endpoint was a change in headache burden, measured using the Headache Impact Test 6 (HIT-6) scale, from baseline to six months. Secondary outcomes included change in headache frequency and intensity, access to care, and resource utilization. The analysis was performed to assess the difference in difference outcomes between the arms at follow up at the individual patient level.

Results: We enrolled 203 adult patients with a confirmed headache diagnosis. At baseline the average HIT-6 scores (standard deviation) in each arm were not significantly different (Intervention – 63 (?6.9) vs. Control - 61.8 (?6.6), $P = .21$). We observed a significant reduction in the headache burden only in the intervention arm at follow-up (3.5-point, $P = .009$). The reduction in the headache burden was not statistically different between groups (difference-in-difference estimate: -1.89 (95% confidence interval -5.0, 1.31; $P = .25$). Similarly, secondary outcomes were not significantly different between groups. Only 11% of alerts were acted upon.

Conclusions: The use of an interruptive electronic alert did not significantly improve headache outcomes. Low use of alerts by providers prompts future alterations of the alert and exploration of alternative approaches. Clinical Trial: Registered at clinicaltrials.gov as registration number NCT05067725.

(JMIR Preprints 25/03/2024:58456)

DOI: <https://doi.org/10.2196/preprints.58456>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

✓ **Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.**

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org>, my preprint will be published in a JMIR journal.

Original Manuscript

Impact of an Electronic Health Record based Interruptive Alert among Patients with Headaches seen in Primary Care: A Cluster Randomized Controlled Trial

Apoorva Pradhan, BAMS, MPH¹, Eric A. Wright, PharmD, MPH^{1,2}, Vanessa A. Hayduk, BS¹, Juliana Berhane, MS³, Mallory Sponenberg, RN⁴, Leeann Webster, RPh, MBA, CDE⁵, Hannah Anderson, MS¹, Siyeon Park, PharmD⁶, Jove Graham, PhD¹, Scott Friedenberg, MD⁷

¹Center for Pharmacy Innovation and Outcomes, Geisinger, Danville, PA, USA

²Department of Bioethics and Decision Sciences, Geisinger, Danville, PA, USA

³Pharmacy Support Services, Geisinger, Danville, PA, USA

⁴Health Information Technology, Geisinger, Danville, PA, USA

⁵Enterprise Pharmacy, Geisinger, Danville, PA, USA

⁶Pharmesol Inc, Auburndale, MA, USA

⁷Department of Neurology, Neuroscience Institute, Geisinger and Geisinger Commonwealth School of Medicine, Danville, PA, USA

Funding: This study was sponsored internally at Geisinger.

Word count (excluding title, abstract, acknowledgment, references, tables, and figure legends): 4193

Corresponding Author:

Apoorva Pradhan, BAMS, MPH

Staff Scientist, Geisinger

100 N Academy Ave, MC 44-00

Danville, PA 17822-4400

Phone: 443-301-4414

Fax: 570-714-6633

Email: apradhan1@geisinger.edu

Institutional URL: www.geisinger.org

ABSTRACT:

Background: Headaches including migraines are one of the most common causes of disability and account for nearly 20-30% of referrals from primary care to neurology. In primary care, electronic health record-based alerts offer a mechanism to influence provider behaviors, manage neurology referrals, and optimize headache care.

Objective: This project aimed to evaluate the impact of an electronic alert implemented in primary care on patients' overall headache management.

Methods: We conducted a stratified cluster-randomized study across 38 primary care clinic sites between December 2021 to December 2022 at a large integrated healthcare delivery system in the United States. Clinics were stratified into six blocks based on region and patient-to-provider ratios and then 1:1 randomized within each block into either control or intervention. Providers practicing at intervention clinics received an interruptive alert in the electronic health record. The primary endpoint was a change in headache burden, measured using the Headache Impact Test 6 (HIT-6) scale, from baseline to six months. Secondary outcomes included changes in headache frequency and intensity, access to care, and resource utilization. We analyzed the difference in the difference between the arms at follow-up at the individual patient level.

Results: We enrolled 203 adult patients with a confirmed headache diagnosis. At baseline the average HIT-6 scores (standard deviation) in each arm were not significantly different (Intervention – 63 (± 6.9) vs. Control - 61.8 (± 6.6), $P = .21$). We observed a significant reduction in the headache burden only in the intervention arm at follow-up (3.5-point, $P = .009$). The reduction in the headache burden was not statistically different between groups (difference-in-difference estimate: -1.89 (95% confidence interval -5.0, 1.31; $P = .25$)). Similarly, secondary outcomes were not significantly different between groups. Only 11% of alerts were acted upon.

Conclusions: The use of an interruptive electronic alert did not significantly improve headache outcomes. Low use of alerts by providers prompts future alterations of the alert and exploration of

alternative approaches.

Trial Registration: Registered at clinicaltrials.gov as registration number NCT05067725.

Keywords: Headache Management; Migraine Management; Electronic Health Record based alerts;

Primary Care



INTRODUCTION

Headache (HA) disorders are a major public health concern with migraine being the second most common cause of disability [1]. The frequency of headaches is nearly twice as common in women, adults aged 18-44, people with low family income (< \$35,000), and Native Americans [2]. Migraine and chronic headache patients experience greater than normal use of emergency department (ED) services, with headaches accounting for approximately 3% of all ED visits annually [2]. Headaches also directly impact workplace productivity, with an economic impact estimated to be \$19.3 billion in 2019 [3].

Headaches account for up to 30% of referrals from primary care to neurology. Of these, nearly 50% of patients are ultimately diagnosed with migraines, and the majority have never adequately trialed first-line therapy prior to referral [4-7]. Barriers to primary care management of headaches include time constraints, access challenges, and lack of expertise to accurately diagnose and give appropriate treatment [8,9]. These barriers result in high referrals to specialized care, an easier option than drug initiation titration, monitoring, and adjustments. This leads to prolonged wait times, resulting in patients often remaining unmanaged and seeking care in urgent or emergent settings for episodic management [10,11].

The Geisinger system was an early adopter of the electronic health record (EHR) system (beginning in 1996); with its EHR implemented across all sites of care. The use of clinician decision support (CDS) tools in the form of alerts within the EHR has increased exponentially in recent years [12]. These alerts have been used across different disease states within primary care to provide useful information to clinicians, shape their behaviors, and positively impact patient safety and outcomes [12-14]. Considering the utility of alerts and the challenges described above, we instituted an EHR-based clinician-facing interruptive alert that provides real-time guidance on managing patients with headaches to improve headache care before neurology referral. This includes assistance in collecting information about headache characteristics, guidance on medication management, and the

opportunity to e-consult with neurology providers. The purpose of this pragmatic randomized control trial was to evaluate the impact of the alert on the management of primary headache disorders in primary care settings. We hypothesized that an electronic alert for the management of headaches improves patient-reported headache burden at six months among headache patients managed in primary care.

METHODS

Setting

Geisinger is an integrated healthcare delivery system located in central and northeastern Pennsylvania serving more than one million patients yearly. Geisinger maintains 44 primary care and over 100 specialty clinics, which include 8 neurology practices with 43 neurology physicians all using the same EHR system.

Study Design

We conducted a prospective stratified cluster-randomized controlled trial across 38 primary care sites to assess the impact of an interruptive electronic alert on the management of primary headache disorders. Eligible patients were enrolled and followed for six months post-index encounter to evaluate headache management-related outcomes. This study was an evaluation of a quality improvement initiative that was undertaken at Geisinger with the intent to affect clinical practice. As a result, it was determined as ‘not human subjects research’ by Geisinger’s Institutional Review Board (IRB). It was uploaded to clinicaltrials.gov and can be found using the registration number NCT05067725. The trial followed Consolidated Standards of Reporting Trials (CONSORT) reporting guidelines for randomized trials. We have also reported the e-CONSORT per the journal requirements [15].

Site Randomization and Patient Enrollment

Thirty-eight of forty-four primary care clinic sites were selected for randomization based on their

ability to participate. The sites were first stratified into 9 possible blocks based on two criteria: geographical location (west, central, or northeast); and patient-to-provider ratio (low, moderate, or high). Note that 3 of the 9 possible blocks did not contain any sites, leaving only 6 blocks for this study. Within each block, simple randomization was then used to assign each site to either the intervention or control arm, using a 1:1 ratio. An Excel random number generator was used to generate the simple randomization allocation sequence. Patients received the treatment to which their site had been assigned, making this a cluster-randomized study with sites as clusters (Multimedia Appendix 1).


Patient Selection and Enrollment

Patients were eligible if: (a) the criteria for the alert were met, which included an in-person or telemedicine encounter AND a headache diagnosis or chief complaint of headache (Multimedia Appendix 2); (b) they were 18 years or older, and (c) post-visit criteria were met, which included a confirmed diagnosis of primary headache disorder by manual chart review and a minimum baseline Headache Impact Test 6 (HIT-6) score of 50 or higher or headache frequency of 12 days or greater in the last three months. Clinical input and the scale guidance document were used to determine thresholds for HIT-6 scores and headache frequency[16]. Manual chart reviews were conducted by study team members (HA, AP, and SP) under the guidance of the neurologist (SF) to confirm headache diagnosis. Types of headaches in the analytic sample included migraine, chronic daily, tension-type, cluster, or medication overuse headaches, collectively grouped as “headaches.” Patients were excluded if they had a serious systemic illness (e.g., headaches due to hypertensive emergencies, hepatic or renal failure, cardiac failure), secondary headaches (e.g., concussion), were pregnant or were actively followed with neurology [17]. Patients with serious systemic illness were excluded from the study as it would have been difficult to distinguish whether headaches observed amongst them were due to primary or secondary factors.

Within two weeks after their scheduled visit, eligible patients were contacted by the Survey Research

and Recruitment Core (SRRC) to collect baseline measures (which determined minimum headache eligibility) and to gain verbal consent. Patient contact was attempted seven times. Figure 1 represents the study's process of patient selection, enrollment, and follow-up.



Figure 1: Patient selection, enrollment, and follow-up.

Intervention

EHR-based Interruptive Alert

A clinician-facing interruptive EHR-based alert in the form of a Best Practice Advisory (BPA) was built based on input from an expert panel of neurologists, primary care providers (PCPs), pharmacists, and informaticians. PCPs include primary care physicians and advanced practitioners such as nurse practitioners and physician assistants. The first iteration of the alert was developed based on input received from the needs assessment conducted with the PCPs. This version was piloted at two primary care sites over three months [18]. Face-to-face education of PCPs at the pilot sites was undertaken to familiarize providers with the utility of the alert [18]. A simple pre-post assessment of the 900 eligible patient encounters during this pilot phase demonstrated a significant reduction in neurology referrals post-implementation of the electronic alert. Following the pilot implementation, feedback was collected from 10 PCPs at these sites about the structure, verbiage, and utility of the electronic alert, subsequent modifications were made based on this feedback before full implementation.

The BPA consisted of four parts: 1) a questionnaire to assist the PCP in characterizing the patient's headache (Multimedia Appendix 3); 2) a "Smart Set" guide for ordering medications, imaging, labs, and referrals utilizing evidence-based guidelines; (Multimedia Appendix 3); 3) a hyperlink to allow for electronic consultation with a neurologist; and 4) a link to the "synopsis" which provided a summary of prior patient headache care. The "Smart Set" guide could be accessed directly within the alert by clicking on the "Open Express Lane" button to facilitate quick ordering for providers. The alert appeared within the EHR when a PCP entered a headache diagnosis for an in-person or telemedicine encounter. All actions taken in the alert were recorded, although providers could dismiss without any action. Suppression criteria were built to lock out and avoid repeated firing of the alert. The alert was suppressed for 24 hours if the dismissal reason was headache-controlled and for 365 days for PCP long-term management. The alert could also be dismissed by acknowledging that the provider was previewing the patients' chart, in this case, the alert would refire during the same encounter.

An invisible or silent alert was activated at control sites that would be triggered by identical criteria to the intervention arm, however, this alert was not visible to the clinicians, so no suppression criteria was developed for control sites. The silent alert was developed to help generate a list of eligible patients for the control group that would be comparable to those identified in the intervention arm. Before full implementation, the primary care leadership team was encouraged to disseminate information about the tool to PCPs in the intervention cohort through local departmental meetings and direct emails, including text and short video instructions.

Data Collection

Data sources included the EHR and survey results from patients at baseline, three months, and six months. The surveys conducted by the SRRC included: six questions from the HIT-6 questionnaire, which measures headache impact on a participant's ability to function in day-to-day life[19]; one question on pain intensity; one on the frequency of headaches; three from the Migraine Treatment Optimization Questionnaire (M-TOQ), which determines if a medication treatment for a participant is optimal; and two asking for sociodemographic information that could not be pulled from the EHR [20]. The M-TOQ questions were modified for our study by replacing the word "migraine" with "headache." Prior to study enrollment, the survey was piloted among a sample (n=10) of patients from pilot sites to ensure appropriateness and check for face validity. Patients had the option to opt-out at any time during the study. See Multimedia Appendix 4 for the full survey guide.

Study Outcomes

Primary Outcome- Headache Burden:

Headache burden was assessed using the HIT-6 scale [7]. The scale ranged from 36 to 78, and a score of 50 or greater represented a greater impact and disruption of life caused by headache [21,22]. The primary endpoint was the comparison of changes in HIT-6 scores between arms from baseline to six months.

Secondary Outcomes:

Secondary outcomes included headache frequency, defined as the self-reported frequency of headache days over the previous three months; average headache pain intensity defined by pain on a 10-point analog scale over the same period (0 = no pain at all, to 10 = the worst pain ever); access to care, defined as the proportion of patients referred to neurology; resource utilization defined as the proportion of patients with an ED-visit (all-cause and headache specific); and medication management, defined as the proportion of patients that initiated preventive or abortive medications for headache [7,23]. Referral to neurology was used as a surrogate measure to represent access to neurological care considering that the availability of a timely neurology appointment impacts the volume of referrals.

Implementation Outcomes:

We assessed the adoption of the alert among providers through actions within the alert. Positive adoption was any activity where the provider interacted with the components of the alert such as completing the headache questionnaire, using the Smart Set, or using the synopsis. We also reviewed feedback provided by PCPs from reports generated by a committee that oversees all EHR-based alerts.

Several modifications to the initial analysis plan were implemented following go-live but before analysis. We omitted some secondary outcomes including time to initiation of treatment in treatment naïve patients and MTOQ outcomes. The data on MTOQ was only available at baseline eliminating any pre-post analysis, and the time to initiation of medication in naïve patients was dropped in favor of initiation of treatment in all patients as a binary outcome. We added a referral to neurology due to its relevance to clinical practice. All these changes were updated on the clinical trials.gov website to ensure reporting transparency. We also updated our analytical approach from Generalized Linear Models (GLM) with robust standard errors in the analytical plan to generalized estimating equations (GEE), a specialized GLM approach. After further internal discussion, we felt given that the outcomes data was longitudinal and correlated due to clustering, GEE was better suited.

Sample size calculation

A minimum difference of equal to or greater than a 2.3-point (Standard Deviation (SD) – 4.3) reduction on the HIT-6 scale represents a clinically important and significant change [22]. We needed to enroll at least a total of 66 patients per arm to detect this change with 80% power at a significance level of 0.05, and an intra-class correlation of 0.02. The sample size was calculated using an online calculator developed by the University of California San Francisco [24].

Statistical Analysis

The total number of patients in each arm was counted to ensure a balanced distribution of patient numbers. Descriptive statistics were computed for continuous variables as means or medians with SD or interquartile ranges (IQR), and frequency and percentages for categorical variables when appropriate for baseline patient-reported outcomes, and socioeconomic and demographic characteristics (i.e., sex, age, race, ethnicity, insurance, education, and income). Appropriate statistical tests were used to compare differences in these characteristics between the two groups and within groups: two-tailed independent t-tests or paired t-tests were used for continuous data, χ^2 for categorical data, and Mann-Whitney for non-normal categorical or continuous data.

We constructed GEE for all outcome comparisons between groups adjusting for effects of treatment, phase (baseline, six months), treatment-by-phase interaction, and demographics while accounting for clustering of patients per primary care site. The GEE model with log link was used considering the skewness of data observed for headache days and pain scores (Multimedia Appendix 5). For all the binary variables, a log link with binomial distribution was used, while an identity link with normal distribution was used for HIT-6 scores. As a first step, we performed an unadjusted GEE model with the study groups as only univariate analysis. Multivariable (adjusted) models only included demographic variables that were significantly different between groups ($P < .05$). For the GEE analysis, we reported odds ratios or estimates and associated confidence intervals. All analysis was conducted using SAS Enterprise Guide 8.3.

The analysis was conducted based on the principle of “intention-to-treat (ITT).” All patients who consented and met the inclusion criteria were included in the analysis per the group they were recruited into. The analysis was performed to assess the difference in outcomes at the individual patient level. For missing data via survey collection, we applied a last observation carried forward (LOCF) technique to maximize the number of observations available for analysis [25]. Results from the LOCF technique were further compared to those from the measured data to assess the impact of loss to follow-up.

RESULTS

The alert was fired for a total of 9239 times in 3183 patients (intervention arm: 2677 times in 1507 patients; control arm: 6562 times in 1676 patients) between December 2021 and February 2022. We excluded 1828 patients for not meeting eligibility criteria. We manually verified the diagnosis of primary headache among 729 of the remaining 1355 patients who were eligible for consent and baseline survey assessment. We consented 221 patients, and subsequently enrolled 203 that either had a HIT-6 score of ≥ 50 at baseline or a headache frequency of > 12 days (Figure 1). At the end of six months, follow-up patient-reported data was available for 88 (43%) patients.

All baseline characteristics were similar between groups for patients enrolled in the study except for mean age, which was higher in the intervention group (43 vs. 39 years, $P = .04$) (Table 1). There was no difference in the HIT-6 scores, headache frequency, and intensity, or patient-reported use of medications at baseline between groups (average HIT-6 scores – 63.0 vs. 61.8; median headache days in the past three months – 18 vs. 15; median pain scores – 7 vs. 7; reported use of preventive or abortive medications – 68.5% vs. 58.2%). The majority of the patients were female, white, non-Hispanic, had a commercial insurance plan, an educational qualification of greater than high school, and an annual household income between \$50,000 to \$100,000. Similarly, there was no statistically significant difference in the baseline characteristics observed between groups for patients that

completed the 6-month follow-up.

Table 1: Patient characteristics at baseline

| | Intervention Arm | Control Arm | P-value |
|---|---|---|---------|
| Total patient (N) | 98 | 105 | |
| Demographics | | | |
| Age, in years Mean (SD) | 43.2 (14.4) | 39.1 (14.4) | .04 |
| Sex, N (%) Males Females | 28 (28.6) 70 (71.4) | 21 (20.0) 84 (80.0) | .19 |
| Race, N (%) White African American Other | 95 (97.0) 2 (2.0) 1 (1.0) | 98 (94.2) 2 (1.9) 4 (3.8) | .56 |
| Ethnicity, N (%) Non-Hispanic Hispanic | 92 (93.9) 6 (6.1) | 95 (91.4) 9 (8.6) | .60 |
| Type of insurance at the most recent encounter, N members (%) Commercial Medicare Medicaid Other | 59 (60.2) 13 (13.3) 26 (26.7) 0 | 56 (53.3) 12 (11.4) 31 (29.5) 6 (5.8) | .32 |
| Highest level of education, N (%) < High school/GED High school/GED Some College or technical program 4-year college (BS/BA) Master's Degree (MS/MA/MPH) Doctorate (PhD/ScD or Professional—(MD/DO/JD) Refuse | 9 (9.2) 32 (32.7) 29 (29.6) 16 (16.3) 8 (8.2) 3 (3.1) 1 (1.0) | 9 (8.6) 38 (36.2) 26 (24.8) 20 (19.1) 9 (8.6) 2 (1.9) 1 (1.0) | .98 |
| Annual household income, N (%) \$25,000 or less Over \$25,000-\$50,000 Over \$50,000-\$100,000 Over \$100,000 Refuse | 20 (20.4) 21 (21.4) 34 (34.7) 14 (14.3) 9 (9.2) | 26 (24.8) 27 (25.7) 21 (20.0) 13 (12.4) 18 (17.1) | .12 |
| HIT-6 scores, Mean (SD) | 63.0 (6.9) | 61.8 (6.6) | .21 |
| Number of Headache days, Median (IQR) | 20.5 (35) | 20.0 (35.0) | .87 |
| Pain Scores, Median (IQR) | 7.0 (2.0) | 7.0 (3.0) | .91 |
| Patient-reported use of preventive or abortive medications, N (%) | 63 (68.5) | 57 (58.2) | .14 |

SD = standard deviation, IQR = Interquartile Range

Primary Outcome- Headache Burden:

HIT-6 scores improved significantly from baseline in the intervention arm at six months (3.5 points, $p = .009$) but not in the control group (1.40 points, $P = .23$); additionally, the improvements from baseline did not differ significantly between groups (Table 2). The intra-cluster correlation coefficient observed was 0.411 for HIT-6 scores.

Table 2: Adjusted primary and secondary outcomes of headache burden, frequency, and intensity.

| Outcome | | | N | Mean (SD) | Change | D-I-D Estimate (Regression Coefficients) (95% Confidence Interval) | P-value |
|----------------------|---------------------|-----------------|-----|-------------|-------------------|--|---------|
| HIT-6 Score | Intervention | <i>Baseline</i> | 98 | 63.0 (6.9) | -3.5 ^a | -1.84 (-5.0 – 1.31) | .25 |
| | | <i>Month 6</i> | 38 | 58.7 (8.6) | | | |
| | Control | <i>Baseline</i> | 105 | 61.8 (6.5) | -1.40 | | |
| | | <i>Month 6</i> | 50 | 60.1 (8.7) | | | |
| Headache Days | Intervention | <i>Baseline</i> | 98 | 30.9 (25.8) | -9.2 | 1.22 (0.82 – 1.81) | .24 |
| | | <i>Month 6</i> | 38 | 22.8 (26.2) | | | |
| | Control | <i>Baseline</i> | 105 | 28.9 (23.5) | -9.3 ^a | | |
| | | <i>Month 6</i> | 50 | 18.4 (18.2) | | | |
| Pain | Intervention | <i>Baseline</i> | 98 | 6.9 (1.8) | -0.395 | 1.00 (0.90 – 1.11) | .99 |
| | | <i>Month 6</i> | 38 | 6.5 (1.8) | | | |
| | Control | <i>Baseline</i> | 105 | 6.9 (1.9) | -0.340 | | |
| | | <i>Month 6</i> | 50 | 6.4 (2.1) | | | |

a- $P = .01$, SD = standard deviation, D-I-D = difference-in-difference

Secondary Outcomes:

Change in number of headache days and pain score did not differ between groups (Table 2).

Similarly, compared to baseline, there was no difference in the proportion of patients being referred to neurology, utilizing ED for all-cause or HA-specific reasons, or initiating new abortive or preventative treatment in the six months post-intervention (Table 3).

Table 3: Percent change in the proportion of patients from baseline to six months and adjusted odds ratio estimates for resource utilization and access to care (note: a negative/positive percentage denotes a decrease/increase post recruitment)

| Outcome | Arm | Phase | Utilization, % | % Change | D-I-D Estimate (Odds Ratio) | P-value |
|-------------------------------------|----------------------------|-----------------|----------------|-------------------|-----------------------------|---------|
| ED Utilization (All-cause) | Intervention (n=98) | <i>Baseline</i> | 19.4 | -2.00 | 1.40 (0.58 – 3.36) | .46 |
| | | <i>Month 6</i> | 17.4 | | | |
| | Control (n=105) | <i>Baseline</i> | 14.3 | -4.8 | | |
| | | <i>Month 6</i> | 9.5 | | | |
| Neurology Referral | Intervention (n=98) | <i>Baseline</i> | 2.04 | 12.3 ^b | 0.52 (0.064 – 4.29) | .55 |
| | | <i>Month 6</i> | 14.3 | | | |
| | Control (n=105) | <i>Baseline</i> | 1.90 | 21.0 ^a | | |
| | | <i>Month 6</i> | 22.9 | | | |
| Medication Initiation | Intervention (n=98) | <i>Baseline</i> | 51.0 | 25.5 ^a | 0.91 (0.51 – 1.60) | .73 |
| | | <i>Month 6</i> | 76.5 | | | |
| | Control (n=105) | <i>Baseline</i> | 46.7 | 28.5 ^a | | |
| | | <i>Month 6</i> | 75.2 | | | |
| ED Utilization (HA-specific) | Intervention (n=98) | <i>Baseline</i> | 4.1 | -2.10 | 0.49 (0.051 – 4.72) | .54 |
| | | <i>Month 6</i> | 2.0 | | | |
| | Control (n=105) | <i>Baseline</i> | 3.81 | 0 | | |
| | | <i>Month 6</i> | 3.81 | | | |

a - $P < .001$, b - $P = .002$, D-I-D = Difference-in-difference

Results from the LOCF models for HIT-6 scores, headache days, and pain intensity are listed in Table 4. Findings from the LOCF models were similar to those from the base models presented in Table 2. It was observed that there was a similar significant improvement in HIT-6 scores in the intervention arm, and a reduction in the headache days in the control arm from baseline to six

months, however, the difference between groups was not statistically significant.

Table 4: Adjusted primary and secondary outcomes of headache burden, frequency, and intensity using the LOCF method.

| Outcome | | | N | Mean (SD) | Change | D-I-D Estimate (Regression Coefficients) (95% Confidence Interval) | P-value |
|----------------------|---------------------|-----------------|-----|-------------|--------------------|--|---------|
| HIT-6 Score | Intervention | <i>Baseline</i> | 98 | 63.0 (6.9) | -1.36 ^a | -1.01 (-2.87 – 0.85) | .29 |
| | | <i>Month 6</i> | 98 | 61.0 (8.2) | | | |
| | Control | <i>Baseline</i> | 105 | 61.8 (6.5) | -0.67 | | |
| | | <i>Month 6</i> | 105 | 60.8 (8.4) | | | |
| Headache Days | Intervention | <i>Baseline</i> | 98 | 30.9 (25.8) | -3.57 | 1.07 (0.87 – 1.32) | .49 |
| | | <i>Month 6</i> | 98 | 27.5 (22.9) | | | |
| | Control | <i>Baseline</i> | 105 | 28.9 (23.5) | -4.41 ^a | | |
| | | <i>Month 6</i> | 105 | 24.0 (22.6) | | | |
| Pain | Intervention | <i>Baseline</i> | 98 | 6.9 (1.82) | -0.153 | 1.02 (0.95 – 1.10) | .60 |
| | | <i>Month 6</i> | 98 | 6.6 (2.01) | | | |
| | Control | <i>Baseline</i> | 105 | 6.9 (1.9) | -0.162 | | |
| | | <i>Month 6</i> | 105 | 6.4 (2.1) | | | |

a- $P=.01$, SD = standard deviation, D-I-D = difference-in-difference

Implementation Outcomes:

Of the 2677 alert firings in the intervention arm, 83% were overridden/dismissed, 5% were ignored, and 11% action was taken. Of the 205 PCPs exposed to the intervention alert, 15 provided feedback, 3 found the alert helpful and 12 expressed dissatisfaction due to alert firing on inappropriate patients

or noting the alert was disruptive to workflow. A manual chart review of 1355 patients revealed that 729 (54%) had a confirmed diagnosis of primary headache disorder. The remaining 626 (46%) did not have either a confirmed diagnosis of primary headache disorder or were already followed by neurology for headaches.

DISCUSSION

In this pragmatic clinical trial across 38 clinics, we found no benefit of the electronic alert as designed on headache outcomes. As the alert was based on evidence-based guidelines for diagnosis and treatment, it seems unlikely that the content of the alert was poor and more likely that the lack of impact was due to gaps in implementation.

This is the first study to offer a CDS tool that guided PCPs in diagnosing and appropriately treating patients with different types of primary headache disorders using an episodic electronic alert. Most of the research studies to date focus on a single type of headache disorder, chronic headaches, or migraines [23,26]. As initial preventative treatment for the most common headache types is similar, our alert was designed to be utilized across them as a one-stop access to help diagnose and treat headaches, and e-consult with neurologists if required.

Even though we did notice improvements in several outcome measures in both the intervention and control arms from baseline to six-month follow-up, on a between-groups comparison, the effects did not hold. A similar trend for improvement from baseline in outcomes such as HIT-6 scores and headache days was observed when an LOCF approach was used, but the effect did not hold for between-group comparisons. Using our pragmatic design, we were able to distinguish between expected improvements in headaches over time and those attributed to the intervention using a comparator group, with largely similar characteristics at baseline except age [27]. Past research studies that have evaluated the impact of headache management programs and found improvement in patients' headache burden and resource utilization have often used a pre-post study design which

makes attribution of success to the intervention difficult [17,23]. Our results have substantial implications, as identification of the ineffectiveness of these types of alerts is necessary to determine which should remain active and which should be either deactivated or revised. Only a well-designed test of comparison with a suitable counter-factual arm (i.e., control) can distinguish improvements due to bias (e.g., regression to the mean), versus the intervention itself.

The total number of unique patients for whom the alert fired was relatively similar between groups, but the alert fired 2.2 times more frequently in the control arm. This is to be expected as in the interventional arm, the interruptive alert that was visible to the providers could be suppressed for the duration of the encounter by acknowledging a reason for dismissal, as opposed to the silent or invisible alert in the control arm. Providers also noted several challenges with alert firing such as confusion with multiple components of the alert, concerns over alert firing for inappropriate patients and appearing at times other than during direct patient contact, and trouble dismissing the alert, highlighting flaws with the existing alert design. Chart reviews illustrated that providers often list headaches as not just chief complaints, but also visit diagnoses despite them often being a symptom of other underlying conditions. The inaccurate use of headache as visit diagnosis led to the alert's misfiring in nearly 40% of our patients, it also complicates firing logic for decision support tools like ours. Research shows that the non-specificity of alerts may desensitize clinicians and lead to habituation, thereby lessening their likelihood to follow alert recommendations or guidance [28,29]. Furthermore, even when firing appropriately, providers reported the location of the alert to be disruptive to their normal workflow. Some of the challenges mentioned could be attributed to the providers not having received sufficient education about the alert. The majority of the providers did not recall having seen or received any education about the alert despite efforts to disseminate this information through emails, factsheets, and videos. This highlights an implementation gap and the need for a more engaged educational outreach in primary care settings.

Much of the feedback and low adoption found in our study affirms prior reports of alert fatigue

experienced by providers in primary care settings and is likely to be the main contributor to the lack of significant impact [30]. With only 11% of alerts being acted upon, we can expect only this fraction to benefit from the intervention. Yet, research does not support the notion that prior alert exposure has a carry-forward impact on the care of future patients; suggesting that future recognition of patients and bypassing of alerts does not have a strong persistence of effect on provider action behaviors [28,29,31].

The lack of positive outcomes within the intervention arm was unexpected considering the detailed human-centered design-build, the engagement of pilot primary care providers, and the early feedback received, which was affirmative for the intervention. Slight modifications to the verbiage within the alert were undertaken based on feedback from the pilot providers before its full-scale implementation. Despite this, we noted low use of the features present within the alert and additional adoption challenges that were not fully uncovered in the pilot period. These results are similar to other studies published using human-centered design which failed to translate into adoption and outcome differences [32-35]. These studies have found that better engagement by pilot providers doesn't always translate to all providers at the time of scaling interventions. Additionally, differences in the mechanism used for delivering training between the pilot and full implementation phase also impact the uptake of the intervention [32-35]. Furthermore, the positive findings and feedback from the pilot phase could be influenced by commitment and congruence bias in addition to observation and measurement bias resulting in a Hawthorne Effect of the intervention [36].

Considering the challenges identified in the current study, post-completion we made significant modifications to the tool. First, we redesigned the identification of the patients to make it more specific (but less sensitive) to the targeted patient population. We repositioned the alert to fire at the time of neurology referral as opposed to when assigning a visit diagnosis or chief complaint. We also recognize that an EHR-only intervention is not the only mechanism to address incomplete headache care, and a pilot initiative involving pharmacist-based medication management has been launched.

Research is currently underway to assess the impact of this pilot.

We recognize several limitations to our study. First, given the single-site design of the study, generalizability to other sites is limited and likely influenced by variations in practice environments and standards of practice. Our use of an alert was largely driven by our integrative healthcare environment, the ubiquitous use of EHR across primary care and specialty providers, and the low cost and light maintenance of the alert. Secondly, while a LOCF method was used to reduce the effect of a type II error, the high attrition rate observed might have affected the magnitude of the impact. We recognize that other interventions to address the optimal treatment of headaches do exist and are worthy of consideration. Programs that involved greater manpower and dedicated professionals for follow-up and management, observed significant improvement in their patients' headache burden and headache-related disability or reduced resource utilization, attributed their success to the dedicated personnel [23,26]. Alternatively, programs that solely relied on physician education, or leveraging existing clinical pathways were while at times cost-saving, did not significantly improve patients' headache burden, or quality of life [7,21]. Most of these existing studies have been standalone initiatives that focused on headache-specific work groups with limited generalizability of results, some of which are no longer in effect due to resource and infrastructure constraints. Our objective was to sustainably change the culture of practice surrounding headache care without creating new standalone avenues. In conclusion, our study found that the implementation of an interruptive electronic alert in primary care to aid diagnosis and management of primary headache disorders did not improve headache care for these patients. Low adoption of the tool by the providers has prompted the development of alternative population health-based approaches to improve headache management.

ACKNOWLEDGEMENTS

We would also like to thank Dr. Neil Holland, Ms. Aubrielle Smith, Mr. Alex Skitolsky, Dr. Maria

Kobylinski, the Survey Research and Recruitment Core (SRRC) at Geisinger, and the primary care providers and patients at Geisinger for their contributions to this study.

CONFLICTS OF INTEREST

The authors have no conflicts of interest to disclose.

ABBREVIATIONS:

| Abbreviations | Meaning |
|---------------|---|
| HA | Headache |
| ED | Emergency Department |
| EHR | Electronic Health Record |
| CDS | Clinician Decision Support |
| IRB | Institutional Review Board |
| CONSORT | Consolidated Standards of Reporting Trials |
| HIT-6 | Headache Impact Test 6 |
| PCP | Primary Care Provider |
| M-TOQ | Migraine Treatment Optimization Questionnaire |
| GLM | Generalized Linear Models |
| GEE | Generalized Estimating Equations |
| ITT | Intention-to-Treat |
| LOCF | Last Observation Carried Forward |
| SD | Standard Deviation |
| IQR | Interquartile Range |
| D-I-D | Difference-in-Difference |
| SRRC | Survey Research and Recruitment Core |
| BPA | Best Practice Alert |

AUTHOR CONTRIBUTIONS:

Conception and design were done by Apoorva Pradhan, Eric A. Wright, Leeann Webster, and Scott Friedenberg. Acquisition of data was done by Apoorva Pradhan, Vanessa A. Hayduk, Hannah Anderson, and Siyeon Park. Analysis and interpretation of data were done by Apoorva Pradhan, Juliana Berhane, Mallory Sponenberg, and Jove Graham. All authors (Apoorva Pradhan, Eric A. Wright, Vanessa A. Hayduk, Juliana Berhane, Mallory Sponenberg, Leeann Webster, Hannah Anderson, Siyeon Park, Jove Graham, and Scott Friedenberg) drafted the manuscript, revised it for

intellectual content, and gave final approval of the completed manuscript.

Multimedia Appendix 1: Site-level distribution based on stratification.

Multimedia Appendix 2: Project Inclusion and Exclusion ICD Codes.

Multimedia Appendix 3: Snapshots of the electronic alert questionnaire and Express Lane/Smart Set tool.

Multimedia Appendix 4: Survey administered by Geisinger's Survey Research and Recruitment Core (SRRC).

Multimedia Appendix 5: Distribution of pain scores and the number of headache days observed in the intervention and control arm of the study.

REFERENCES:

1. Stovner LJ, Hagen K, Linde M, et al. The global prevalence of headache: an update, with analysis of the influences of methodological factors on prevalence estimates. *J Headache Pain*. 2022; 23(1):34. PMID: 35410119.
2. Burch R, Rizzoli P, Loder E. The Prevalence and Impact of Migraine and Severe Headache in the United States: Figures and Trends From Government Health Studies. *Headache*. 2018 Apr; 58(4):496-505. PMID: 29527677.
3. Yucel A, Thach A, Kumar S, et al. Estimating the Economic Burden of Migraine on US Employers. *The American journal of managed care*. 2020 Dec 1; 26(12):e403-e408. PMID: 33315334.
4. World Health Organizations. Headache Disorders. 2016 April 8. <https://www.who.int/news-room/fact-sheets/detail/headache-disorders> [accessed May 17, 2023].
5. Tepper SJ, Dahlöf CGH, Dowson A, et al. Prevalence and Diagnosis of Migraine in Patients Consulting Their Physician With a Complaint of Headache: Data From the Landmark Study. *Headache*. 2004 Oct; 44(9):856-864. PMID: 15447694.
6. Fernandes L, Tanti M, Randall M, et al. 111 An audit of headache referrals from primary care, and subsequent management in the outpatient clinic. *Journal of neurology, neurosurgery and psychiatry*. 2019 Dec; 90(12):e33.
7. Rua T, Mazumder A, Akande Y, et al. Management of chronic headache with referral from primary care to direct access to MRI compared with Neurology services: an observational prospective study in London. *BMJ open*. 2020 Oct 16; 10(10):e036097. PMID: 33067273.
8. Matchar DB, Harpole L, Samsa GP, et al. The Headache Management Trial: A Randomized Study of Coordinated Care. *Headache*. 2008 Oct; 48(9):1294-1310. PMID: 18547268.
9. Minen M, Shome A, Halpern A, et al. A migraine management training program for primary care providers: An overview of a survey and pilot study findings, lessons learned, and considerations for

further research. Headache. 2016 Apr; 56(4):725-740. PMID: 27037903.

10. Roy S, Keselman I, Nuwer M, et al. Fast Neuro: A Care Model to Expedite Access to Neurology Clinic. Neurology. Clinical practice. 2022 Apr; 12(2):125-130. PMID: 35747888.

11. Dall T, Storm M, Chakrabarti R, et al. Supply and demand analysis of the current and future US neurology workforce. Neurology. 2013 Jul 30; 81(5):470-478. PMID: 23596071.

12. Powers EM, Shiffman RN, Melnick ER, et al. Efficacy and unintended consequences of hard-stop alerts in electronic health record systems: a systematic review. Journal of the American Medical Informatics Association: JAMIA. 2018 Nov 1; 25(11):1556-1566. PMID: 30239810.

13. Konerman MA, Thomson M, Gray K, et al. Impact of an electronic health record alert in primary care on increasing hepatitis c screening and curative treatment for baby boomers. Hepatology (Baltimore, Md.). 2017 Dec; 66(6):1805-1813. PMID: 28714196.

14. Downing NL, Rolnick J, Poole SF, et al. Electronic health record-based clinical decision support alert for severe sepsis: a randomised evaluation. BMJ Quality & Safety. 2019 Sep 1; 28(9):762-768. PMID: 30872387.

15. Eysenbach G. CONSORT-EHEALTH: Improving and Standardizing Evaluation Reports of Web-based and Mobile Health Interventions. Journal of Medical Internet Research 2011 Dec 31; 13(4):e126-e126. PMID: 22209829.

16. QualityMetric, Inc. and GlaxoSmithKline Group of Companies. HIT-6. <https://headaches.org/wp-content/uploads/2018/02/HIT-6.pdf> [accessed May 4, 2023].

17. Blumenfeld A, Tischio M. Center of Excellence for Headache Care: Group Model at Kaiser Permanente. Headache 2003 May; 43(5):431-440. PMID: 12752747.

18. Patel A, Sponenberg M, Webster L, et al. Using Design Thinking to Understand the Reason for Headache Referrals and Reduce Referral Rates. Neurology Clinical Practice (Forthcoming).

19. Yang M, Rendas-Baum R, Varon SF, et al. Validation of the Headache Impact Test (HIT-6™) across episodic and chronic migraine. Cephalalgia. 2011 Feb 1; 31(3):357-367. PMID: 20819842.

20. Lipton RB, Kolodner K, Bigal ME, et al. Validity and reliability of the Migraine-Treatment Optimization Questionnaire. *Cephalalgia*. 2009 Jul; 29(7):751-759. PMID: 19239676.
21. Smelt AF, Blom JW, Dekker F, et al. A proactive approach to migraine in primary care: a pragmatic randomized controlled trial. *Canadian Medical Association Journal (CMAJ)*. 2012; 184(4):224. PMID: 22231680.
22. Coeytaux RR, Kaufman JS, Chao R, et al. Four methods of estimating the minimal important difference score were compared to establish a clinically significant change in Headache Impact Test. *Journal of clinical epidemiology*. 2006; 59(4):374-380. PMID: 16549259.
23. Harpole LH, Samsa GP, Jurgelski AE, et al. Headache Management Program Improves Outcome for Chronic Headache. *Headache*. 2003 Jul; 43(7):715-724. PMID: 12890125.
24. Kohn MA, Senyak J. Sample Size Calculators [website]. UCSF CTSI. 13 April 2024. <https://www.sample-size.net/> [accessed April 17, 2024].
25. Liu X. *Methods and applications of longitudinal data analysis*: Academic Press; 2016.
26. Veenstra P, Kollen BJ, de Jong G, et al. Nurses improve migraine management in primary care. *Cephalalgia*. 2016 Jul; 36(8):772-778. PMID: 26487468.
27. Austrian J, Mendoza F, Szerencsy A, et al. Applying A/B Testing to Clinical Decision Support: Rapid Randomized Controlled Trials. *Journal of medical Internet research*. 2021 Apr 9; 23(4):e16651. PMID: 33835035.
28. Kawamanto K, Flynn MC, Kukhareva P, et al. A Pragmatic Guide to Establishing Clinical Decision Support Governance and Addressing Decision Support Fatigue: a Case Study. *AMIA Annual Symposium proceedings*. 2018; 2018:624-633. PMID: 30815104.
29. Kawamoto K, McDonald CJ. Designing, Conducting, and Reporting Clinical Decision Support Studies: Recommendations and Call to Action. *Annals of internal medicine*. 2020 Jun 2; 172(11_Supplement):S101-S109.
30. Backman R, Bayliss S, Moore D, et al. Clinical reminder alert fatigue in healthcare: a systematic

literature review protocol using qualitative evidence. *Systematic Reviews*. 2017 Dec 13; 6(1):255. PMID: 29237488.

31. Flottorp S, Håvelsrud K, Oxman AD. Process evaluation of a cluster randomized trial of tailored interventions to implement guidelines in primary care—why is it so hard to change practice?. *Family practice*. 2003 Jun 1; 20(3):333-339. PMID: 12738704.

32. Beets MW, Weaver RG, Ioannidis JPA, et al. Identification and evaluation of risk of generalizability biases in pilot versus efficacy/effectiveness trials: a systematic review and meta-analysis. *The international journal of behavioral nutrition and physical activity*. 2020 Feb 11; 17(1):19. PMID: 32046735.

33. Holeman I, Kane D. Human-centered design for global health equity. *Information Technology for Development*. 2019 Sept 29; 26(3):477-505.

34. Das M, Eichner J. Challenges and Barriers to Clinical Decision Support (CDS) Design and Implementation Experienced in the Agency for Healthcare Research and Quality CDS Demonstrations (Prepared for the AHRQ National Resource Center for Health Information Technology under Contract No. 290-04-0016.) AHRQ Publication No. 10-0064-EF. Rockville, MD: Agency for Healthcare Research and Quality. March 2010.


35. Byrne, C., Sherry, D., Mercincavage, L., et al. Advancing clinical decision support - Key lessons in clinical decision support implementation. Office of the National Coordinator for Health IT, Office of Policy and Planning. 2012.

36. Berkhout C, Berbra O, Favre J, et al. Defining and evaluating the Hawthorne effect in primary care, a systematic review and meta-analysis. *Frontiers in medicine*. 2022 Nov 8; 9:1033486.

Supplementary Files

Figures

Patient selection, enrollment, and follow-up.



Multimedia Appendixes

Site-level distribution based on stratification.

URL: <http://asset.jmir.pub/assets/e68172a62dea73b34a49bc76f88cf062.docx>

Project Inclusion and Exclusion ICD Codes.

URL: <http://asset.jmir.pub/assets/b94ea0b1c9342f5a5bffc92e85aaa2b.docx>

Snapshots of the electronic alert questionnaire and Express Lane/Smart Set tool.

URL: <http://asset.jmir.pub/assets/650773bca6f2f3dcea3064fd67ff536c.docx>

Survey administered by Geisinger's Survey Research and Recruitment Core (SRRC).

URL: <http://asset.jmir.pub/assets/d84767170df6a661c26e7266d64f105f.docx>

Distribution of pain scores and the number of headache days observed in the intervention and control arm of the study.

URL: <http://asset.jmir.pub/assets/7379f3c3408541b52753767c26b61a51.docx>

CONSORT (or other) checklists

E-CONSORT form.

URL: <http://asset.jmir.pub/assets/856130ced41ebef5e2e6a05e5acea712.pdf>