

Exploring the Barriers and Facilitators to Implementing a Smartphone Application for Physicians to Improve the Management of Acute Myocardial Infarctions: A Multi-Centre Mixed-Methods Observational Study

Katelyn J Cullen, Hassan Mir, Madhu K Natarajan, Marija Corovic, Karen Mosleh, Jacob Crawshaw, Mathew Mercuri, Hassan Masoom, JD Schwalm

Submitted to: JMIR mHealth and uHealth
on: May 06, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 21

 Multimedia Appendixes..... 22

 Multimedia Appendix 1..... 22

 Multimedia Appendix 2..... 22

 Multimedia Appendix 3..... 22

 Multimedia Appendix 4..... 22

 Multimedia Appendix 5..... 22

 Multimedia Appendix 6..... 22

Exploring the Barriers and Facilitators to Implementing a Smartphone Application for Physicians to Improve the Management of Acute Myocardial Infarctions: A Multi-Centre Mixed-Methods Observational Study

Katelyn J Cullen^{1, 2*} MPH; Hassan Mir^{3, 4*} MD, MHI, MPH; Madhu K Natarajan^{1, 2, 5} MD, MSc; Marija Corovic^{1, 2} MPH; Karen Mosleh^{1, 2} MSc; Jacob Crawshaw⁶ PhD; Mathew Mercuri^{1, 7} PhD, MSc; Hassan Masoom⁸ MD, MSc; JD Schwalm^{1, 2, 5} MD, MSc

¹ McMaster University Hamilton CA

² Hamilton Health Sciences Hamilton CA

³ University of Ottawa Heart Institute Ottawa CA

⁴ University of Ottawa Ottawa CA

⁵ Population Health Research Institute Hamilton CA

⁶ The Ottawa Hospital Ottawa CA

⁷ University of Toronto Toronto CA

⁸ William Osler Health System Brampton CA

*these authors contributed equally

Corresponding Author:

Hassan Mir MD, MHI, MPH

University of Ottawa Heart Institute
40 Ruskin St
Ottawa
CA

Abstract

Background: The delivery of timely and appropriate care is crucial for survival in patients experiencing ST-elevation myocardial infarction (STEMI). Efficient communication and exchange of test results between the referring emergency medicine (EM) physician or Emergency Medical Service (EMS) paramedic and the interventional cardiologist (IC) is essential to achieving timely care. In many communities, sharing of some information crucial to decision making, such as electrocardiograms (ECG), relies on using fax or text message. The SMART AMI App was developed to streamline communication and ensure that information is shared in a secure manner. The application is simple to use, privacy-compliant, and allows for rapid ECG sharing between healthcare providers, enabling timely decision-making.

Objective: This paper details the results of targeted pre-implementation surveys to establish the barriers and enablers of using a smartphone application to transmit ECG images among ICs, EM physicians and EMS paramedics to help tailor implementation interventions.

Methods: To assess the proposed acceptability and uptake of the application, pre-implementation surveys were disseminated to ICs, EM physicians and EMS paramedics in one region of Ontario, Canada. Questions were generated based on selected components of the Consolidated Framework for Implementation Research, results from a pilot study carried out at a regional hospital where the SMART-AMI App was previously implemented, and predicted barriers based on expert guidance. The pre-implementation surveys consisted of both 7-point Likert scale questions (1=strongly disagree and 7=strongly agree) and open-ended questions. Open-ended data was extracted verbatim and analyzed using an inductive qualitative approach, with transcripts coded into descriptive qualitative codes and then collapsed into themes.

Results: Uptake of the survey was acceptable, with 9 of the invited 10 ICs, 51 of the invited 223 EM physicians, and 93 of the invited 1138 EMS paramedics responding. Survey findings demonstrated a need for an App, as all groups recognized that current practices for sharing ECGs allowed room for improvement, accepting that fax can be inconvenient and text messages may not be secure. When asked whether there was a need for a smartphone application to transmit ECGs, ICs (M=6.67,

SD=0.50), EM physicians (M=5.57, SD=1.30) and EMS paramedics (M=5.79, SD=1.45) consistently agreed. Commonly reported barriers were concerns over technological challenges, privacy issues, and cell phone reception strength. Through identification of the barriers in each stakeholder group, implementation strategies were developed that facilitated the scale-up of this system-change intervention.

Conclusions: Results from the three online pre-implementation surveys to identify key barriers and enablers to implementation of the App helped inform the selection of tailored implementation strategies to support roll-out of the App across the health region. The surveys identified key barriers around technology, privacy concerns, and access to required WiFi that needed to be addressed during App implementation to facilitate uptake and use. Results from the surveys, and ongoing evaluation of effectiveness, are informing expansion of the App intervention to local ambulance services and other health regions. Clinical Trial: <https://clinicaltrials.gov/study/NCT05290389>

(JMIR Preprints 06/05/2024:60173)

DOI: <https://doi.org/10.2196/preprints.60173>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [a JMIR journal](#)

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in [a JMIR journal](#)

Original Manuscript

Exploring the Barriers and Facilitators to Implementing a Smartphone Application for Physicians to Improve the Management of Acute Myocardial Infarctions: A Multi-Centre Mixed-Methods Observational Study

Authors: Katelyn J Cullen*, Hassan Mir*, Madhu K Natarajan, Marija Corovic, Karen Mosleh, Jacob Crawshaw, Mathew Mercuri, Hassan Masoom, JD Schwalm

*Katelyn J Cullen and Hassan Mir contributed equally to this paper and will serve as co-first authors.

Background:

The delivery of timely and appropriate care is crucial for survival in patients experiencing ST-elevation myocardial infarction (STEMI). Efficient communication and exchange of test results between the referring emergency medicine (EM) physician or Emergency Medical Service (EMS) paramedic and the interventional cardiologist (IC) is essential to achieving timely care. In many communities, sharing of some information crucial to decision making, such as electrocardiograms (ECG), relies on using fax or text message. The SMART AMI App was developed to streamline communication and ensure that information is shared in a secure manner. The application is simple to use, privacy-compliant, and allows for rapid ECG sharing between healthcare providers, enabling timely decision-making.

Objective:

This paper details the results of targeted pre-implementation surveys to establish the barriers and enablers of using a smartphone application to transmit ECG images among ICs, EM physicians and EMS paramedics to help tailor implementation interventions.

Methods:

To assess the proposed acceptability and uptake of the application, pre-implementation surveys were disseminated to ICs, EM physicians and EMS paramedics in one region of Ontario, Canada. Questions were generated based on selected components of the Consolidated Framework for Implementation Research, results from a pilot study carried out at a regional hospital where the SMART-AMI App was previously implemented, and predicted barriers based on expert guidance. The pre-implementation surveys consisted of both 7-point Likert scale questions (1=strongly disagree and 7=strongly agree) and open-ended questions. Open-ended data was extracted verbatim and analyzed using an inductive qualitative approach, with transcripts coded into descriptive qualitative codes and then collapsed into themes.

Results:

Uptake of the survey was acceptable, with 9 of the invited 10 ICs, 51 of the invited 223 EM physicians, and 93 of the invited 1138 EMS paramedics responding. Survey findings demonstrated a need for an App, as all groups recognized that current practices for sharing ECGs allowed room for improvement, accepting that fax can be inconvenient and text messages may not be secure. When asked whether there was a need for a smartphone application to transmit ECGs, ICs ($M=6.67$, $SD=0.50$), EM physicians ($M=5.57$, $SD=1.30$) and EMS paramedics ($M=5.79$, $SD=1.45$) consistently agreed. Commonly reported barriers were concerns over technological challenges, privacy issues, and cell phone reception strength. Through identification of the barriers in each stakeholder group, implementation strategies were developed that facilitated the scale-up of this system-change intervention.

Conclusions:

Results from the three online pre-implementation surveys to identify key barriers and enablers to implementation of the App helped inform the selection of tailored implementation strategies to support roll-out of the App across the health region. The surveys identified key barriers around technology, privacy concerns, and access to required WiFi that needed to be addressed during App implementation to facilitate uptake and use. Results from the surveys, and ongoing evaluation of effectiveness, are informing expansion of the App intervention to local ambulance services and other health regions.

Trial Registration: <https://clinicaltrials.gov/study/NCT05290389>

Keywords: ST-elevation myocardial infarction, digital health, mHealth, barriers and enablers

Introduction

Each year, approximately 7,000 people in Ontario experience ST-elevation myocardial infarctions (STEMIs) requiring timely reperfusion of the affected area of the heart to restore blood flow through the coronary arteries[1]. Reperfusion can occur by a procedure called percutaneous coronary intervention (PCI) or with thrombolytic medications. When possible, PCI is preferable, and guidelines recommend that a patient presenting to a centre that can perform PCI (PCI-capable centre) should have this procedure performed within 90 minutes of the patient presenting to the hospital. Those who present to a non-PCI-capable centre should be transferred to a centre that can perform this procedure within 120 minutes[2]. However, among the 270 health care institutions in Ontario, only 20 centres have PCI capabilities onsite[3]. Thus, achieving timely care can be challenging, particularly for patients who live in rural areas where access to a PCI-capable centre is limited. Where on-site PCI is not available, patients must be transferred to a partner PCI capable hospital. Patients receiving care above evidence based best practice time thresholds show increased risk of mortality, repeat myocardial infarction, congestive heart failure, and hospitalization[4], [5], [6], [7].

STEMI is diagnosed based on an electrocardiogram (ECG) and care for patients with STEMI requires rapid coordination of several services, including Emergency Medical Services (EMS), emergency medicine (EM) physicians and staff, the interventional cardiologist (IC) and the PCI laboratory staff. This can range across several centres and/or jurisdictions that rarely share electronic medical records. Previous research shows that efficient coordination of these services through timely communication leads to improved outcomes for STEMI patients[2]. The current process for STEMI activation for many centres in Canada relies on a variety of technologies that are poorly integrated or inefficient. For example, STEMI activation at the Hamilton General Hospital (HGH), a high-volume quaternary cardiac care hospital in Ontario, Canada, utilized fax for electrocardiogram (ECG) transmission, which is secure but slow, inconvenient, and sometimes unavailable. To improve communication and efficiency in this pathway, a quick, easy-to-use, privacy-compliant smartphone application was developed, the Strategic Management of Acute Reperfusion and Therapies in Acute Myocardial Infarction (SMART-AMI) smartphone application (App), for rapid ECG sharing between healthcare providers, enabling timely decision-making.

The SMART AMI App has been pilot tested with ICs and select EM physicians since May 2020 for patients presenting to Niagara Health and requiring transfer to HGH for primary PCI. Although preliminary data shows promise [8], additional research is required to promote the technology more widely as an evidence-based innovation for care. Similar smartphone applications with capabilities of sending ECG images to ICs for decision-making have demonstrated STEMI patient outcome improvement and reduction in decision time[9], [10], [11], [12], [13]. However, limited evidence exists from the users' perspectives that assesses barriers and enablers to implementing such an App; perspectives that impact acceptability and sustained App use. To promote the App as the standard means to communicate ECGs in the setting of STEMI patients beyond the pilot sites in the Niagara region, we sought to gather an understanding of the perceived usefulness and acceptability of the technology in routine practice from prospective App users across a wider region (i.e., Southeastern Ontario)[14]. This paper details the findings from the pre-implementation surveys, which will be used to assist

with the potential scale-up and implementation of the App across other provincial STEMI systems in Ontario and beyond.

Methods

Study Design

Development and administration of the survey was led by a collaborative team consisting of medical professionals, public health researchers, and experts in implementation science. A multi-centre cross-sectional study was conducted to assess the need, usefulness, and acceptability of the SMART AMI App in practice through the development and distribution of online surveys to key stakeholders, including ICs, EM physicians, and EMS paramedics who care for patients presenting with STEMI within the HGH cardiac catheterization laboratory catchment area. The catchment area includes Hamilton, Niagara, Haldimand, and Brant regions, which cover an area of 6,473 km² and a population of 1.8 million people[15], [16]. Figure 1 illustrates the partner hospitals from which EM physicians were sent survey invitations, as well as a geographical map depicting the locations where sampled EMS paramedics respond to emergency calls. Interventional cardiologists were sampled from the primary PCI hospital indicated on the map as Hamilton Health Sciences.

Survey Development

In 2021, three surveys were developed and tailored for each stakeholder group (interventional cardiologist, emergency medicine physicians, and EMS paramedics). These surveys were focused on gathering a better understanding of the barriers and facilitators that may impact the successful implementation of the SMART-AMI App intervention on a larger scale (see Supplementary File 1-3 for the surveys). Development of survey questions, analysis and interpretation were guided by results from the Niagara Health pilot study, predicted barriers based on expert guidance from study team members (including implementation science researchers, interventional cardiologists, EMS paramedics and EM physicians) and the Consolidated Framework for Implementation Research (CFIR)[17]. The CFIR is a determinant framework, with a list of 39 constructs organized across five major domains, and provides a practical structure to identify multi-level factors that can impact implementation[18]. CFIR is a widely used framework in implementation science, with great breadth in its application across study design, objectives, settings and outcome measures [19]. Through an iterative process involving input from research team members, questions were drafted and selected for inclusion in the surveys around the potential need for an App to transmit ECG images, the fit of an App into existing workflow processes, the potential for an App to improve patient care and reduce treatment times, and the overall comfort level of using an App in the treatment of suspected STEMI cases. The pre-implementation surveys consisted of both 7-point Likert scale questions and open-ended questions. Likert scales were utilized to identify participant opinions on App implementation and usefulness. We applied a 7-point scale (1=strongly disagree and 7=strongly agree), as this has been shown to reach the upper limits of the scale's reliability, in comparison to a 5-point scale [20]. The surveys developed for each group are fully presented in the Multimedia Appendix 1. Additional questions on the usage of the STEMI activation system (known locally as the Heart Investigation Unit Hotline), were included, however, results from

those questions will not be reported in this paper.

Sample Population

Our study targets the healthcare providers who use the STEMI activation system for a Quaternary Care Hospital to activate and respond to a STEMI. These include ICs, EM physicians and EMS paramedics. All IC and EM physicians working at one of the referring centres (Figure 1)

Survey Distribution

The pre-implementation surveys were hosted in REDCap and questionnaires were designed for anonymous completion. Senior leaders of each survey group utilized a list of physician and EMS paramedic contact details provided by administrative distribution lists to develop the survey sampling frame. Senior leaders were asked to distribute a survey invitation by e-mail to their staff, which included a summary of the evaluation project, a link to the survey, and details about consent; they were also encouraged to send out a reminder email to staff approximately 1 month after the initial invitation was sent.

Data Analysis

Quantitative data were analyzed using descriptive statistics of frequencies through Excel Microsoft 365 ©, with continuous variables presented as means and standard deviations. Open-ended data was extracted verbatim and analyzed using an inductive qualitative approach, with transcripts coded into descriptive qualitative codes and then collapsed into themes using qualitative data analysis software (NVivo 12, QSR International). Initial coding was completed independently by two researchers (KC and KM) and development of themes was done collaboratively by two researchers (KC and KM) with themes then validated by a third researcher (JC), to ensure rigor and enhance trustworthiness of data analysis.

Ethics

Consent was obtained from participants through the use of a question at the beginning of each survey. As survey responses were anonymous, participants could not withdraw their data after submission, however, participants were notified that they could withdraw consent by exiting the survey without submitting any responses. There was no compensation provided for participation. This study was approved by the Hamilton Integrated Research Ethics Board (REB #7691). That approval did not include the ability to publish verbatim survey responses.

Results

Quantitative

Between the period of June and August 2021, senior leaders of each survey group sent out survey requests to physicians and EMS paramedic staff, with response rates varying among each of the stakeholder groups sampled. Of the 10 interventional cardiologists at the Hamilton General Hospital, nine (90%) completed the survey. 51 (23%) of the 223 invited EM physicians

completed the survey, and 93 (8%) of the 1138 invited EMS paramedics completed the survey.

Of the 51 EM physicians, 96% reported that they currently submit ECGs to ICs. Nearly all (98%) did so through text message, while 39% also reported sending ECGs via fax. All nine ICs indicated that they received ECGs via text message, while 33% reported receiving ECGs also via fax. EMS paramedics reported that they rarely submitted pre-hospital ECGs to ICs (7.5%), and physical ECGs were typically exchanged in person at patient hand off.

Among all groups, there was recognition that current practices for sharing ECGs allowed room for improvement, accepting that fax can be inconvenient and text messages are not secure. When asked whether there was a need for a smartphone application to transmit ECGs, ICs ($M=6.67$, $SD=0.50$), EM physicians ($M=5.57$, $SD=1.30$) and EMS paramedics ($M=5.79$, $SD=1.45$) consistently agreed one was needed. Furthermore, among ICs, EM physicians and EMS paramedics, use of the App was believed to fit well into existing workflow processes, with mean Likert responses of 6.67 ($SD=0.50$), 5.65 ($SD=1.31$) and 5.56 ($SD=1.77$), respectively. Staff also noted feeling comfortable with using an App for ECG transmission and review, with mean Likert responses of 6.44 ($SD=0.88$), 5.51 ($SD=1.92$), and 5.22 ($SD=2.05$) reported. Figure 2 highlights viewpoints on the implementation of a smartphone application to transmit ECGs from each group.

Qualitative

Through inductive qualitative analysis, 24 codes were identified for ICs, 46 codes were identified for EM physicians, and 57 codes were identified for EMS paramedics (Table 1). The codes were grouped into four key themes, which housed both barriers and enablers to App implementation: 1) technological issues and constraints; 2) perceived capabilities, alternatives, and comparisons of the App; 3) attitudes and beliefs towards the App; 4) impact on current workflow and requirements for implementation (Table 2).

Commonly reported barriers to implementation were concerns over technological failures within the App. Given STEMI activation is a medical emergency, there were concerns that App updates could be required during urgent scenarios, making the App unusable and thus affect clinical care. EMS paramedic providers and physicians practicing at hospitals with unreliable internet signal also expressed concerns about their ability to reliably call and transmit ECGs using their smartphone. Another important barrier was the uncertainty about the proven risks and benefits of the App as this is a new technology without extensive real-world data. Privacy and security of the tool to transmit PHI was raised. There was concern that the App could take more time than the current clinical workflows thus worsening the process of STEMI activation. Respondents also expressed a desire to follow-up with referring or accepting physicians to discuss the case in future, which is a feature not offered via the App.

Discussion

Principal Results

Responses from the pre-implementation surveys addressed the objectives of this study, which

sought to develop an understanding of the barriers and enablers to implementing an App that facilitates emergency cardiac communication. The primary use case is to securely transmit ECGs images and enable telephone calls directly via the App in the case of suspected STEMI in a large health region in Ontario, Canada. Although previous studies have examined the diagnostic utility of ECG image transmission via an app for STEMI care [11], [21], [22], this paper is the first to report on the barriers and enablers to implementing such an App from target users, allowing for tailored implementation strategies to be utilized in future to increase uptake. Overall, survey findings demonstrated agreement on the perceived need for an App, and consensus that the App would fit well within existing workflow processes, reduce time to STEMI treatment, and reduce inappropriate activation of the on-call IC team. Commonly reported barriers were concerns over technological challenges, privacy issues, and cell phone reception strength.

While all three groups shared concerns over potential technological challenges, privacy issues and cell phone reception strength, there were differences in the barriers and enablers reported by group (Table 2). As the App was previously implemented in the IC group during the pilot phase in 2020, several of the barriers expressed by the EM physicians and EMS paramedics were not reported by the ICs. For example, the ICs were aware of App functionality and ECG image quality and so did not report these as a barrier to use. However, the majority of EM physicians and all EMS paramedics had no prior experience with using the App at time of survey collection, and so responses illustrate apprehension of the unknown. Across each stakeholder group, common enablers were the expected speed of ECG image sharing and the associated timely access to that knowledge distribution. All groups reported comments that were both in favour and against App implementation. Given the rapid growth of smartphone technology in recent decades and its increasing adoption among consumers, it is unsurprising that survey responses reflected a high level of comfort with using an App to transmit ECG images. Furthermore, mobile health technology has the potential to enhance system efficiencies and streamline workflow processes, which may account for our findings indicating both a recognized need for an App to transmit ECG images and its perceived compatibility with existing practices. From the barriers and enablers that were identified in the survey responses, select CFIR constructs were found to be represented more frequently than others (Table 2). Under the technological issues and constraints theme, CFIR constructs such as complexity and knowledge and beliefs about the intervention were commonly referenced. This indicated that the individual's personal perception on how difficult the App would be to use, and their personal attitudes toward utilizing the App for cases of suspected STEMI, were going to be a contributing factor into whether they took the necessary steps to download and use the App. In the perceived capabilities and comparisons theme, CFIR constructs of compatibility and relative advantage were frequently cited. Participants' willingness to use the App depended on whether they saw it as valuable, fulfilling a clinical need, and offering advantages over existing ECG transmission methods. In the attitudes and beliefs toward the intervention theme, there were no dominant CFIR constructs, however, responses indicated that participant perception on the evidence supporting the need for an App to transmit ECGs images influenced whether they themselves would download and use it. Finally, in the impact on current workflow and requirements for implementation theme, complexity and relative advantage were again most commonly referenced, demonstrating that the perceived difficulty of incorporating the App into their current workflow processes and their perception on the benefits to using the App, would impact uptake. Our findings of the barriers identified in the survey responses, combined with our analysis of the most commonly cited CFIR constructs, facilitated the development of implementation strategies for App use and scale-up.

Results from the online surveys allowed for the application of stakeholder-specific implementation science interventions that supported the uptake of the App on a wider scale across the health region. For example, we obtained an understanding of the preferred mechanism of training on App use and applied appropriate education strategies, such as offering Zoom drop-in question and answer training sessions at accessible times during the day, to accommodate shift work hours. Furthermore, we cultivated relationships with opinion leaders (EM Chiefs), early adopters, and sought advice from experts in implementation science research to maximize survey responses, and increase uptake and use of the App. To address password recall concerns among ICs, biometric authentication was introduced to streamline login. Finally, key barriers related to concerns over technological challenges and privacy issues were addressed with the development and distribution of educational materials, such as user guides that highlighted multiple successful security tests from internal privacy, security, and information technology staff along with external third-party vendors. Table 3 illustrates the barriers by theme and the implementation strategies that were utilized to facilitate uptake and use of the App.

This study focuses on a specific region in Ontario, Canada, however, the results can be generalized to other regions or countries where an App is being considered for inclusion in patient care by healthcare providers, as noted by the similarity of findings to published literature. A systematic review of barriers and facilitators to using digital health technologies representing North America, Europe, Asia, Africa and Latin America, highlighted infrastructure and technical barriers as well as privacy and security as commonly reported concerns by healthcare providers when using App-based technology [23]. As reported in our findings, concerns over insufficient network and connectivity speed were found, with a higher prevalence of these barriers being reported by those working in rural or remote regions. Furthermore, a published study on App use among nurses reported similar findings, highlighting perceived usefulness and ease of use as key factors influencing adoption [24]. The alignment of this study's findings with previously published research on barriers and enablers to App use by healthcare providers underscores the importance of engaging users before implementation, which can assist in developing targeted strategies to enhance adoption and integration of new technology.

These findings are relevant to clinical practice improvements as they highlight the potential for mobile health technology to enhance communication in STEMI care and improve coordination among healthcare providers. By addressing identified barriers and leveraging enablers, the successful integration of an App for ECG transmission could improve the efficiency of STEMI activation, reduce unnecessary patient transfers, and ultimately enhance patient outcomes by minimizing delays in critical care. While emerging technology can allow for the integration of a machine learning model to enhance the ECG diagnosis of acute coronary occlusion, research testing the precision and sensitivity of such models combined the judgement of trained health professionals for ECG review [25]. The findings presented in this paper can support the integration of new technologies, including artificial intelligence (AI)-enabled Apps, into acute STEMI care, as without understanding how to effectively implement and maintain an App in the acute management of STEMI, similar barriers are likely to hinder the successful adoption of AI technologies in this setting.

Following this analysis of barriers and enablers to implementation, further research is needed to evaluate the impact of mobile health technology on the care of suspected STEMI cases.

Specifically, it is important to assess whether addressing the identified barriers and leveraging enablers leads to successful implementation strategies that enhance App adoption and sustained use. Further analysis from this implementation will explore data on App uptake, utilization, and effectiveness, with findings to be published in subsequent papers.

Limitations

As the surveys were distributed prior to the majority of EM physicians and EMS paramedics gaining access to the App, there was limited understanding of App features, capabilities and usability among these two groups. This made it difficult for participants to answer certain questions and was reflected in survey responses that noted inadequate knowledge affected their ability to reliably answer some questions. Nonetheless, this provided valuable, unbiased responses to questions aimed at understanding the need for such a tool. Additionally, as links to the survey were shared directly by senior leaders of each stakeholder group, it was challenging for the research team to manage follow-up emails that may have improved survey completion rates, as well as calculate accurate response rates. However, this was the preferred means of communication by the senior leaders to maintain anonymity, confidentiality, and avoiding coercion to respond to questionnaires. It is also possible that response rates were improved as participants were being directly asked to complete the survey by the chief or chair of their department. This study leveraged a convenience sampling strategy, which can introduce selection and sampling bias. As such, caution is necessary in generalization of the results. Reassuringly, there was significant representation from each hospital and paramedic region with similar themes and responses for both the qualitative and quantitative questions.

Conclusions

The implementation of system-level interventions in health care can be affected by unexpected barriers, leading to poor uptake and acceptance. In order to circumvent the challenges of implementing change to an organization's culture, understanding the variables that influence or have an impact on the population's willingness to accept and adopt the change is critical. This study exemplifies the importance of identifying barriers faced by the target population and utilizing that knowledge to create successful implementation strategies to optimize research interventions. Results from the pre-implementation surveys disseminated to ICs, EM physicians and EMS paramedics helped inform the strategies for the SMART AMI App implementation program in Hamilton and the surrounding region. They will be vital in informing active scaling of the App into additional areas throughout the Province and local ambulance services.

Acknowledgments

We wish to thank the Emergency Medicine physician and Emergency Medical Service Paramedic chiefs who assisted in distributing the survey invitation email. Their contribution made it possible to gather important insight into the barriers and facilitators to implementation of the smartphone application technology.

Funding Source

This study was funded through the Hamilton Academic Health Sciences Organization

Innovation Fund received in 2021-2024 (HAH-21-015).

Conflicts of Interest

Dr. Hassan Mir is an inventor and sole owner of the background intellectual property of the smartphone application used in this study. There is no specific financial conflict of interest to declare as the tool is being used for research. However, it may be commercialized in future. None of the other authors have a relevant conflict of interest to declare.

Author Contributions

HM, MKN, JDS, and MM contributed to the design of the study. KJC, KM and MC managed the ethics application, implementation, and evaluation of the App. KJC, KM and JC contributed to the analysis of survey findings. All authors contributed to writing the manuscript and approved the final version.

Data Availability

The datasets generated or analyzed during this study are available from the corresponding author on reasonable request. All relevant data are included in this article.

References

- [1] Ministry of Long-Term Care, "Training Bulletin STEMI Hospital Bypass," 118, Jan. 2016. [Online]. Available: https://www.health.gov.on.ca/en/pro/programs/emergency_health/docs/ehs_training_blltn118_en.pdf
- [2] G. C. Wong *et al.*, "2019 Canadian Cardiovascular Society/Canadian Association of Interventional Cardiology Guidelines on the Acute Management of ST-Elevation Myocardial Infarction: Focused Update on Regionalization and Reperfusion," *Can. J. Cardiol.*, vol. 35, no. 2, pp. 107–132, Feb. 2019, doi: 10.1016/j.cjca.2018.11.031.
- [3] CorHealth Ontario, "Cardiac, Stroke & Vascular Centres." Accessed: Apr. 18, 2023. [Online]. Available: <https://www.corhealthontario.ca/cardiac-stroke-&-vascular-centres>
- [4] L. Lambert, K. Brown, E. Segal, J. Brophy, J. Rodes-Cabau, and P. Bogaty, "Association between timeliness of reperfusion therapy and clinical outcomes in ST-elevation myocardial infarction," *JAMA*, vol. 303, no. 21, pp. 2148–2155, Jun. 2010, doi: 10.1001/jama.2010.712.
- [5] R. L. McNamara *et al.*, "Effect of door-to-balloon time on mortality in patients with ST-segment elevation myocardial infarction," *J. Am. Coll. Cardiol.*, vol. 47, no. 11, pp. 2180–2186, Jun. 2006, doi: 10.1016/j.jacc.2005.12.072.
- [6] B. R. Brodie *et al.*, "When is door-to-balloon time critical? Analysis from the HORIZONS-AMI (Harmonizing Outcomes with Revascularization and Stents in Acute Myocardial Infarction) and CADILLAC (Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications) trials," *J. Am. Coll. Cardiol.*, vol. 56, no. 5, pp. 407–413, Jul. 2010, doi: 10.1016/j.jacc.2010.04.020.

- [7] CorHealth Ontario, "Sharing Best Practices. Timely Access to Reperfusion Therapy for STEMI in Ontario." 2018. [Online]. Available: <https://www.corhealthontario.ca/Sharing-Best-Practices.pdf>
- [8] H. Mir *et al.*, "Smartphone application for STEMI activation: A pilot study," *Can. J. Cardiol.*, vol. 37, no. 10, p. S8, Oct. 2021, doi: 10.1016/j.cjca.2021.07.032.
- [9] J. J. Park *et al.*, "Reduction of Ischemic Time for Transferred STEMI Patients Using a Smartphone Social Network System," *J. Am. Coll. Cardiol.*, vol. 68, no. 13, pp. 1490–1492, Sep. 2016, doi: 10.1016/j.jacc.2016.07.733.
- [10] M. A. Astarcioglu *et al.*, "Time-to-reperfusion in STEMI undergoing interhospital transfer using smartphone and WhatsApp messenger," *Am. J. Emerg. Med.*, vol. 33, no. 10, pp. 1382–1384, Oct. 2015, doi: 10.1016/j.ajem.2015.07.029.
- [11] P. R. Bradley *et al.*, "Smartphone application for transmission of ECG images in pre-hospital STEMI treatment," in *2012 IEEE Systems and Information Engineering Design Symposium*, Apr. 2012, pp. 118–123. doi: 10.1109/SIEDS.2012.6215144.
- [12] C.-C. Chao *et al.*, "Smartphone transmission of electrocardiography images to reduce time of cardiac catheterization laboratory activation," *J. Chin. Med. Assoc.*, vol. 81, no. 6, pp. 505–510, Jun. 2018, doi: 10.1016/j.jcma.2017.11.009.
- [13] M. Studencan *et al.*, "Significant benefits of new communication technology for time delay management in STEMI patients," *PLOS ONE*, vol. 13, no. 11, p. e0205832, Nov. 2018, doi: 10.1371/journal.pone.0205832.
- [14] "Smartphone Application for Pre-Hospital ECG Transmission in STEMI Activation: Design and Rationale," JMIR Preprints. Accessed: May 03, 2024. [Online]. Available: <https://preprints.jmir.org/preprint/55506>
- [15] S. C. Government of Canada, "Profile table, Census Profile, 2021 Census of Population - Toronto, City (C) [Census subdivision], Ontario; Hamilton Niagara Haldimand Brant [Home and Community Care Support Services], Ontario." Accessed: Apr. 27, 2023. [Online]. Available: <https://www12.statcan.gc.ca/census-recensement/2021/dp-pd/prof/index.cfm?Lang=E>
- [16] "Census Profile, 2016 Census - Hamilton Niagara Haldimand Brant, [Health region, December 2017], Ontario and Ontario [Province]." Accessed: Apr. 27, 2023. [Online]. Available: <https://www12.statcan.gc.ca/census-recensement/2016/dp-pd/prof/details/page.cfm?Lang=E&Geo1=HR&Code1=3504&Geo2=PR&Code2=35&SearchText=Hamilton%20Niagara%20Haldimand%20Brant&SearchType=Begins&SearchPR=01&B1=All&GeoLevel=PR&GeoCode=3504&TABID=1&type=0>
- [17] L. J. Damschroder, D. C. Aron, R. E. Keith, S. R. Kirsh, J. A. Alexander, and J. C. Lowery, "Fostering implementation of health services research findings into practice: a consolidated framework for advancing implementation science," *Implement. Sci.*, vol. 4, no. 1, p. 50, Aug. 2009, doi: 10.1186/1748-5908-4-50.
- [18] L. J. Damschroder, C. M. Reardon, M. A. O. Widerquist, and J. Lowery, "The updated Consolidated Framework for Implementation Research based on user feedback," *Implement. Sci.*, vol. 17, no. 1, p. 75, Oct. 2022, doi: 10.1186/s13012-022-01245-0.
- [19] M. A. Kirk, C. Kelley, N. Yankey, S. A. Birken, B. Abadie, and L. Damschroder, "A systematic review of the use of the Consolidated Framework for Implementation Research," *Implement. Sci.*, vol. 11, no. 1, p. 72, May 2016, doi: 10.1186/s13012-016-0437-z.
- [20] M. A. A. Malik *et al.*, "Optimal Reliability and Validity of Measurement Model in Confirmatory Factor Analysis: Different Likert Point Scale Experiment," *J. Contemp. Issues Thought*, vol. 11, pp. 105–112, Jul. 2021, doi: 10.37134/jcit.vol11.9.2021.
- [21] A. Barbagelata *et al.*, "Smartphone ECG for evaluation of ST-segment elevation myocardial

- infarction (STEMI): Design of the ST LEUIS International Multicenter Study,” *J. Electrocardiol.*, vol. 51, no. 2, pp. 260–264, 2018, doi: 10.1016/j.jelectrocard.2017.10.011.
- [22] S. Mahajan *et al.*, “Smartphone based 12 lead ECG as decision support in STEMI: a prospective, cross sectional study, non-randomised, single blinded and single-center study,” *Int. J. Community Med. Public Health*, vol. 10, pp. 1410–1414, Mar. 2023, doi: 10.18203/2394-6040.ijcmph20230916.
- [23] I. J. Borges do Nascimento *et al.*, “Barriers and facilitators to utilizing digital health technologies by healthcare professionals,” *Npj Digit. Med.*, vol. 6, no. 1, pp. 1–28, Sep. 2023, doi: 10.1038/s41746-023-00899-4.
- [24] S. Nezamdoust, M. Abdekhoda, and A. Rahmani, “Determinant factors in adopting mobile health application in healthcare by nurses,” *BMC Med. Inform. Decis. Mak.*, vol. 22, no. 1, p. 47, Feb. 2022, doi: 10.1186/s12911-022-01784-y.
- [25] S. S. Al-Zaiti *et al.*, “Machine learning for ECG diagnosis and risk stratification of occlusion myocardial infarction,” *Nat. Med.*, vol. 29, no. 7, pp. 1804–1813, Jul. 2023, doi: 10.1038/s41591-023-02396-3.
- [26] Ontario Local Health Integration Network, “Hamilton Niagara Haldimand Brant LHIN: 2017-2018 Annual Report.” 2018. [Online]. Available: <https://healthcareathome.ca/wp-content/uploads/2022/11/04-2017-18-HNHBLHIN-AR-EN.pdf>

Abbreviations

CFIR: Consolidated Framework for Implementation Research

ECG: electrocardiogram

EM: Emergency medicine

ERIC: Expert Recommendations for Implementing Change

HGH: Hamilton General Hospital

HHS: Hamilton Health Sciences

IC: Interventional cardiologist

SMART AMI: Strategic Management of Acute Reperfusion and Therapies in Acute Myocardial Infarction

STEMI: ST-elevation myocardial infarctions

Figures and Files Legend

Figure 1: Map of Hamilton General Hospital (HGH) and the Non-PCI Hospitals in the region

Figure 2: Summary of survey responses from interventional cardiologist, emergency medicine physicians, and EMS paramedics

Supplementary File 1: Table 1. Codebook created for the analysis of themes on the barriers and enablers to App implementation

Supplementary File 2: Table 2. Barriers and enablers to implementation

Supplementary File 3: Table 3. Mapping barriers identified from survey responses to implementation strategies

Supplementary File 4: STEMI activation questionnaire (Interventional Cardiologists)

Supplementary File 5: STEMI activation questionnaire (Emergency Medicine Physicians)

Supplementary File 6: STEMI Activation Questionnaire (Paramedics)

Figure 1. Primary PCI hospital STEMI partnership map, adapted from the 2017-2018 Ontario Local Health Integration Network Annual Report[26]





Figure 2. Means and standard deviations on the implementation of a smartphone application to transmit ECGs in cases of suspected STEMI

Mean and Standard Deviation Scores from 7-Point Likert Scale Responses on the Implementation of a Smartphone Application to Transmit ECGs in Cases of Suspected STEMI



Supplementary Files

Multimedia Appendixes

Table 1. Codebook created for the analysis of themes on the barriers and enablers to App implementation.

URL: <http://asset.jmir.pub/assets/5f2a7d829f78d9c65282d4190de43c4d.docx>

Table 2. Barriers and enablers related to the implementation of a smartphone application to transmit ECGs in cases of suspected STEMI across all sampled groups.

URL: <http://asset.jmir.pub/assets/393d97b54f10059c15e3f5cd0fcd7729.docx>

Table 3. Mapping barriers identified from survey responses to implementation strategies.

URL: <http://asset.jmir.pub/assets/9d5445fb610bf42eb861a1d80e80199f.docx>

STEMI Activation Questionnaire: Interventional cardiologists.

URL: <http://asset.jmir.pub/assets/42d8bf89fc976168771bfe2ca200bd30.docx>

STEMI Activation Questionnaire: Emergency Medicine Physicians.

URL: <http://asset.jmir.pub/assets/2e3e1d2aa5271b1ce390d6998de6a5c4.docx>

STEMI Activation Questionnaire: Paramedics.

URL: <http://asset.jmir.pub/assets/a34c2300933e1c9896e9760c75c099e4.docx>