

# **Understanding the occurrence and fate of atmospheric microplastics and their potential risks to human health: A study protocol for cross-sectional analysis**

Shaikh Sharif Hasan, Abdus Salam, Mohammad Moniruzzaman, Aynul Bari, Nirupam Aich, Farjana Jahan, Mahbubur Rahman, Zubayer Islam, Md. Humayun Kabir, Md. Aftab Ali Shaikh, Rubhana Raqib, Sarker Masud Parvez

Submitted to: JMIR Research Protocols  
on: May 07, 2024

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 27

    Multimedia Appendixes ..... 28

        Multimedia Appendix 1..... 28

        Multimedia Appendix 2..... 28

        Multimedia Appendix 3..... 28

        Multimedia Appendix 4..... 28

# Understanding the occurrence and fate of atmospheric microplastics and their potential risks to human health: A study protocol for cross-sectional analysis

Shaikh Sharif Hasan<sup>1\*</sup> MBBS, MPM; Abdus Salam<sup>2</sup>; Mohammad Moniruzzaman<sup>3</sup>; Aynul Bari<sup>4</sup>; Nirupam Aich<sup>5</sup>; Farjana Jahan<sup>1</sup>; Mahbubur Rahman<sup>1</sup>; Zubayer Islam<sup>2</sup>; Md. Humayun Kabir<sup>3</sup>; Md. Aftab Ali Shaikh<sup>3</sup>; Rubhana Raqib<sup>6</sup>; Sarker Masud Parvez<sup>7\*</sup> MPH

<sup>1</sup>Environmental Health and WASH Health System and Population Studies Division icddr,b Dhaka BD

<sup>2</sup>Department of Chemistry University of Dhaka Dhaka BD

<sup>3</sup>Bangladesh Council of Scientific and Industrial Research (BCSIR) Dhaka BD

<sup>4</sup>Department of Environmental & Sustainable Engineering University at Albany State University of New York Albany US

<sup>5</sup>Department of Civil and Environmental Engineering University of Nebraska–Lincoln Lincoln US

<sup>6</sup>Nutrition Research Division icddr,b Dhaka BD

<sup>7</sup>Children's Health and Environment Program, Child Health Research Centre University of Queensland South Brisbane AU

\*these authors contributed equally

## Corresponding Author:

Sarker Masud Parvez MPH

Children's Health and Environment Program, Child Health Research Centre

University of Queensland

62 Graham St

South Brisbane

AU

## Abstract

**Background:** Plastic pollution has reached alarming magnitude, defining the contemporary era as the 'Plastic Age.' Uncontrolled plastic production and inadequate recycling processes have led to widespread contamination of the environment with micro and nanoplastics.

**Objective:** Understanding the extent of environmental exposure to plastic additives and consequential health outcomes for providing knowledge and developing effective control measures.

**Methods:** A cross-sectional exploratory study design was used, consisting of exposed and non-exposed groups in plastic recycling sites in Dhaka city. With a systematic approach involving plastic recycling hotspot detection, management pathway mapping, and detecting the presence of environmental microplastic (MP), the study will establish possible associations between different health consequences and MP exposure. MPs and heavy metals will be detected from environmental samples using fluorescence microscopy, Fourier-transform infrared spectroscopy (FTIR), and inductively coupled plasma mass spectrometry (ICP-MS). Human exposure will be assessed by detecting the metabolites of bisphenol and phthalates from urine samples using Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) and thoroughly evaluating endocrine, reproductive, respiratory, and renal functions. The sample size was derived from the mean concentrations of urinary bisphenol and phthalates metabolites, requiring the participation of 168 respondents. A 1:1 exposure to non-exposed stratification would be sufficient to meet our study objectives, considering the conventional level of power and confidence interval.

**Results:** This study protocol (PR#22111) has received approval from the Research Review Committee and Ethical Review Committee of the icddr,b. We enrolled 84 adult plastic recycling workers with at least five years of exposure history and 84 non-exposed participants who were not involved with plastic recycling activities. We anticipate completing the sample analysis by the second half of 2024.

**Conclusions:** The findings would provide valuable insights into the adverse impacts of microplastic pollution on both the environment and human health, aiding in better understanding the extent of the issue.

(JMIR Preprints 07/05/2024:60289)

DOI: <https://doi.org/10.2196/preprints.60289>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#)

## Original Manuscript

# Understanding the occurrence and fate of atmospheric microplastics and their potential risks to human health: A study protocol for cross-sectional analysis

Shaikh Sharif Hasan,<sup>1</sup> Abdus Salam,<sup>2</sup> Mohammad Moniruzzaman,<sup>3</sup> Aynul Bari,<sup>4</sup> Nirupam Aich,<sup>5</sup> Farjana Jahan,<sup>1</sup> Mahbubur Rahman,<sup>1</sup> Zubayer Islam,<sup>2</sup> Md. Humayun Kabir,<sup>3</sup> Md. Aftab Ali Shaikh,<sup>2,3</sup> Rubhana Raqib,<sup>6</sup> Sarker Masud Parvez,<sup>1,7\*</sup>

1. Environmental Health and WASH, Health System and Population Studies Division, icddr,b, Dhaka 1212, Bangladesh
2. Department of Chemistry, University of Dhaka, Dhaka 1000, Bangladesh
3. Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka 1205, Bangladesh
4. Department of Environmental & Sustainable Engineering, University at Albany, State University of New York, Albany, NY 12222, United States
5. Department of Civil and Environmental Engineering, University of Nebraska–Lincoln, W150E Nebraska Hall, 900 N 16th St Lincoln, NE 68588-0531
6. Nutrition Research Division, icddr,b, Dhaka 1212, Bangladesh
7. Children's Health and Environment Program, Child Health Research Centre, The University of Queensland, South Brisbane, QLD, Australia

**Target Journal:** JMIR Research Protocols (Impact factor: 1.7)

**Corresponding author:**

Sarker Masud Parvez

Child Health Research Centre, The University of Queensland, QLD, Australia

Email: [s.parvez@uq.edu.au](mailto:s.parvez@uq.edu.au)

Mobile: +61480365175

**Keywords:** Plastic pollution, Microplastic, Bisphenol, Phthalate, Environment, Human Health

**Abstract (word count: 276)**

**Introduction:** Plastic pollution has reached alarming magnitude, defining the contemporary era as the 'Plastic Age.' Uncontrolled plastic production and inadequate recycling processes have led to widespread contamination of the environment with micro and nanoplastics.

**Methods:** A cross-sectional exploratory study design was used, consisting of exposed and non-exposed groups in plastic recycling sites in Dhaka, Bangladesh. With a systematic approach involving plastic recycling hotspot detection, management pathway mapping, and detecting the presence of environmental microplastic (MP), the study will establish possible associations between different health consequences and MP exposure. MPs and heavy metals will be detected from environmental samples using fluorescence microscopy, Fourier-transform infrared spectroscopy (FTIR), and inductively coupled plasma mass spectrometry (ICP-MS). Human exposure will be assessed by detecting the metabolites of bisphenol and phthalates from urine samples using Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) and thoroughly evaluating endocrine, reproductive, respiratory, and renal functions. The sample size was derived from the mean concentrations of urinary bisphenol and phthalates metabolites, requiring the participation of 168 respondents. A 1:1 exposure to non-exposed stratification would be sufficient to meet our study objectives, considering the conventional level of power and confidence interval.

**Results:** This study protocol (PR#22111) has received approval from the Research Review Committee and Ethical Review Committee of the icddr,b. We enrolled 84 adult plastic recycling workers with at least five years of exposure history and 84 non-exposed participants who were not involved with plastic recycling activities. We anticipate completing the sample analysis by the second half of 2024.

**Conclusion:** The findings would provide valuable insights into the adverse impacts of microplastic pollution on both the environment and human health, aiding in better understanding the extent of the issue.

## Introduction

Plastic has become essential to our 'throwaway' lifestyle. Due to its ubiquitous presence in our environment, the current era is better known as the 'Plastic Age' [1]. In 2019, around 368 million metric tons of plastics were produced worldwide, which is expected to reach 33 billion tonnes by 2050 [2]. Plastic has invaded our natural resources and ecosystems to become a global environmental and public health threat. It is reported that 79% of global plastic products have yet to be appropriately recycled and are eventually released into landfills or natural environments [3]. About 0.8 million tons of plastic waste are generated annually in Bangladesh, among which 36% are recycled, 39% are landfilled, and the rest are considered leakage or unattended [4]. A study reported that 5 to 13 million metric tons of plastic waste was released into the ocean by 192 coastal countries in 2010 [5]. If urgent action isn't taken, the societal cost of plastic production could hit US\$ 7.1 trillion by 2040, surpassing 2018's global health spending and the combined 2019 gross domestic product (GDP) of Germany, Canada, and Australia [6].

Plastic degrades into powdery fragments and microscopic-sized plastics (less than 5 mm) called microplastics [7]. The pollution of the environment with degraded plastics and microplastics is an emerging concern attracting global attention [8]. Exposure to microplastic particles (MP) results in adverse outcomes, which can be classified into physical effects (such as those related to particle size, shape, and concentration) and chemical effects (involving harmful chemicals associated with MPs) [9]. Cox et al. reported the presence of an average of 0.44 MPs/g of sugar, 0.11 MPs/g of salt, and 0.09 MPs/g of bottled water in the American diet [10]. Typically, humans consume 80 g of MPs/per day via fruits and vegetables that accumulate MPs through uptake from polluted soil [11]. Among numerous toxic chemicals, bisphenol and phthalates are most commonly used as plasticizers [12], where MPs work as vectors for these chemicals. Plastic and its additives harm human health and the environment [13, 14]. Moreover, many heavy metals like zinc (Zn), nickel (Ni), mercury (Hg), cadmium (Cd), nickel (Ni), and lead (Pb) are present in plastic compounds as additives and pigments [15]. Long-term exposure to these metals has adverse health consequences [16-18]. In Bangladesh, MP was detected in different fish and shrimp species from the mangrove estuary and the Northern Bay of Bengal [19, 20]. An abundant number of MP was also detected in the beach sediments of Cox's Bazar, Bangladesh [21]. To our knowledge, limited research exists worldwide on the impacts of atmospheric microplastics, and the study of their human health impacts in Bangladesh is also scarce.



Proper plastic recycling plays a vital role in mitigating the environmental impact of plastic waste and promotes a sustainable future for generations to come. However, the mechanical friction and abrasion of the plastic recycling industries produce tons of MPs [22]. Plastic waste recycling and its harmful effects on human and environmental health should be adequately addressed in Bangladesh. Epidemiological studies indicate that plastic particles and their chemical additives (e.g., bisphenol, phthalates, flame retardants, etc.) may have adverse impacts on human health [23]. However, the underlying mechanism still needs to be clearly understood. Understanding the extent of environmental exposure to plastic additives and consequential health outcomes would provide valuable knowledge and help to develop effective control measures.

**Objectives:**

The fundamental objective of this study is to determine the extent of environmental and human health consequences resulting from exposure to plastics (microplastics) and their associated additives. The specific objectives are: 1) to identify the factors that determine the location of hotspots with higher exposures and concentrations of MPs in Dhaka city; 2) to map the management pathways of plastic waste in Dhaka city specifically from collection to disposal, 3) to characterize the nature of plastics that entry into the environmental system of Dhaka city by identifying the types of MPs (composition, size, and shape), their sources, entry points, and contamination levels in different environmental samples; 4) to evaluate the possible health impacts of MP exposure among workers in the plastic recycling industry.

**Methods:*****Study design and Participant recruitment:***

We will employ a cross-sectional exploratory study consisting of an exposed and a non-exposed comparison group. This study will be conducted at plastic recycling sites in Dhaka City. The research team will identify major plastic recycling sites. Participants will be recruited according to the following inclusion criteria: 1) workers involved in plastic recycling processes for at least five years and 2) aged 18 years or older at the time of data collection. We will exclude irregular part-time workers (e.g., the workers who occasionally work in plastic recycling factories as day labor but primarily rely on other occupations for livelihood) and children under 18 years old. We will also exclude the participants who will be on antibiotic treatment as it may mask symptoms of infection as well as alter the immune response [24, 25]. Data collectors will clarify the study

objectives, procedures, and intended outcomes to the participants and obtain written consent. Additionally, we will select a non-exposed control site within Dhaka city, where no plastic processing factory exists. Since plastic has become embedded in our daily lives, having a population free from plastic pollution's effects seems unreasonable. All non-exposed participants will be other occupational workers, including daily laborers, masons, carpenters, etc., who have had no exposure to the plastic industry.

***Sample size:***

We calculated the sample size based on the primary outcomes, including bisphenol and phthalate metabolites in biological samples. Mean concentrations of urinary bisphenol and phthalate metabolites from studies with similar outcomes were used [26, 27]. Among the phthalate metabolites, urinary mono isononyl phthalate (MiNP) had a mean and standard deviation (SD) of  $10.12 \pm 4.91$   $\mu\text{g/L}$  in plastic recyclers and  $12.22 \pm 4.69$   $\mu\text{g/L}$  in the general population. The estimated sample size was 168 (84 per group) using the 2-sample mean test, considering 80% power and 95% confidence interval.

***Data collection methods:***

The detailed study activities by study objectives are discussed below—

***Objective 01: Hotspots identification***

Hotspot identification is the initial and crucial step in plastic waste management. The field team has identified the plastic recycling facilities across Dhaka and recorded the GPS coordinates. This data was compiled and analyzed using the mapping software ArcGIS Pro (version 3.3; Environmental Systems Research Institute, Inc.). The research team also sought permission from the plastic recycling factory owners to conduct the research within their facilities.

***Objective 02: Mapping the management pathway:***

Plastics are typically distributed into three categories: (1) plastics in active use, (2) post-consumer managed plastic waste, and (3) mismanaged plastic waste [28]. Usually, plastic waste that is managed undergoes disposal methods such as recycling, incineration, or landfilling. However, improperly managed plastic waste is either discarded into the environment or inadequately disposed of and often ends up in rivers/oceans, contributing adversely to the ecosystem [29]. To identify the current challenges within Dhaka city's plastic waste management system, the research team will interview key informants involved in collection, refurbishment, reuse, crude recycling, and disposal. The team will collect information on who collects plastics,

the primary source of collection (such as homes, institutions, dump sites, and transfer stations), the types of plastics collected, transportation methods to the recycling center, which part of plastics are reused and recycled, and the ultimate disposal site and methods of disposal (burying, incineration, chemical melting). Purposive sampling techniques will be used, and data will be collected using two tools: in-depth and key informant interviews. Sampling will continue until information redundancy or saturation is attained [30].

***Objective 03: Quantifying the extent of environmental pollution***

The research team will collect different environmental samples, including floor dust and topsoil (0-5 cm depth) from the plastic recycling and dismantling sites, sediments from the water vicinity (lake/river/pond) where such facilities are operated, and air samples (table 1). The detailed sampling procedures are described below—

**Table 1: Human participants and environmental sampling plan**

| Human subjects    | Proposed number | Environmental samples | Proposed number |
|-------------------|-----------------|-----------------------|-----------------|
| Exposed group     | 84              | Air                   | 186             |
| Non-exposed group | 84              | Soil                  | 20              |
|                   |                 | Dust                  | 20              |
|                   |                 | Sediment              | 10              |

**Air sampling:** The field research team conducted air sampling at two plastic recycling industries/shops and one outdoor location for MPs and heavy metals (HMs) detection. Air sampling was performed using a filter-based low-volume PM<sub>2.5</sub> sampler for heavy metal detection in nylon 6.6 filter paper, and a total suspended particulate (TSP) sampler was used for microplastic detection in the quartz fiber filter. The team set up PM<sub>2.5</sub> and TSP samplers in each selected industry/shop and outdoor location with a duration of six hours and a frequency of once every six days to assess the work-related exposure of factory workers. We collected air samples in both dry (January to March) and wet seasons (June to August) to explore seasonal variation. After sampling, quartz fiber filters were kept in glass filter holders and nylon filters in Ziplock bags. All the samples were placed in a cool box and kept at four °C till transported to the laboratory. The samples were preserved at a negative 20°C temperature at the laboratory for future analysis.

**Soil Sampling:** We will develop a 10x10 meter grid map of selected plastics recycling factories to collect soil samples. The team will collect 20 soil samples from the selected areas of the grid map. We will identify the closest open-air bare soil surface using a whisk broom (non-plastic)

and clean about 1m<sup>2</sup> area. Using the stainless-steel hand trowel, the clean surface will be divided into six equal quadrangles. Six subsamples will be collected from these quadrangles at depths ranging from 0 – 5 cm depth using a stainless-steel auger and placed in a metal tray. After thoroughly mixing the subsamples and removing stones and other non-soil materials, approximately 100 gm of soil sample will be enclosed in aluminum foil, put in a Ziploc bag, and labeled appropriately. Samples will be kept in a cooler box until carrying them to the laboratory, where they will be preserved at negative 20°C for later analysis.

**Dust sampling:** The field team will collect indoor dust samples from the plastic recycling shop/industry where we have recruited our study participants. Our field research team will approach the shop/industry at the end of working hours but before cleaning to collect dust samples. We will sweep the floor under desks, shelves, and other areas, including furniture, machinery, and walls, collecting dust using a dustpan with a pre-cleaned non-plastic brush. Dust from After mixing the sample with a metallic hand trowel, a 100-gm homogenous sample will be collected. These samples will be enclosed in aluminum foil, put in a Ziploc bag, and labeled appropriately. We will keep the samples in a cooler box until carrying them to the laboratory and preserve them at negative 20°C for later analysis.

**Sediment sampling:** The team will collect sediment samples from the nearest water body in the selected plastic recycling industries. Following the same 10x10 meter grid map, the field team will identify ten sampling locations. Surface sediments will be collected at 0-20 cm depth using a metallic handheld sediment sampler. Samples will be collected in a glass jar labeled appropriately, put in a Ziplock bag, and transported to the laboratory for later analysis.

**Environmental sample analysis:** For microplastic detection from environmental samples, an adequate amount of samples will be combined with about 20 mL of 35% H<sub>2</sub>O<sub>2</sub>, heated at 60°C for an hour for organic matter removal. For density separation, 20 mL NaCl solution will be added, ultrasonicated for 20 minutes, and left overnight for organic matter removal. The top solution will be filtered through quartz and examined under an Oxion Inverso microscope (40x to 100x). The filter, post-examination, will be stained using Nile red solution (1-5 mg/mL) and observed under green excitation fluorescence (540-580 nm). Microplastic sizing will be conducted using ImageJ software.

For heavy metal detection, environmental samples will be analyzed using inductively coupled plasma mass spectrometry (ICP-MS) following the acid digestion method. Samples will be dried

at 105°C until reaching a stable weight, followed by disaggregation using a mortar and pestle and then sieving. Following the US Environmental Protection Agency (EPA) Method 3051, fine fraction samples weighing 0.3 g will be subjected to acid digestion using a microwave oven digester. The resulting digested samples will be analyzed using an ICP-MS (NexION® 2000, Perkin Elmer, USA).

#### ***Objective 4: Health risk characterization***

Plastic particles enter into the human body through inhalation, ingestion, and dermal contact [31]. In the human body, they cause oxidative stress [32, 33], cytotoxicity [34, 35], and even translocation to different tissues [36]. There is limited evidence on direct human health effects of MP exposure, but mathematical modeling, animal studies, and in vitro cell cultures revealed potential health risks like altered metabolism, immune disruption, neurotoxicity, reproductive toxicity [37]. Previous biomonitoring approaches have demonstrated the presence of bisphenol, phthalates, and other plastic additives and their metabolites in the human population. This study focuses on two common types of plastic additives— phthalates and bisphenol metabolites and various heavy metals. Based on the literature review, the human health systems most affected are identified and listed in Table 2. To assess the impact on human health, we will collect biological samples, conduct anthropometric measurements, measure blood pressure, and assess the lung function of study participants.

#### ***Biological sample collection, assessment, and survey:***

Upon informed consent, a trained team collected 50 mL of fasting urine samples in urine collection pots to detect bisphenol and phthalate metabolites [38]. Total bisphenol (free and conjugated) and phthalate metabolites will be determined by Liquid Chromatography with Tandem Mass Spectrometry (LC-MS/MS) (6420 LC/TQ, Agilent, Germany). The trained field team also collected 5 mL fasting venous blood samples using trace metal-free vacutainers, and an additional 3 mL was collected using a serum tube. All the samples were carried in a cooler box (4-8°C) during transportation and stored in a negative 80 freezer until analysis.

We also developed a data collection tool and piloted it before applying it in the field. Our research team administered electronic questionnaires, gathering data across multiple domains. This includes general characteristics such as age, gender, income, education, and living environment, as well as detailed exposure histories and attitudes toward plastic waste. We also recorded lifestyle factors like smoking, and medication, along with self-reported health

conditions such as hypertension, diabetes, and liver diseases. Interviews were conducted at the participant's workplaces.

**Table 2:** Major systems affected and potential health impact assessment plan following microplastic and their additives exposure.

| System              | Health-related outcome  | Disease indicators   | Biological Sample | Procedure                                 |
|---------------------|---|--|-------------------|---|
| Female Reproduction | Hormonal imbalance  | ↑ Progesterone, ↑ Luteinizing hormone (LH)   | Fasting blood     | Enzyme-linked immunosorbent assay (ELISA) |
|                     | Undesirable pregnancy outcomes  | ↑ Risk of recurrent miscarriage (RM), ↑ Spontaneous preterm birth (PTB), ↑ Preterm premature rupture of membrane (pPROM) | N/A               | Self-reported questionnaire               |
| Male Reproduction   | Hormonal imbalance  | ↓ Serum total testosterone (TT), ↑ Prolactin   | Fasting blood     | ELISA                                     |
|                     | Sexual dysfunction and Infertility  | ↓ sexual function, ↓ sexual desire, ↓ erectile ability, ↑ premature ejaculation  | N/A               | Self-reported questionnaire               |
| Endocrine           | Metabolic Disorder (Any 3 from 5 features suggestive of metabolic Disorder) | Height, weight, waist, hip circumference.  | N/A               | Anthropometry                             |
|                     |   | Triglycerides  | Fasting blood     | Semi-automatic biochemistry analyzer      |
|                     |   | HDL Cholesterol  | Fasting blood     | Semi-automatic biochemistry analyzer      |
|                     |   | Blood pressure   | N/A               | Automated blood pressure monitor          |
|                     |   | Fasting blood glucose  | Fasting blood     | Hexokinase method                         |
|                     | Thyroid function  | ↓T4, ↓TSH  | Fasting blood     | ELISA                                     |
| Respiratory         | Asthma  | ↓ Lung function  | N/A               | Spirometry                                |
| Renal               | Impaired renal function   | ↑ Albuminuria  | Fasting urine     | Dipstick test for Albumin                 |
|                     |   | Urinary creatinine   | Fasting urine     | Automated hematology analyzer             |
|                     |   | Urinary micro-albumin  | Fasting           | Automated                                 |

|                  |            |   |               |                     |
|------------------|------------|---|---------------|---------------------|
|                  |            |   | urine         | hematology analyzer |
|                  |            | ↓ estimated Glomerular filtration rate (eGFR) | Blood         | CKD-EPI equation    |
| Oxidative stress | DNA damage | 8-OhDG  | Fasting urine | ELISA               |

### ***Blood pressure measurement:***

The team used an automated blood pressure monitor to assess the blood pressure of all participants, ensuring that all necessary precautions were taken care of. Participants were instructed to refrain from consuming caffeine (such as tea, coffee, and carbonated beverages), engaging in heavy physical activities, and smoking for at least 30 minutes before measuring blood pressure. They were then asked to rest for 5 minutes while seated on a chair with arm support. Different cuff sizes and calibrated instruments were used according to participants' age groups. Blood pressure measurements were conducted thrice: initially from the left arm, followed by the right arm, and then once more from the left arm. Systolic blood pressure and diastolic blood pressure readings were noted separately, and then the readings were averaged.

### ***Respiratory measurement:***

The research team collected information on smoking habits and medical history, including any previous diseases. The questionnaire inquired about acute symptoms, such as eye irritation, nasal congestion, sore throat, and headache. Lung function was assessed using a portable digital spirometer, adhering to standard operating procedures. Measurements encompassed forced vital capacity (FVC), forced expiratory volume in one second (FEV<sub>1</sub>), and peak expiratory flow rate (PEFR).

### ***Dietary diversity assessment***

Prior research findings indicate that humans are commonly exposed to microplastics through inhalation, ingestion, and skin contact [23]. Thousands of chemicals and heavy metals are added to plastics as plasticizers; they enter the human body directly with microplastics. It was estimated that an American consumes an average of 39,000 to 52,000 microplastic particles annually, where water was found to be the most prominent source [10]. Even our regular food items like chicken meat and fish were found to be contaminated with microplastics [39]. So, exploring environmental pollution and health consequences will be incomplete without a dietary assessment. Dietary diversity was measured considering the one-week recall period. This assessment will guide us to the probable sources and help us measure the risks more precisely.

**Statistical methods and data analysis:**

Data analysis will be performed using the 'R' programming language (version 3.5.2 for Windows; R Foundation for Statistical Computing) and Stata 14 (StataCorp, College Station, TX, USA). A codebook will be prepared based on data collection questionnaires. All variable labels, value labels, and composite variable calculations will be constructed using do files and 'R' scripts. Throughout the data collection phase, quality control measures will be implemented. Any anomalies will be promptly addressed through discussion and resolution. Data will be summarized using descriptive statistical tools such as the mean/standard deviations for quantitative symmetric variables, median/IQR for quantitative asymmetric variables, and frequencies/proportions for qualitative variables. The Chi-square test will be used to assess the association between two categorical variables [40]; independent t-tests will be used to determine the significance of the differences between the means of two groups for normally distributed data [41], whereas the Kruskal-Wallis rank test will be used to compare the median between two groups or variables for non-normally distributed data [42]. After adjusting for the relevant covariates, inferential analyses such as regression and risk ratio will be calculated to draw more definitive conclusions. Confounding variables will be identified from the literature, and different types of bi-variate analysis will be included in the regression model.

For qualitative data analysis, we will employ ATLAS.ti (version 5.2; ATLAS.ti Scientific Software Development) to organize our data and conduct thematic analysis [43]. Our approach will begin with transcription and familiarization with interview content, followed by selecting significant quotations and identifying keywords. We will develop a coding system by iteratively reading and coding the transcripts to uncover patterns and themes. These coded segments will be entered into ATLAS.ti to facilitate theme identification and further analysis. Themes will be developed by grouping related codes and interpreting their meaning in context, aiming to construct a conceptual model that illustrates connections and insights from the data. Throughout this process, we will compare themes across interviews to ensure comprehensive coverage and triangulate findings for enhanced validity and reliability [44].

**Ethical Consideration:**

The Research Review Committee (RRC) and Ethical Review Committee (ERC) of the icddr,b have reviewed and approved this study protocol (PR#22111). Eligible participants who consent to participate will be asked to provide written consent in the local language (Bengali). There will



be no compensation for their participation in the research. Interviews will be conducted in private or in a comfortable setting to minimize the risk of confidentiality breaches. The interviewers will use culturally appropriate terminology and make every effort to ensure the subjects' comfort. Participants will be informed of their option to discontinue their participation in the study at any time if they feel uncomfortable. To protect participants' privacy, anonymous datasets without personal identifiers will be developed, and deidentified data will be used during storage, analysis, and dissemination.

### Results:

Following approval from the RRC and ERC, the data collectors were recruited, and a comprehensive exposure survey and sample collection training was provided. Before starting fieldwork, we developed and piloted the data collection tool. We initiated air sample collection in January 2023 and completed this phase by August 2023. Simultaneously, biological sample collection and exposure surveys were started in June 2023 and completed by October 2023. Additional environmental samples, including soil, dust, and sediment from the exposure sites, will be collected by May 2024, and analysis is scheduled for the second half of 2024.

We successfully enrolled a total of 84 exposed and 84 non-exposed respondents, with socio-demographic information detailed in Table 3. The mean age of exposed respondents was 34 years (SD: 6.47), slightly younger than the non-exposed respondents (mean, 36 years, SD: 13.3) population. The majority of the respondents were male, and almost half of them did not have any formal education across exposed and non-exposed groups. The BMI of non-exposed respondents was slightly higher than exposed respondents; however, this difference was not significant ( $p \geq 0.05$ ).

Currently, all biological and environmental samples are undergoing analysis in the laboratory, and we anticipate obtaining the results by July 2024.

**Table 3:** Participants characteristics of exposed and nonexposed population.

| Characteristics         | Exposed population<br>(N=84), n (%) | Nonexposed population<br>(N=84), n (%) | P value          |
|-------------------------|-------------------------------------|--|------------------|
| Age in years, mean (SD) | 34.3 (6.47)                         | 35.7 (13.3)                            | .9 <sup>a</sup>  |
| Sex, male, n (%)        | 49 (58.3)                           | 49 (58.3)                              | .01 <sup>b</sup> |
| <b>Education, n (%)</b> |                                     |  |                  |
| No education            | 53 (63.1)                           | 47 (56)                                | .19 <sup>b</sup> |
| Up to primary school    | 28 (33.3)                           | 28 (33.3)                              |                  |
| Above primary school    | 3 (3.6)                             | 9 (10.7)                               |                  |

|  |                   |                   |                  |
|--|-------------------|-------------------|------------------|
| Married, n (%)                         | 72 (85.7)         | 71 (84.5)         | .31 <sup>b</sup> |
| Self-owned home, n (%)                 | 29 (34.5)         | 32 (38.1)         | .52 <sup>b</sup> |
| Family members, mean (SD)              | 4.31 (1.92)       | 4.55 (1.70)       | .39 <sup>a</sup> |
| Monthly income (US \$), median (IQR)   | 182 (136, 255)    | 182 (168, 273)    | .05 <sup>c</sup> |
| <b>Households have own, n (%)</b>      |                   |                   |                  |
| Television                             | 0 (0)             | 1 (1.2)           | .32 <sup>b</sup> |
| Mobile phone                           | 27 (32.1)         | 37 (44)           | .11 <sup>b</sup> |
| Refrigerator                           | 37 (44)           | 46 (54.8)         | .17 <sup>b</sup> |
| Electric fan                           | 82 (97.6)         | 83 (98.8)         | .56 <sup>b</sup> |
| Current smoker, n (%)                  | 31 (36.90)        | 22 (26.19)        | .14 <sup>b</sup> |
| Alcohol consumption, n (%)             | 10 (11.9)         | 3 (3.57)          | .04 <sup>b</sup> |
| BMI (kg/m <sup>2</sup> ), median (IQR) | 22.9 (19.5, 25.6) | 23.6 (21.4, 26.4) | .66 <sup>c</sup> |

<sup>a</sup> Independent sample *t* test

<sup>b</sup> Chi-square test

<sup>c</sup> Mann-Whitney *U* test

### Dissemination:

The study results will be disseminated to the Director General of Health and Family Services, the Department of Environment (DoE), and partnering civil society organizations in Bangladesh that are dedicated to addressing health and environmental issues, aiming to uphold environmental safety. Findings will be used to explore the boundaries and challenges of plastic recycling and management in Bangladesh. Moreover, strategies at both technical and policy levels will be formulated alongside efforts to enhance public awareness of the environmental and health risks linked with plastic recycling. We will also develop initiatives to build capacity in this area. Abstracts will be prepared for international conferences to disseminate our findings globally, and manuscripts will be drafted and submitted to peer-reviewed journals for publication.

### Discussion:

Several studies have examined the issue of plastic pollution in Bangladesh, with a primary focus on marine and aquatic environments. However, no study has investigated the potential human health impacts resulting from microplastic exposure [45]. Earlier evidence indicates the presence of MP in fish [20], and shrimp [46] from the northern Bay of Bengal. Additionally, microplastics have been detected in sediments from the sea beaches and major rivers of Bangladesh [47]. While exploring the sources of these microplastics, it was revealed that the plastic recycling industry also contributes due to the mechanical process and inadequate effluent treatments [48]. The presence of microplastic particles in our surrounding environment is evident, yet numerous unexplored aspects remain in measuring exposure and assessing health outcomes. While most of

the health impacts have been hypothesized based on animal studies, only a few have been investigated through epidemiological research in humans. The proposed study aims to contribute evidence on human microplastic exposure and possible health impacts suggested by animal studies. In 2022, researchers from the Netherlands reported the first-ever detection of plastic particles in human blood, with 77% of the tested blood samples containing such particles [49]. Instead of directly detecting microplastic from biological samples, this study will focus on the metabolites of bisphenol and phthalates [50]. Moreover, this study is designed to generate evidence on the presence of microplastic particles in environmental media, the extent of human exposure, and associated health outcomes. Collecting and analyzing a wide range of environmental sample types will contribute to a more comprehensive understanding of microplastic distribution in the environment and potential routes of human exposure. The results from this study will serve as a foundation for further research into monitoring the long-term ecological impacts of plastic debris. The multifaceted approach will also enhance the ability to develop targeted strategies for future mitigation and regulatory measures.

Nonetheless, this study has some limitations. We intend to identify the presence of bisphenol and phthalate metabolites in biological samples, representing an indirect method of assessing exposure. To address this limitation, we plan to collect environmental samples from the respondents' workplaces to detect the presence of microplastic particles, which may validate our exposure assessment. This study is cross-sectional, from which causality between exposures and outcomes cannot be inferred.

**Conclusion:**

The findings from this study could be valuable for regulatory authorities in understanding the magnitude of the problem. The proposed analysis of multiple systems, including hormonal, renal, lung function, and oxidative stress indicators, may enhance our understanding of the mechanism between exposures and biological responses. Additionally, the detailed environmental investigation would provide initial evidence of contamination in the plastic industry, which might lead to negative consequences for the environment and human health. These findings could also be crucial in implementing occupational safety measures to mitigate adverse impacts, enabling informed decisions in formulating effective waste management and resource utilization strategies.

## Author affiliation

<sup>1</sup>Environmental Health and WASH, Health System and Population Studies Division, icddr,b, Dhaka 1212, Bangladesh

<sup>2</sup>Department of Chemistry, University of Dhaka, Dhaka 1000, Bangladesh

<sup>3</sup>Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka 1205, Bangladesh

<sup>4</sup>Department of Environmental & Sustainable Engineering, University at Albany, State University of New York, Albany, NY 12222, United States

<sup>5</sup>Department of Civil and Environmental Engineering, University of Nebraska–Lincoln, W150E Nebraska Hall, 900 N 16th St Lincoln, NE 68588-0531

<sup>6</sup>Nutrition Research Division, icddr,b, Dhaka 1212, Bangladesh

<sup>7</sup>Children's Health and Environment Program, Child Health Research Centre, The University of Queensland, South Brisbane, QLD, Australia

## Acknowledgments

The author acknowledges the study collaborators and the cooperation of the study participants, who were visited several times for data collection. This research study was funded by the 'Integrated Health Science Research and Development Fund Activity' from the Directorate General of Medical Education Branch, Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh. We also acknowledge our core donors for their support and commitment to icddr,b's research efforts.

## Contributors

SSH and SMP developed the study concept. SMP, SSH, AS, and FJ developed the study design. AS, AB, NI, MR, ZI, MHK, RR, and MM provided input in environmental and biological sample collection and analysis. SSH and SMP provided input in statistical analysis. SSH drafted the paper, and all authors read and commented on it and agreed with the final version.

## Data Availability

Data will be available upon request.

## **Conflicts of interests**

None declared.



**References:**

1. Fok L, Cheng INY, Yeung YY. Mismanaged Plastic Waste: Far Side of the Moon. In: So WWM, Chow CF, Lee JCK, editors. *Environmental Sustainability and Education for Waste Management: Implications for Policy and Practice*. Singapore: Springer Singapore; 2019. p. 57-71.
2. Rochman CM, Browne MA, Halpern BS, Hentschel BT, Hoh E, Karapanagioti HK, et al. Classify plastic waste as hazardous. *Nature*. 2013 2013/02/01;494(7436):169-71. doi: 10.1038/494169a.
3. Geyer R, Jambeck JR, Law KL. Production, use, and fate of all plastics ever made. *Science advances*. 2017 Jul;3(7):e1700782. PMID: 28776036. doi: 10.1126/sciadv.1700782.
4. Waste Concern. National workshop on sustainable management of plastic to leverage circular economy and achieve SDG in Bangladesh. Waste Concern; 2019 [cited 2021 23 June]; Available from: <https://wasteconcern.org/national-workshop-on-sustainable-management-of-plastic-to-leverage-circular-economy-and-achieve-sdg-in-bangladesh/>.
5. Jambeck JR, Geyer R, Wilcox C, Siegler TR, Perryman M, Andrady A, et al. Plastic waste inputs from land into the ocean. *Science*. 2015;347(6223):768-71. doi: doi:10.1126/science.1260352.
6. DeWit W, Burns ET, Guinchard J-C, Ahmed N. *Plastics: the costs to society, the environment and the economy*. Gland, Switzerland: World Wide Fund for Nature (WWF), 2021.
7. Barnes DK, Galgani F, Thompson RC, Barlaz M. Accumulation and fragmentation of plastic debris in global environments. *Philos Trans R Soc Lond B Biol Sci*. 2009 Jul 27;364(1526):1985-98. PMID: 19528051. doi: 10.1098/rstb.2008.0205.
8. Enyoh CE, Verla AW, Verla EN, Ibe FC, Amaobi CE. Airborne microplastics: a review study on method for analysis, occurrence, movement and risks. *Environmental Monitoring and Assessment*. 2019 2019/10/24;191(11):668. doi: 10.1007/s10661-019-7842-0.
9. Campanale C, Massarelli C, Savino I, Locaputo V, Uricchio VF. A Detailed Review Study on Potential Effects of Microplastics and Additives of Concern on Human Health. *International journal of environmental research and public health*. 2020 Feb 13;17(4). PMID: 32069998. doi: 10.3390/ijerph17041212.
10. Cox KD, Covernton GA, Davies HL, Dower JF, Juanes F, Dudas SE. Human Consumption of Microplastics. *Environmental Science & Technology*. 2019 2019/06/18;53(12):7068-74. doi: 10.1021/acs.est.9b01517.
11. Ebere EC, Wirnkor VA, Ngozi VEJWSN. Uptake of Microplastics by Plant: a Reason to Worry or to be Happy? 2019;131:256-67.
12. Flaws J, Damdimopoulou P, Patisaul HB, Gore A, Raetzman L, Vandenberg LN. *Plastics, EDCs & Health: Authoritative Guide*. Endocrine Society, 2020 December 10. Report No.
13. Yee MS, Hii LW, Looi CK, Lim WM, Wong SF, Kok YY, et al. Impact of Microplastics and Nanoplastics on Human Health. *Nanomaterials (Basel)*. 2021 Feb 16;11(2). PMID: 33669327. doi: 10.3390/nano11020496.
14. Vethaak AD, Legler J. Microplastics and human health. *Science*. 2021;371(6530):672-4.

doi: doi:10.1126/science.abe5041.

15. Hahladakis JN, Velis CA, Weber R, Iacovidou E, Purnell P. An overview of chemical additives present in plastics: Migration, release, fate and environmental impact during their use, disposal and recycling. *Journal of Hazardous Materials*. 2018 2018/02/15/;344:179-99. doi: <https://doi.org/10.1016/j.jhazmat.2017.10.014>.
16. Parvez SM, Jahan F, Abedin J, Rahman M, Hasan SS, Islam N, et al. Blood lead, cadmium and hair mercury concentrations and association with soil, dust and occupational factors in e-waste recycling workers in Bangladesh. *International Journal of Hygiene and Environmental Health*. 2024 2024/04/01/;257:114340. doi: <https://doi.org/10.1016/j.ijheh.2024.114340>.
17. Parvez SM, Jahan F, Brune MN, Gorman JF, Rahman MJ, Carpenter D, et al. Health consequences of exposure to e-waste: an updated systematic review. *Lancet Planet Health*. 2021 Dec;5(12):e905-e20. PMID: 34895498. doi: 10.1016/S2542-5196(21)00263-1.
18. Parvez SM, Hasan SS, Moniruzzaman M, Jahan F, Rahman M, Aich N, et al. Trace metal concentrations, geoaccumulation index and risk assessment from soil of electronic waste (e-waste) exposed and non-exposed area, Bangladesh. *ISEE Conference Abstracts*.2023(1). doi: 10.1289/isee.2023.MP-028.
19. Sultan MB, Rahman MM, Khatun MA, Shahjalal M, Akbor MA, Siddique MAB, et al. Microplastics in different fish and shellfish species in the mangrove estuary of Bangladesh and evaluation of human exposure. *Science of The Total Environment*. 2023 2023/02/01/;858:159754. doi: <https://doi.org/10.1016/j.scitotenv.2022.159754>.
20. Hossain MS, Sobhan F, Uddin MN, Sharifuzzaman SM, Chowdhury SR, Sarker S, et al. Microplastics in fishes from the Northern Bay of Bengal. *Sci Total Environ*. 2019 Nov 10;690:821-30. PMID: 31302547. doi: 10.1016/j.scitotenv.2019.07.065.
21. Rahman SMA, Robin GS, Momotaj M, Uddin J, Siddique MAM. Occurrence and spatial distribution of microplastics in beach sediments of Cox's Bazar, Bangladesh. *Mar Pollut Bull*. 2020 Nov;160:111587. PMID: 32871432. doi: 10.1016/j.marpolbul.2020.111587.
22. Brown E, MacDonald A, Allen S, Allen D. The potential for a plastic recycling facility to release microplastic pollution and possible filtration remediation effectiveness. *Journal of Hazardous Materials Advances*. 2023 2023/05/01/;10:100309. doi: <https://doi.org/10.1016/j.hazadv.2023.100309>.
23. Zarus GM, Muianga C, Hunter CM, Pappas RS. A review of data for quantifying human exposures to micro and nanoplastics and potential health risks. *Sci Total Environ*. 2021 Feb 20;756:144010. PMID: 33310215. doi: 10.1016/j.scitotenv.2020.144010.
24. Young VB. The role of the microbiome in human health and disease: an introduction for clinicians. *BMJ*. 2017;356:j831. doi: 10.1136/bmj.j831.
25. Murphy RA. Prevalence of Inappropriate Antibiotic Prescriptions Among US Ambulatory Care Visits, 2010-2011. *Journal of Emergency Medicine*. 2016;51(3):337. doi: 10.1016/j.jemermed.2016.07.019.
26. Kouidhi W, Thannimalay L, Soon CS, Ali Mohd M. Occupational exposure to bisphenol A (BPA) in a plastic injection molding factory in Malaysia. *Int J Occup Med Environ Health*.

2017 Jul 14;30(5):743-50. PMID: 28584331. doi: 10.13075/ijomeh.1896.00917.

27. Pilka T, Petrovicova I, Kolena B, Zatko T, Trnovec T. Relationship between variation of seasonal temperature and extent of occupational exposure to phthalates. *Environ Sci Pollut Res Int*. 2015 Jan;22(1):434-40. PMID: 25081008. doi: 10.1007/s11356-014-3385-7.

28. Lebreton L, Andrady A. Future scenarios of global plastic waste generation and disposal. *Palgrave Communications*. 2019 2019/01/29;5(1):6. doi: 10.1057/s41599-018-0212-7.

29. United Nations Environment Programme. National guidance for plastic pollution hotspotting and shaping action - Introduction to the methodology. Nairobi, Kenya: United Nations Environment Programme,; 2020. ISBN: 978-92-807-3793-6.

30. Saunders B, Sim J, Kingstone T, Baker S, Waterfield J, Bartlam B, et al. Saturation in qualitative research: exploring its conceptualization and operationalization. *Quality & Quantity*. 2018 2018/07/01;52(4):1893-907. doi: 10.1007/s11135-017-0574-8.

31. Revel M, Châtel A, Mouneyrac C. Micro(nano)plastics: A threat to human health? *Current Opinion in Environmental Science & Health*. 2018 2018/02/01;1:17-23. doi: <https://doi.org/10.1016/j.coesh.2017.10.003>.

32. Wright SL, Kelly FJ. Plastic and Human Health: A Micro Issue? *Environmental Science & Technology*. 2017 2017/06/20;51(12):6634-47. doi: 10.1021/acs.est.7b00423.

33. Schirizzi GF, Perez-Pomeda I, Sanchis J, Rossini C, Farre M, Barcelo D. Cytotoxic effects of commonly used nanomaterials and microplastics on cerebral and epithelial human cells. *Environ Res*. 2017 Nov;159:579-87. PMID: 28898803. doi: 10.1016/j.envres.2017.08.043.

34. Anbumani S, Kakkar P. Ecotoxicological effects of microplastics on biota: a review. *Environ Sci Pollut Res Int*. 2018 May;25(15):14373-96. PMID: 29680884. doi: 10.1007/s11356-018-1999-x.

35. Galloway TS. Micro- and Nano-plastics and Human Health. Bergmann M, Gutow L, Klages M, editors. Cham: Springer International Publishing; 2015 2015/. 343-66 p. ISBN: 978-3-319-16510-3.

36. Fiorentino I, Gualtieri R, Barbato V, Mollo V, Braun S, Angrisani A, et al. Energy independent uptake and release of polystyrene nanoparticles in primary mammalian cell cultures. *Exp Cell Res*. 2015 Jan 15;330(2):240-7. PMID: 25246129. doi: 10.1016/j.yexcr.2014.09.017.

37. Rahman A, Sarkar A, Yadav OP, Achari G, Slobodnik J. Potential human health risks due to environmental exposure to nano- and microplastics and knowledge gaps: A scoping review. *Sci Total Environ*. 2021 Feb 25;757:143872. PMID: 33310568. doi: 10.1016/j.scitotenv.2020.143872.

38. Rebai I, Fernandes JO, Azzouz M, Benmohammed K, Bader G, Benmbarek K, et al. Urinary bisphenol levels in plastic industry workers. *Environ Res*. 2021 Nov;202:111666. PMID: 34265347. doi: 10.1016/j.envres.2021.111666.

39. Huang Y, Chapman J, Deng Y, Cozzolino D. Rapid measurement of microplastic contamination in chicken meat by mid infrared spectroscopy and chemometrics: A feasibility study. *Food Control*. 2020 2020/07/01;113:107187. doi: 10.1016/j.foodcont.2020.107187.

40. Franke TM, Ho T, Christie CA. The Chi-Square Test: Often Used and More Often



- Misinterpreted. *American Journal of Evaluation*. 2012;33(3):448-58. doi: 10.1177/1098214011426594.
41. Kim TK, Park JH. More about the basic assumptions of t-test: normality and sample size. *kja*. 2019 04;72(4):331-5. doi: 10.4097/kja.d.18.00292.
  42. Ostertagová E, Ostertag O, Kováč J. Methodology and Application of the Kruskal-Wallis Test. *Applied Mechanics and Materials*. 2014;611:115-20. doi: 10.4028/[www.scientific.net/AMM.611.115](http://www.scientific.net/AMM.611.115).
  43. Naeem M, Ozuem W, Howell K, Ranfagni S. A Step-by-Step Process of Thematic Analysis to Develop a Conceptual Model in Qualitative Research. *International Journal of Qualitative Methods*. 2023;22:16094069231205789. doi: 10.1177/16094069231205789.
  44. Campbell R, Goodman-Williams R, Feeney H, Fehler-Cabral G. Assessing Triangulation Across Methodologies, Methods, and Stakeholder Groups: The Joys, Woes, and Politics of Interpreting Convergent and Divergent Data. *American Journal of Evaluation*. 2020;41(1):125-44. doi: 10.1177/1098214018804195.
  45. Islam T, Li Y, Rob MM, Cheng H. Microplastic pollution in Bangladesh: Research and management needs. *Environmental Pollution*. 2022 2022/09/01/;308:119697. doi: <https://doi.org/10.1016/j.envpol.2022.119697>.
  46. Hossain MS, Rahman MS, Uddin MN, Sharifuzzaman SM, Chowdhury SR, Sarker S, et al. Microplastic contamination in Penaeid shrimp from the Northern Bay of Bengal. *Chemosphere*. 2020 2020/01/01/;238:124688. doi: 10.1016/j.chemosphere.2019.124688.
  47. Hossain MB, Banik P, Nur A-AU, Rahman T. Abundance and characteristics of microplastics in sediments from the world's longest natural beach, Cox's Bazar, Bangladesh. *Marine Pollution Bulletin*. 2021 2021/02/01/;163:111956. doi: <https://doi.org/10.1016/j.marpolbul.2020.111956>.
  48. Suzuki G, Uchida N, Tuyen LH, Tanaka K, Matsukami H, Kunisue T, et al. Mechanical recycling of plastic waste as a point source of microplastic pollution. *Environmental Pollution*. 2022 2022/06/15/;303:119114. doi: <https://doi.org/10.1016/j.envpol.2022.119114>.
  49. Leslie HA, van Velzen MJM, Brandsma SH, Vethaak AD, Garcia-Vallejo JJ, Lamoree MH. Discovery and quantification of plastic particle pollution in human blood. *Environ Int*. 2022 May;163:107199. PMID: 35367073. doi: 10.1016/j.envint.2022.107199.
  50. Thompson RC, Moore CJ, vom Saal FS, Swan SH. Plastics, the environment and human health: current consensus and future trends. *Philos Trans R Soc Lond B Biol Sci*. 2009 Jul 27;364(1526):2153-66. PMID: 19528062. doi: 10.1098/rstb.2009.0053.

## Abbreviations:

|                        |  |
|------------------------|--|
| <b>DoE</b>             | : Department of Environment              |
| <b>ERC</b>             | : Ethical review committee               |
| <b>EPA</b>             | : Environmental Protection Agency        |
| <b>ELISA</b>           | : Enzyme-linked immunosorbent assay      |
| <b>eGFR</b>            | : estimated Glomerular filtration rate   |
| <b>FEV<sub>1</sub></b> | : Forced expiratory volume in one second |

|                 |  |
|-----------------|--|
| <b>FTIR</b>     | : Fourier-transform infrared spectroscopy        |
| <b>FVC</b>      | : Forced vital capacity                          |
| <b>HM</b>       | : Heavy metal                                    |
| <b>ICP-MS</b>   | : Inductively coupled plasma mass spectrometry   |
| <b>IQR</b>      | : Interquartile range                            |
| <b>LC-MS/MS</b> | : Liquid Chromatography-Tandem Mass Spectrometry |
| <b>LH</b>       | : Luteinizing hormone                            |
| <b>MiNP</b>     | : Mono isononyl phthalate                        |
| <b>MP</b>       | : Microplastic particles                         |
| <b>PTB</b>      | : Preterm birth                                  |
| <b>PEFR</b>     | : Peak expiratory flow rate                      |
| <b>pPROM</b>    | : Preterm premature rupture of membrane          |
| <b>RM</b>       | : Recurrent miscarriage                          |
| <b>RRC</b>      | : Research review committee                      |
| <b>SD</b>       | : Standard deviation                             |
| <b>TSP</b>      | : Total suspended particulate                    |
| <b>TT</b>       | : Total testosterone                             |

## Supplementary Files

## Multimedia Appendixes

Comments from the research review committee.

URL: <http://asset.jmir.pub/assets/9ac72c4757769e9ec43ca81d5235d274.pdf>

Approval of the research review committee.

URL: <http://asset.jmir.pub/assets/f843dfe8d1de6f9c43aee64661d930d6.pdf>

Comments from the ethical review committee.

URL: <http://asset.jmir.pub/assets/12021a172a93b773aa64159425e9ae3d.pdf>

Approval from the ethical review committee.

URL: <http://asset.jmir.pub/assets/55c67860d0ff9e28831e38c615d04538.pdf>