

Veterans Affairs Explores Contactless Vital Signs Collection in Video Telehealth Visits: Usability Pilot among Providers and Patients

Lynn Garvin, Eric Richardson, Leonie Heyworth, D. Keith McInnes

Submitted to: JMIR Formative Research
on: May 13, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 22

 Figures 23

 Figure 1..... 24

 Figure 2..... 25

 Figure 3..... 26

 Multimedia Appendixes 27

 Multimedia Appendix 1..... 28

 Multimedia Appendix 2..... 28

 Multimedia Appendix 3..... 28

 Multimedia Appendix 4..... 28

Veterans Affairs Explores Contactless Vital Signs Collection in Video Telehealth Visits: Usability Pilot among Providers and Patients

Lynn Garvin^{1, 2*} MBA, PhD; Eric Richardson^{1*} PhD; Leonie Heyworth^{3, 4*} MPH, MD; D. Keith McInnes^{2, 5*} MS, ScD

¹Center for Healthcare Organization and Implementation Research Veterans Affairs Boston Healthcare System Boston US

²Department of Health Law, Policy and Management Boston University School of Public Health Boston US

³Office of Connected Care/Telehealth US Department of Veterans Affairs Central Office Washington US

⁴Department of Medicine University of California, San Diego School of Medicine San Diego US

⁵Center for Healthcare Organization and Implementation Research VA Bedford Healthcare System Bedford US

*these authors contributed equally

Corresponding Author:

Lynn Garvin MBA, PhD

Center for Healthcare Organization and Implementation Research

Veterans Affairs Boston Healthcare System

Boston

US

Abstract

Background: To expand Veterans' access to healthcare, the Veterans Affairs (VA) Office of Connected Care explored a novel software feature called "Vitals" on its VA Video Connect telehealth platform. Vitals uses contactless video-based remote photoplethysmography (rPPG) through the infrared camera on Veterans' smartphones (and other devices) to automatically scan their faces to provide real-time vital statistics on screen to both the provider and patient.

Objective: The purpose of this project was to assess VA clinical provider and Veteran patient attitudes regarding the usability of Vitals.

Methods: We conducted a mixed methods evaluation of Vitals among VA providers and patients, collecting data in July and August 2023 at the VA Boston Healthcare System and VA San Diego Healthcare System. We conducted analyses in October 2023. In-person usability testing sessions consisted of a "think aloud" procedure while using the software, a semi-structured interview and a 26-item online survey.

Results: Usability test sessions with 20 VA providers and 13 patients demonstrated that both groups found Vitals "useful" and "easy to use," and rated its usability highly (86 and 82 points, respectively, on a 100-point scale). Regarding acceptability or willingness/intent to use, providers and patients generally expressed confidence and trust in Vitals readings, with high ratings of 90 and 85 points, respectively. Providers and patients rated Vitals highly for its feasibility and appropriateness for context (90 and 90 points, respectively on both measures). Finally, providers noted that Vitals' flexibility makes it appropriate and advantageous for implementation in a wide range of clinical contexts, particularly in specialty care. Providers believed that most clinical teams would readily integrate Vitals into their routine workflow because it saves time; delivers accurate, consistently collected vitals; and may reduce reporting errors. Providers and Veterans suggested training and support materials that could improve Vitals adoption and implementation.

Conclusions: While remote collection of vital readings has been described in the literature, this is one of the first accounts of testing a contactless vitals tool among providers and patients. If ongoing initiatives demonstrate accuracy in its readings, Vitals could enhance virtual care by: providing accurate, automatic reporting and recording of vitals, sending patients' vital readings (pending provider approval) directly to their electronic medical record; saving provider and patient time; and potentially reducing necessity of some home-based biometric devices. Understanding these issues prior to FDA approval of Vitals and its implementation could contribute to a seamless introduction of Vitals to VA providers and patients.

(JMIR Preprints 13/05/2024:60491)

DOI: <https://doi.org/10.2196/preprints.60491>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/preprint/60491>

Original Manuscript

Veterans Affairs Explores Contactless Vital Signs Collection in Video Telehealth Visits: Usability Pilot among Providers and Patients

Lynn A. Garvin, Eric Richardson, Leonie Heyworth, D. Keith McInnes


Introduction

Veterans' access to high quality healthcare is a priority for the US Department of Veterans Affairs (VA) which serves over 9 million enrolled Veterans each year.¹ VA has built its telehealth services over the past 20 years, with 40% of Veterans now receiving some portion of their VA care through telehealth.² During this time, VA and U.S. policy makers have sought to address the digital divide faced by many Veterans due to demographic, clinical and geographic challenges (e.g., travel distance to care in rural settings).^{3,4} In 2016, VA's Office of Rural Health and Office of Connected Care (OCC) launched a national initiative to distribute video-enabled tablets to Veterans facing access barriers.^{5,6} The MISSION Act of 2018 eliminated geographic licensure constraints, permitting VA providers to care for Veterans across state lines in the U.S., its possessions and territories.⁷ In 2019, VA partnered with mobile carriers to reduce data fees for Veterans using VA video telehealth.⁸ To meet the sudden demand for virtual care posed by the COVID-19 pandemic, in 2020 VA's video architecture was rapidly scaled to allow over 3000% growth in video telehealth in one year.⁶ Simultaneously, OCC launched its Digital Divide consult which sustained tablet distribution to eligible Veterans⁹ and, in 2021, partnered with the U.S. Department of Housing and Urban Development-VA Supportive Housing (HUD-VASH) program to distribute smartphones and tablets to homeless Veterans to support their virtual care access.¹⁰ Current expansion of VA video telehealth continues to focus on quality and equity.^{11,12}

One critical virtual care solution is the VA Video Connect (VVC) secure videoconferencing app, designed to help Veterans meet with their healthcare teams on any smartphone, computer or tablet.¹³ Studies have shown that video telehealth can offer effective delivery of mental healthcare,¹⁴⁻¹⁶ primary care,^{17,18} and specialty ambulatory care.^{19,20} Patient populations who face demographic, clinical and geographic challenges (e.g., travel distance to care in rural settings) can benefit from video telehealth.^{3,4}

The VA Office of Connected Care which manages VVC is constantly exploring new innovations. A novel VVC software feature called "Vitals" uses contactless video-based remote photoplethysmography (rPPG) technology through the infrared camera on Veterans' smartphones (and other devices) to automatically scan their faces when incorporated into the video platform. (Figure 1) Within 45 seconds, it delivers vital statistics on screen to both the provider and patient. Vitals' statistics include: blood pressure, heart rate, respiratory

Figure 1. rPPG technology on smartphones can scan patient faces to deliver vital statistics for patient and provider view



rate, pulse and temperature. U.S. Food and Drug Administration (FDA) approval for Vitals software is pending. If Vitals demonstrates accuracy in its readings (as tested through a separate project), it could enhance virtual care by: providing accurate, automatic reporting and recording of vitals, sending patients' vital readings (pending provider approval) directly to their electronic medical record; saving provider and patient time; and potentially reducing necessity of some home-based biometric devices. There are a number of current VA initiatives in which remote vital sign collection would be beneficial, such as in pre-anesthesia patient evaluations.²¹ And while remote collection of vital readings has been described in the literature,²² this is one of the first accounts of testing a contactless vitals tool among providers and patients.

Hence, the objective of this project was to evaluate VA provider and Veteran attitudes regarding the usability of, and intent to use, Vitals. Related attitudes of trust, privacy, credibility, personal preference and practice were explored. Providers and Veterans (hereafter "patients") might voice a range of interests, questions and/or concerns regarding Vitals and its potential to support accurate and transparent health information collection, data sharing and use. Understanding these issues prior to FDA approval and implementation could contribute to a seamless introduction of Vitals to VA providers and patients.

Methods

Overview

We conducted a mixed methods evaluation of VA provider and patient attitudes regarding the usability of Vitals and their intent to use. In-person usability testing sessions consisted of a "think aloud" procedure and a semi-structured interview, followed by a quantitative survey. All procedures were approved and considered quality improvement by the Institutional Review Board at the VA Boston Healthcare System in March 2023. We collected data in July and August 2023 and conducted analysis in October 2023.

Participants

We recruited participants from the primary and specialty care programs of the VA Boston Healthcare System and VA San Diego Healthcare System. A prospective sample of 15 providers (7-8 per site) included 5 primary care and 10 specialty care providers. Based on studies of specialty care telemedicine^{19,20,23} and guidance from OCC, 10 specialty care providers were selected from across specialties that could most benefit from Vitals: dermatology, endocrinology (e.g. Diabetes -T1 & T2), oncology, gastroenterology, pulmonology (e.g. chronic obstructive pulmonary disease (COPD)), nephrology (e.g. chronic kidney disease), cardiology (e.g. congestive heart failure), neurology (e.g. functional neurological disorder), anesthesiology, pain management, rehabilitation (e.g. physical or speech therapy), and emergency/urgent care. A prospective sample of 20 patients (10 per site) included 10 "Less Experienced" patients who had had 1-2 VVC visits, and 10 "More Experienced" patients who had had 3+ VVC visits within the past year on tablets (no phones or computers). The sample reflected a variety of health conditions relevant to Vitals statistics, e.g., diabetes or COPD, and provided a broad sociodemographic representation. Based on qualitative research standards, "data saturation" is the point at which no new information or themes are observed. We determined that saturation could be reached with 5-10 individual interviews per sample.^{24,25} All participants provided verbal informed consent for their participation and audio recording.


Usability Testing among Providers and Patients

We conducted 30-minute in-person sessions with providers, and 60-minute in-person sessions with patients. Two researchers (LG and ER) conducted sessions in Boston and one researcher (DM) conducted sessions in San Diego. Sessions consisted of two steps: 1) usability testing through a think aloud protocol and interview, and 2) a structured survey.

Step 1: Usability testing and semi-structured interviews

Usability testing combined two qualitative methods: a “think aloud” procedure and an interview. In a think aloud exercise, the participant is asked to voice his/her thoughts while examining or navigating the use of a technical device or software.²⁶⁻²⁸ To make the think aloud experience as real as possible, we conducted mock VVC visits among patients with the researcher playing the role of “the provider” collecting the patient’s vital signs using Vitals. Patients were given Vitals-enabled smartphones while the researcher acting as “the provider” used a Vitals-enabled clinical tablet. Alternatively, the researcher played the role of “the patient” with VA providers. This procedure enabled us to gather participants’ real-time impressions while observing their facial cues and body language while they were using Vitals. Static visuals (storyboards, see Figure 2) of the providers’ and patients’ views of Vitals on a tablet served as a backup tool for interviews in case issues arose that prevented the tablet, VVC or Vitals from operating.

Figure 2. Views of Vitals on Veterans’ smartphones and on VA Providers’ tablets (The purple arrows point to the Vital sign readings on both devices)



The semi-structured interviews that followed documented participants’ attitudes about Vitals’ based on constructs of the Unified Theory of Acceptance and Use of Technology (UTAUT)²⁹ constructs and other themes (detailed in “Measurement and instrument”).

Step 2: Quantitative survey

We concluded our session with the collection of provider and patient responses to a brief quantitative survey composed of validated scales for intervention usability, acceptability (willingness to use), appropriateness (fit for purpose), and feasibility. Researchers read survey questions aloud to patients and entered responses to the online survey hosted on VA’s secure Research Electronic Data Capture (REDCap) platform.³⁰ Patient participants received a \$25 CVS gift card as reimbursement for completing the session. Being familiar with REDCap, providers entered their own responses to the online survey.

Measurement and instrument

The semi-structured interview guides for providers and patients were informed by the UTAUT model which proposes that performance expectancy, effort expectancy, social influence and facilitating conditions (positive and negative) influence a person's intent to use, the strongest predictor of technology adoption.²⁹ Other themes explored included participants' attitudes on Vitals' measurement accuracy, data privacy, system security, and their trust in the VVC app. Interview questions asked whether any training, educational materials or technical support were needed; and probed for barriers and facilitators to Vitals' acceptance and sustained use (e.g. impact on clinical workflow). Table 1 summarizes the constructs, measures, and their sources. (See Multimedia Appendices 1-4 for Interview Guides and Online Surveys for providers and patients.)

Table 1. Constructs, measures, and sources: Usability evaluation

Constructs	Measures
(Step 1a) Think aloud protocol	
User perceptions	Unfiltered user impressions of the usability and acceptability of Vitals tool.
(Step 1b) Qualitative measures – Patient and provider usability testing interviews based on UTAUT constructs	
Performance expectancy	Perceived effectiveness/usefulness, e.g., provider time saved in recording accurate blood pressure reading; Equivalent care quality for VVC visit as an in-person visit.
Effort expectancy	Design simplicity/ease of use, e.g., patient convenience of not having to travel for care; upon provider approval, Vitals automatically adds the reading to patient health record.
Social influence	Community and provider support for use, e.g., availability of technology and training.
Facilitating conditions	Facilitators or barriers to technology use, e.g., friends who assist with technology is a positive facilitating condition, while depression can be a negative facilitating condition.
(Step 2) Quantitative Measures – Survey administered at the conclusion of patient and provider interviews. All measures use Likert-style questions on a 5-point continuum	
Usability	Factors that affect <u>providers' and patients' ability</u> to use the solution to achieve specified goals. The 10-item Intervention Usability Scale (IUS), ³¹ appropriate for combined interpersonal and technology solutions, is an adaptation of the widely used System Usability Scale. ^{32,33}
Acceptability	Factors that affect <u>providers' and patients' willingness/intent to use</u> a solution deemed enjoyable/comfortable to use. 4-item Acceptability of Intervention Measure (AIM). ³⁴
Feasibility	Motivation and ability to introduce and support the solution, and the extent to which the <u>solution is practical and can be successfully used or completed</u> . Feasibility will be assessed with the 4-item Feasibility of Intervention Measure (FIM). ³⁴
Appropriateness	Factors that affect the <u>perceived fit (relevance, suitability) of the solution</u> to a given context, a given provider and/or patient, or solving a particular issue or problem. 4-item Intervention Appropriateness Measure (IAM). ³⁴

The 26-item survey quantitative survey that concluded each session was composed of four validated scales: the Intervention Usability Scale (IUS),³¹ the Acceptability of Intervention Measure (AIM),³⁴ the Feasibility of Intervention Measure (FIM),³⁴ and the Intervention Appropriateness Measure (IAM).³⁴ All measures use Likert-style questions on a 5-point continuum. Based on research on the System Usability Scale^{32,33} (the parent scale for IUS), a score above a 68 Points is considered above average and anything below 68 Points is below average.^{32,35}

Data Analysis

For qualitative analysis, audio recordings of usability testing sessions were transcribed. These, along with session notes, were coded thematically using rapid qualitative inquiry method.^{36,37} Two researchers (LG and ER) coded interviews, performing inductive and deductive content analysis. A priori categories included UTAUT constructs (performance expectancy, effort expectancy, social influence and facilitating conditions) and other themes from the literature (see "Measurement

Instruments”). Emergent themes from participants’ think aloud comments generated additional codes. The research team conducted a thematic analysis that assessed the patterns of attitudes and experiences and reached consensus on interpretation.

For quantitative analysis, we calculated total scores for the survey measures following the published guidelines for scoring. For ease of interpretation, scores for the AIM, FIM, and IAM were converted to a 100-point scale to match the IUS scores. We then calculated descriptive statistics (e.g., range and mean) of the total scores on measures. We calculated descriptive statistics of individual items to capture some additional insights into participant responses. Analysis was conducted using R statistical software version 4.2.³⁸

Results

Sample characteristics

We interviewed 20 VA providers whose average age was 47 years (Range=30-63). Over half (60%) identified as female. Equal numbers identified their race as Asian (35%) or White (35%). Service line representation was 45% primary care, and 25% specialty care, with remaining indicating a telehealth specific position. Specialty care included urgent care, neurology physical therapy and surgery. Their experience with telehealth was broadly distributed.




Figure 3. VA Provider and Veteran average score (out of 100 points) for Vitals on scales for usability, acceptability, feasibility and appropriateness to context

We also interviewed 13 patients with an average age of 67 (Range=38-84) all of whom identified as male. Race/ethnicity was evenly divided among patients who self-identified as either: Black/African

American, Hispanic/Latinx or Asian. All patients had attained at least a high school diploma. Most patients (90%) lived 30 to 60 minutes from their VA facility.

Usability of Vitals

As seen in Figure 3, providers and patients highly rated Vitals' usability (86 Points and 82 Points, respectively). Overall, providers thought that Vitals was "useful" and "easy to use."

I like it. That's my biggest thing with doing VVCs is sometimes not having vitals...One of the biggest things I see is hypertension, and the blood pressure is always elevated. And so that's like the biggest thing I would want to see is the blood pressure... (Provider 202, NP)

I think -- it seems pretty easy.. I think having batteries and making sure you find a finger if they have... circulation issues, that this [Vitals] is almost easier than using the pulse oximeter. (Provider 105, RN)

Providers liked that Vitals "helps to provide some valuable context" and "helps to triage" patient's general health and particular conditions.

I do actually like the stress level. That's something we don't do in person. We ask Veterans manually about what their stress levels are...So, more of an objective look at stress levels...So, that's one I think that's in here that I don't think we get normally and I would like. (Provider 211, MD)

Patients found Vitals both easy to use and more convenient than conventional methods of reading Vitals remotely, e.g., using a blood pressure (BP) cuff or oxygen (O2) sensor.

It's a lot easier to sit still and relax in front of a computer screen monitor than it is holding your phone. (Veteran 201)

Well, right now you have to do a BP cuff and an O2 sensor, my scale, and then my finger for my glucose to get all the readings to provide... (Veteran 201)

While Vitals usability scored high overall, providers rated two items on the IUS slightly lower, indicating the need for technical assistance at the introduction of the Vitals, and evidence of Vitals' accuracy based on its validity (measurement to standard) and reliability (consistently valid measurement over time). During the structured interviews, providers also emphasized that educational materials distributed to both providers and patients before use would be critical to

effective use. Examples included a Vitals data sheet, Vitals onscreen prompts on patients' smart devices to guide them through use experience, and video training on both the VA website and social media platforms, e.g., YouTube.

For providers in general, I'd probably start with evidence base...Like I would say "We can get this data and here is the basis of how this data is collected...Here is what the system, the software is doing to give us this data and here's how it's been validated." (Provider 211, MD)

[Need to provide patients with a prompt on screen before Vitals starts] ...Maybe some sort of automated technical cue] "We are now collecting your vitals, so...Please make sure your face...stays centered"...or whatever you need to do to make sure you get the appropriate readings. (Provider 211, MD)

Patients did not express any concerns with usability of Vitals. However, several patients mentioned the challenge of logging in or maintaining an internet connection throughout the visit. Moreover, some patients mentioned that it would be challenging to hold up their smartphone or tablet for the required 45-60 seconds (to obtain the vitals reading) and suggested that VA provide a tabletop stand to hold and stabilize the device during the reading.

I think if -- releasing it [Vitals] with the blood pressure, you get way more utilization with it. (Veteran 201)

...my thought just being that if you want consistency between the readings, there's actually three variables. The head, the camera, and the hand. So, if you can remove variability for the camera and the hand, now it's just the head. (Veteran 201)

Even just something like a clipboard tripod. You know the old things that the secretaries would use... That simple stand somebody used to put a steno pad in and do their notes on. (Veteran 203)

Acceptability of Vitals

Providers and patients rated Vitals high on the Acceptability of Intervention Measure (AIM) reflecting their willingness to use Vitals (Figure 3). They perceived Vitals readings as being accurate, resulting in trust. Neither group expressed concern with security. Providers liked the fact that patients could take their own vitals anytime with the app, that it educates and engages them in their health, an important part of patient-centered care.

I feel like I could trust it. Yeah. To me it's all cool technology. Other big tech companies are doing this stuff, so I'm sure there is good evidence and research behind the technology. Yeah, I trust it. (Provider 201, DPN)

I like the fact that the Veterans can see it themselves, too, which is helpful because they can get a little patient education with it, learn every time they do a VVC, they can see it.

(Provider 206, Licensed Vocational Nurse (LVN))

I would at least be open it, like if there was an issue with blood pressure or something like that, because a lot of times a lot of our Veterans have hypertension...so I know accurate readings are important...I think she's [primary care physician] more in fear of 'What if it's inaccurate and I have to treat it and something happens to the patient.' And it was – 'Did it need to be treated.' Or maybe 'I didn't treat something because it should've been treated.' I think that's more of a concern.... (Provider 206, LVN)

Patients felt that Vitals was acceptable for their care stating that it helped them keep track of their vitals between visits, was easier than other ways of collecting vitals and helped with their virtual care visits. They did not, however, want this to replace the option of seeing a doctor in-person. One Veteran expressed this qualified acceptance, saying:

...as long as the provider didn't default and decline to see us when we felt like we needed to have an in-person visit. (Veteran 201).

Patients offered suggestions on how to promote the technology to them, such as having providers explain Vitals' value in collecting patient metrics, and VA offering materials on how Vitals readings compare with familiar consumer monitoring devices they use, e.g., Fitbit and Apple Watch.

[Need for providers to offer some initial explanation of Vitals' value to patients.] If you just give me data, I'm either going to be pissed off or I'm going to -- it's going to usurp my confidence in the app, or both. And if it wasn't important, they wouldn't be measuring it. (Veteran 203)

It was very close to heart rate [comparing Vitals to his Fitbit reading]. This had 97, that was one 98, so it's very close. I don't know how accurate this is sometimes...it seemed pretty accurate. (Veteran 203)

Feasibility of Vitals

In both the surveys and interviews, providers and patients found Vitals highly feasible and practical for home-based use. One patient expressed that Vitals could also be empowering. (Figure 1).

I think it's pretty cool. I think it would be nice because we don't -- unless the patient has a blood pressure machine at home, we -- and they can't -- they have to visually show us checking it, we can't technically count it...So, it'd be nice because not everyone has a blood pressure cuff at home."
(Provider 206, LVN)

It seems pretty cool. I like the technology to be able to do that at home instead of having to come in here. Like if I'm having a panic attack, it would be nice to see what's going on with my heart rate and everything. Is the heart actually rapid or was it just a panic attack... 'cause it does feel like a heart attack sometimes [laughing]. (Veteran 201)

Appropriateness of Vitals

Finally, providers and patients rated Vitals high on appropriateness, i.e., fit for purpose, per the Intervention Appropriateness Measure (IAM) (Figure 1). Providers believed that most clinical teams would readily integrate Vitals into their routine workflow because it saves time and delivers accurate, consistently collected vitals and may reduce reporting errors in the electronic health record.

You could just say like 'This has been validated and however it's been validated'...then maybe references for people to look up. So, if I am a clinician who maybe is not necessarily believing and I wanna look at factual...strength of evidence... I can't state that enough that it's reproducible, valid, and correlates with what I would get if I had taken vital signs in person...knowing that both have problems.

(Provider 211, MD)

I want integration into CPRS, and I want to make sure that it'll work with Genesis... (Provider 202, NP) [Note: VA's Computerized Patient Record System (CPRS) is being replaced by the Cerner Millennium solution that powers the Department of Defense's Military Health System (MHS GENESIS).³⁹]

Moreover, providers from different specialties indicated that Vitals is appropriate in a variety of clinical contexts. For example, several specialty care providers mentioned how Vitals would enhance their workflow, help with billing for virtual visits, and keep patients safer for preparation appointments. Specialty providers said Vitals could be used for virtual physical therapy appointments and medication management with the pharmacy.

So, we could finish our exercise, we could say hit the heart button and then their vitals would be -- just be popping up as we go. Or I mean every [inaudible]. It would simplify what we do quite a bit.
(Provider 203, DPN)

“I know that with pharmacy, they see Veterans for hypertension to monitor their blood pressure so they can taper or increase or decrease their medication. So, I mean I don’t know how far it’s gonna go or who’s gonna access it, but I know it could be beneficial to like many people, not just primary care, but I know like there’s a lot of people who follow-up on this data...” (Provider 206, LVN)

Also, primary care providers expressed that Vitals could help improve virtual visits and help streamline the collection of patient vitals. They also expressed interest in an application for follow-up visits.

I think we're trying to move towards check-ins, VVC check-ins, like, mirroring how it happens in real life. Like, when you go to a doctor, LVN or nurse comes and checks you in, takes your vitals. So, we're trying to mirror that or get to a point where we can mirror that with our virtual appointments, where an LVN will be checking them in. So, I think at that point would be a good point to get these vitals. (Provider DPN)

Yes. It's vastly [better] because there are no vitals. If we have any, it's because we're looking for the Veterans to provide those. You don't know if their instruments are up to date. I always ask, “How old is your blood pressure cuff?”... And you also wonder if they're coming to the provider, they have to give a good blood pressure reading...you want to be able to see something and to do it in real time, I think that's awesome. (Provider 201, LPN)

We have a lot of screening questions. So, if it would be -- it’s perfect if it works and they can talk and we can continue through like the check-in process, because that’s when we would be using in. (Provider 206, LVN)

Discussion

Principal Findings

Overall, healthcare providers and Veteran patients at two VA facilities found that contactless remote collection of vital signs presents a usable, acceptable, feasible and appropriate way to collect patient health information. Our findings highlight some of the similarities and unique perspectives concerning usability and intent to use among healthcare providers and patients. The inclusion of patients presents a full description of usability of contactless collection of vital signs which is critical for developing implementation plans. This is one of the first accounts of testing a virtual vitals tool with patients.

First, providers and patients expressed strong support for Vitals' usability in their discussion and survey scores. Through hands-on experience in a Vitals demonstration, they found the app to be useful and easy to use. Providers liked that Vitals "helps to provide some valuable context" and "helps to triage" patient's general health and particular conditions. For successful adoption of Vitals, however, providers indicated their need for technical assistance during the Vitals' introduction. Providers and patients emphasized that educational materials distributed before use would be critical to effective use. Providers would need a detailed report of Vitals' accuracy based on its validity (measurement to standard) and reliability (consistently valid measurement over time) with references to peer-reviewed articles. Patients would need a clear, concise fact sheet. Providers and patients indicated that they would also benefit from video trainings (at VA provider- and patient-facing sites or YouTube). Providers suggested that patients' ease of use could be increased if Vitals displayed onscreen prompts on patients' digital devices (e.g., to guide the patient through the Vitals process and display). Patients found Vitals more convenient than conventional remote monitoring because it required less equipment and set-up. However, patients cautioned that VA and providers should make clear that they always have the choice of an in-person visit rather than a virtual visit. Providers and patients recognized that use of Vitals could eliminate the cost and inconvenience of some home-based peripherals (e.g., blood pressure cuff), and potentially reduce the risk of patients' misreading their vitals. Patients and providers urged support for Veteran patients with mobility challenges (e.g., hand tremors, loss of an arm), suggesting that VA offer a platform to steady the patient's tablet or phone for accurate, consistent readings.

Second, regarding acceptability or willingness to use, providers generally expressed confidence and trust in Vitals readings. Still, they requested technical evidence, including citations to academic studies, to assure Vitals' accuracy. Providers felt that use of Vitals would save time, potentially increase the accuracy of vitals records, and promote more consistent clinical practice in taking vitals at the start of all visits. Patients also found Vitals acceptable, offered suggestions on how to present the technology to make it more acceptable to them, such as training providers to explain the Vitals app and its metrics during its introduction to patients, and offering a device stand to steady the cameras of patients' devices, and encouraging patients to compare Vitals' readings with those of their Fitbit or Apple Watch.

Finally, providers noted that Vitals' flexibility makes it feasible, appropriate and advantageous for implementation in a wide range of clinical contexts, particularly in specialty care. Providers and patients both gave Vitals high marks for its feasibility, pleased that it could eliminate use of in-home peripherals and allow patients to take their vitals at any time. Providers believed that most clinical teams would readily integrate Vitals into their routine workflow because it saves time; delivers accurate, consistently collected vitals; and may reduce reporting errors. Providers signaled their readiness to implement Vitals once FDA approval was given, and anticipated potential future capabilities. Specialty care providers, in particular, elaborated on how Vitals would enhance their workflow, help with billing for virtual visits, and keep patients safer for preparation appointments. Primary care providers expressed that Vitals could improve virtual visits and standardize vitals collection.

Limitations

Several limitations to our findings should be considered. The small size of both provider and patient samples may have introduced potential bias. For example, all 13 patient participants were male, and they self-identified as either Black/African American, Hispanic/Latinx or Asian. Offering some balance in representation, over half of providers identified as female and providers identified across all race/ethnicity categories. Participants were associated with only two VA healthcare systems, both

in urban settings that precluded a rural patient- or provider perspective. Future work should employ larger, more geographically diverse samples of Vitals users, both providers and patients, as well as investigate the experiences of those who tried Vitals once but did not return, to understand why and how to promote sustained use.

Conclusions

An essential step in VA's expanded access for Veterans is the increase in utility and adoption of virtual care tools such as VA Video Connect.¹¹ The contactless Vitals feature on their smart devices has the potential to make vitals collection faster and easier for VA providers and patients, and provide accurate, real-time vitals reporting and recording. Integration of this feature would first require FDA approval and technical development but offers the potential to support additional clinical use cases, especially for Veterans with reduced mobility, those in underserved communities, and those in geographically remote areas to complete appointments when in-person care is not required. This ensures greater access to healthcare including treatment and prescribed medications and greater healthcare equity.

Acknowledgements

The authors thank Cortney Frosch, Alex Mujica, Alexander Hamilton, David Yanez and Gabriela Betancourt-Flores of Iron Bow Technologies, LLC under the direction of the VA Office of Connected Care (VA OCC), and Kathleen Craig, BSN, RN of VA Boston Healthcare System for their help in this evaluation. Views expressed are those of the authors and the contents do not necessarily represent the views of the US Department of Veterans Affairs or the United States Government.

This work was supported by the VA OCC and the VA Virtual Care Consortium of Research (VC CORE). LG and DM were supported by VA OCC and VC CORE. ER was supported by the US Department of Veterans Affairs Office of Academic Affiliations Advanced Fellowship in Health Services Research. DM and ER were also supported by Project EBP 22-104 of the Evidence, Policy, and Implementation Center (EPIC), VA Quality Enhancement Research Initiative (QUERI).

Data Availability

Due to US Department of Veterans Affairs (VA) regulations and our ethics agreements, the analytic data sets used for this study are not permitted to leave the VA firewall without a Data Use Agreement. This limitation is consistent with other studies based on VA data. However, VA data are made freely available to researchers with an approved VA study protocol. For more information, please visit <https://www.virec.research.va.gov> or contact the VA Information Resource Center at VIREC@va.gov.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Semi-structured interview guide for providers

Multimedia Appendix 2

Semi-structured interview guide for patients.

Multimedia Appendix 3

Web-based survey for providers.

Multimedia Appendix 4

Web-based survey for patients.


References

1. Veterans Health Administration. URL: <https://www.va.gov/health/> [accessed 02-07-2024].
2. Heyworth L, Shah N, Galpin K. 20 Years of Telehealth in the Veterans Health Administration: Taking Stock of Our Past and Charting Our Future. *J Gen Intern Med*. Feb 20 2024;doi:10.1007/s11606-024-08617-w
3. Garvin LA, Hu J, Slightam C, McInnes DK, Zulman DM. Use of video telehealth tablets to increase access for veterans experiencing homelessness. *J Gen Intern Med*. 2021;1-9.
4. Shura RD, Brearly TW, Tupler LA. Telehealth in Response to the COVID-19 Pandemic in Rural Veteran and Military Beneficiaries. *The Journal of Rural Health*. 2021;37:200-204.
5. Zulman DM, Wong EP, Slightam C, et al. Making connections: nationwide implementation of video telehealth tablets to address access barriers in veterans. *JAMIA Open*. 2019;2(3):323-329.
6. Heyworth L, Kirsh, S., Zulman, D. M., Ferguson, J. M., Kizer, K. W. Expanding access through virtual care: the VA's early experience with Covid-19. *NEJM Catalyst Innovations in Care Delivery*. 2020;1(4)
7. 115th Congress of the United States of America. John S. McCain III, Daniel K. Akaka, and Samuel R. Johnson VA Maintaining Internal Systems and Strengthening Integrated Outside Networks Act of 2018. S.2372. Published 1994. Accessed December 12, 2023. <http://www.copyright.gov/legislation/s505.pdf>.
8. Zulman DM. Expanding Access for Veterans through VA-Issued Video Telehealth Tablets, Research Brief to VA/Stanford Tech Research and Programs. 2021;
9. Dhanani Z, Ferguson JM, Van Campen J, et al. Overcoming Access Barriers for Veterans: Cohort Study of the Distribution and Use of Veterans Affairs' Video-Enabled Tablets Before and During the COVID-19 Pandemic. Original Paper. *J Med Internet Res*. 2023;25:e42563. doi:10.2196/42563
10. Wray CM, Van Campen J, Hu J, Slightam C, Heyworth L, Zulman DM. Crossing the digital divide: a veteran affairs program to distribute video-enabled devices to patients in a supportive housing program. *JAMIA Open*. 2022;5(2)doi:10.1093/jamiaopen/ooac027
11. Slightam C, Wray C, Tisdale RL, Zulman DM, Gray C. Opportunities to Enhance the Implementation of Veterans Affairs Video-Based Care: Qualitative Perspectives of Providers from Diverse Specialties. *J Med Internet Res*. Apr 24 2023;25:e43314. doi:10.2196/43314
12. Wray CM, Ferguson JM, Greene L, et al. Overcoming Obstacles: Barriers to Virtual Care Use Among Video-Enabled Tablet Recipients in the Veterans Health Administration. *J Gen Intern Med*. 2023/11/01 2023;doi:10.1007/s11606-023-08468-x
13. Zulman DM, Verghese A. Virtual care, telemedicine visits, and real connection in the era of COVID-19: unforeseen opportunity in the face of adversity. *JAMA*. 2021;325(5):437-438.
14. Connolly SL, Sullivan, J.L., Lindsay, J.A. et al. Factors influencing uptake of telemental health via videoconferencing at high and low adoption sites within the Department of Veterans Affairs during COVID-19: a qualitative study. *Implement Sci Commun*. 2022;3:66. doi:<https://doi.org/10.1186/s43058-022-00318-x>
15. Shreck E, Nehrig N, Schneider JA, et al. Barriers and facilitators to implementing a US Department of Veterans Affairs Telemental Health (TMH) program for rural veterans. *Journal of Rural Mental Health*. 2020;41(1):1.
16. Gujral K, Van Campen, J., Jacobs, J., Kimerling, R., Blonigen, D., & Zulman, D. M. Mental health service use, suicide behavior, and emergency department visits among rural US veterans who received video-enabled tablets during the COVID-19 pandemic. *JAMA network*

open. 2022;5(4):e226250-e226250.

17. Gujral K, Scott, J. Y., Ambady, L., Dismuke-Greer, C. E., Jacobs, J., Chow, A., ... & Yoon, J. A Primary Care Telehealth Pilot Program to Improve Access: Associations with Patients' Health Care Utilization and Costs *Telemedicine and e-Health*. 2021;
18. Reddy A, Gunnink E, Deeds SA, et al. A rapid mobilization of 'virtual' primary care services in response to COVID-19 at Veterans Health Administration. *Healthcare*. 2020;8(4):100464.
19. Ferguson JM, Wray, C. M., Jacobs, J., Greene, L., Wagner, T. H., Odden, M. C., Freese, J., Van Campen, J., Asch, S. M., Heyworth, L., Zulman, D. M. Variation in Initial and Continued use of Primary, Mental Health, and Specialty Video Care among Veterans. *Health Serv Res*. Nov 7 2022;
20. Mackwood MB, Nagpal AS, Yuen J, Cancino RS. Virtual Access to Subspecialty Care. *Prim Care*. Dec 2022;49(4):557-573. doi:10.1016/j.pop.2022.05.001
21. Barbeito A, Raghunathan K, Connolly S, et al. Barriers to Implementation of Telehealth Pre-anesthesia Evaluation Visits in the Department of Veterans Affairs. *Fed Pract*. Jul 2023;40(7):210-217a. doi:10.12788/fp.0387
22. Qiao D, Ayesha, A.H., Zulkernine, F.H., Jaffar, N., & Masroor, R. ReViSe: Remote Vital Signs Measurement Using Smartphone Camera. *IEEE Access*. 2022;10:131656-131670.
23. Drake C, Lian T, Cameron B, Medynskaya K, Bosworth HB, Shah K. Understanding telemedicine's "new normal": variations in telemedicine use by specialty line and patient demographics. *Telemedicine and e-Health*. 2022;28(1):51-59.
24. Hamilton AB, Finley EP. Qualitative methods in implementation research: an introduction. *Psychiatry research*. 2019;280:112516.
25. Hennink MM, Kaiser BN. Sample sizes for saturation in qualitative research: A systematic review of empirical tests. *Social Science & Medicine*. November 2 2021;114523.
26. Fonteyn ME, Kuipers B, Grobe SJ. A description of think aloud method and protocol analysis. *Qualitative health research*. 1993;3(4):430-441.
27. Van Someren MW, Barnard YF, Sandberg JAC. *The think aloud method: a practical approach to modelling cognitive*. AcademicPress; 1994.
28. Davies EL, Martin J, Foxcroft DR. Development and Acceptability of a Co-Produced Online Intervention to Prevent Alcohol Misuse in Adolescents: A Think Aloud Study *JMIR Human Factors*. 2015;2(2):e13.
29. Venkatesh V, Morris MG, Davis GB, Davis FD. User acceptance of information technology: Toward a unified view. *MIS quarterly*. 2003:425.
30. Harris PA, Taylor, R., Thielke, R., Payne, J., Gonzalez, N., Conde, J.G. Research Electronic Data Capture (REDCap) - A Metadata-Driven Methodology and Workflow Process for Providing Translational Research Informatics Support. *J Biomed Inform*. Apr 2009;42(2):377-81.
31. Lyon AR. Intervention Usability Scale (IUS). *University of Washington*. 2016;
32. Brooke J. *System Usability Scale (SUS): Usability Evaluation in Industry*. Taylor and Francis; 1996.
33. Lewis JR. The system usability scale: past, present, and future. *International Journal of Human-Computer Interaction*. 2018;34(7):577-590. *International Journal of Human-Computer Interaction*. 2018;34(7):577-590.
34. Weiner BJ, Lewis CC, Stanick C, et al. Psychometric assessment of three newly developed implementation outcome measures. *Implementation Science*. 2017;12(1):1-12.


35. Brooke J. SUS: a retrospective. *Journal of usability studies*. 2013;8(2):29-40.
36. Beebe J. *Rapid Qualitative Inquiry: A Field Guide to Team-Based Assessment (2nd Ed.)*. Rowman & Littlefield; 2014.
37. Gale RC, Wu J, Erhardt T, et al. Comparison of rapid vs in-depth qualitative analytic methods from a process evaluation of academic detailing in the Veterans Health Administration. *Implementation Science*. 2019;14(1):11.
38. White CN, Kauffman BY, Acierno R. Factors contributing to veterans' satisfaction with PTSD treatment delivered in person compared to telehealth. *Journal of telemedicine and telecare*. 2021:1357633X20987704.
39. VA Achieves Critical Milestone in Its Electronic Health Record Modernization Program. URL: <https://digital.va.gov/ehr-modernization/news-releases/va-achieves-critical-milestone-in-its-electronic-health-record-modernization-program/> [accessed 2024-02-07].




Supplementary Files

Figures


rPPG technology on smartphones can scan patient faces to deliver vital statistics for patient and provider view.



Views of Vitals on Veterans' smartphones and on VA Providers' tablets (The purple arrows point to the Vital sign readings on both devices).



VA Provider and Veteran average score (out of 100 points) for Vitals on scales for usability, acceptability, feasibility and appropriateness to context.



Multimedia Appendixes

Semi-structured interview guide for providers.

URL: <http://asset.jmir.pub/assets/c7e5bf057447edd47ac99068bef4e264.docx>

Semi-structured interview guide for patients.

URL: <http://asset.jmir.pub/assets/97c9ecbf01a985756996771f74138c82.docx>

Web-based survey for providers.

URL: <http://asset.jmir.pub/assets/1171b4fb385e5acf96cc2c864900a48e.docx>

Web-based survey for patients.

URL: <http://asset.jmir.pub/assets/46da6fbd32f856353ef4cb7045a85ac4.docx>

