

# **Effect of Semaglutide on Body Composition, Physical Function, and Biomarkers of Aging in Older, Overweight Adults with Insulin Resistance: Protocol for an Open-labeled Randomized Controlled Trial**

Tiffany Cortes, Libia Vasquez, Monica C Serra, Ronna Robbins, Allison Stepanenko, Kevin Brown, Hannah Barrus, Annalisa Campos, Sara E Espinoza, Nicolas Musi

Submitted to: JMIR Research Protocols  
on: May 28, 2024

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

*Table of Contents*

---

**Original Manuscript..... 5**  
**Supplementary Files..... 20**  
    Multimedia Appendixes ..... 21  
        Multimedia Appendix 1..... 21



# Effect of Semaglutide on Body Composition, Physical Function, and Biomarkers of Aging in Older, Overweight Adults with Insulin Resistance: Protocol for an Open-labeled Randomized Controlled Trial

Tiffany Cortes<sup>1,2,3</sup> MD; Libia Vasquez<sup>4</sup> MD; Monica C Serra<sup>2,3,5</sup> PhD; Ronna Robbins<sup>6</sup> PhD; Allison Stepanenko<sup>2</sup> NP; Kevin Brown<sup>2</sup> BS; Hannah Barrus<sup>2</sup> BS; Annalisa Campos<sup>2</sup> BS; Sara E Espinoza<sup>7</sup> MD, MSc; Nicolas Musi<sup>7</sup> MD

<sup>1</sup>Division of Endocrinology Department of Medicine University of Texas Health Science Center San Antonio San Antonio US

<sup>2</sup>Sam & Ann Barshop Institute for Longevity & Aging Studies University of Texas Health Science Center San Antonio San Antonio US

<sup>3</sup>San Antonio Geriatric Research, Education, and Clinical Center South Texas Veterans Health Care System San Antonio US

<sup>4</sup>Texas Diabetes Institute University Health System San Antonio US

<sup>5</sup>Division of Geriatrics, Gerontology & Palliative Medicine Department of Medicine University of Texas Health Science San Antonio San Antonio US

<sup>6</sup>Department of Nutrition and Food Science Texas Woman's University Denton US

<sup>7</sup>Center for Translational Geroscience, Diabetes and Aging Center Department of Medicine Cedars-Sinai Medical Center Los Angeles US

## Corresponding Author:

Tiffany Cortes MD

Division of Endocrinology

Department of Medicine

University of Texas Health Science Center San Antonio

7703 Floyd Curl Dr

San Antonio

US

## Abstract

**Background:** Older adults with type 2 diabetes mellitus (T2DM) or prediabetes are at increased risk of adverse changes in body composition, physical function, and concentrations of biomarkers of aging compared with those with normal glucose tolerance. Semaglutide is a glucagon-like peptide 1 receptor agonist that has been approved for T2DM and chronic weight management. Although semaglutide is effective for weight loss and T2DM management, its effects on lean body mass, physical function, and biomarkers of aging are understudied in older adults.

**Objective:** This randomized, open-label, controlled trial aims to compare the effects of semaglutide in addition to lifestyle counseling vs. lifestyle counseling alone on body composition, physical function, and biomarkers of aging in 20 older adults (n=10 per group) with elevated BMI (27-40 kg/m<sup>2</sup>) and glucose intolerance (HgbA1c 5.7-7.5%).

**Methods:** Older adults (>65 y) with prediabetes or well-controlled T2DM will be recruited, stratified by gender, and randomized to one of the two groups (semaglutide plus lifestyle counseling vs lifestyle counseling alone) and followed for five months. Those in the semaglutide group will be titrated to 1 mg weekly, as tolerated, for 12 weeks. Lifestyle counseling will be given by Registered Dietitians and is based on the Diabetes Prevention Program Lifestyle Change Program. Our primary outcomes include changes in lean mass, physical function, and biomarkers of aging. Body composition will be measured by dual-energy X-ray absorptiometry and include total fat mass and lean mass. Physical function will be measured by 6-minute walk distance, grip strength, and short physical performance battery. Biomarkers of aging will be measured in blood, skeletal muscle, and abdominal adipose tissue to include C-reactive protein, interleukin-6, tumor necrosis factors  $\alpha$  and  $\gamma$ , and  $\beta$ -galactosidase staining.

**Results:** The study was funded in December 2021 with a projected data collection period from April 2023 through Summer 2024.

**Conclusions:** Despite the elevated risk of adverse changes in body composition, physical function, and biomarkers of aging among older adults with glucose intolerance and elevated adiposity, the benefits, and risks of commonly prescribed anti-hyperglycemic or weight loss medications such as semaglutide are understudied. This study aims to fill this knowledge gap to inform clinicians about non-glycemic effects of semaglutide and optimize management in this population.

(JMIR Preprints 28/05/2024:62667)

DOI: <https://doi.org/10.2196/preprints.62667>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#)

## Original Manuscript

## **Effect of Semaglutide on Physical Function, Body Composition, and Biomarkers of Aging in Older, Overweight Adults with Insulin Resistance: Protocol for an Open-labeled Randomized Controlled Trial**

**Tiffany M. Cortes, MD<sup>1,3,4</sup>; Libia Vasquez, MD<sup>1</sup>; Monica C. Serra, PhD<sup>2-4</sup>; Ronna Robbins, PhD<sup>6</sup>; Allison Stepanenko, APRN<sup>3</sup>; Kevin Brown<sup>3</sup>; Annalisa Campos<sup>3</sup>; Hannah Barrus<sup>3</sup>; Sara Espinoza, MD<sup>2,7,9</sup>; Nicolas Musi, MD<sup>7-8</sup>**

<sup>1</sup>Division of Endocrinology, <sup>2</sup>Division of Geriatrics, Gerontology & Palliative Medicine, <sup>3</sup>Sam & Ann Barshop Institute for Longevity & Aging Studies, Department of Medicine, UT Health San Antonio,

San Antonio, TX

<sup>4</sup>San Antonio Geriatric Research, Education, and Clinical Center (GRECC), South Texas Veterans Health Care System, San Antonio, TX

<sup>5</sup>Texas Diabetes Institute, University Health System, San Antonio, TX

<sup>6</sup>Department of Nutrition and Food Science, Texas Woman's University, Denton, TX

<sup>7</sup>Center for Translational Geroscience, <sup>8</sup>Diabetes and Aging Center, Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA

### **Corresponding Author:**

**Tiffany Cortes, MD**

**7703 Floyd Curl. MC 7755**

**San Antonio, TX 78229**

**[cortest@uthscsa.edu](mailto:cortest@uthscsa.edu)**

**210-450-8102**

## Abstract

**Background:** Older adults with type 2 diabetes mellitus (T2DM) or prediabetes are at increased risk of adverse changes in body composition, physical function, and aging-related biomarkers compared to those with normal glucose tolerance. Semaglutide is a glucagon-like peptide 1 receptor agonist that has been approved for T2DM and chronic weight management. Although semaglutide is effective for weight loss and T2DM management, its effects on lean body mass, physical function, and biomarkers of aging are understudied in older adults.

**Objective:** This trial aims to compare the effects of lifestyle counseling with and without semaglutide on body composition, physical function, and biomarkers of aging in older adults.

**Methods:** This is a randomized, open-label, controlled trial. Twenty older adults ( $\geq 65$  y) with elevated BMI ( $27\text{--}40\text{ kg/m}^2$ ) and prediabetes or well-controlled T2DM (HgbA1c  $5.7\text{--}7.5\%$ ) are recruited, stratified by sex, and randomized 1:1 to one of two groups (semaglutide plus lifestyle counseling vs lifestyle counseling alone) and followed for five months. Those in the semaglutide group are titrated to 1 mg weekly, as tolerated, for 12 weeks. Lifestyle counseling is given by Registered Dietitians and based on the Diabetes Prevention Program Lifestyle Change Program. Our primary outcomes include changes in lean mass, physical function, and biomarkers of aging. Body composition is measured by dual-energy X-ray absorptiometry and includes total fat mass and lean mass. Physical function is measured by 6-minute walk distance, grip strength, and short physical performance battery. Biomarkers of aging are measured in blood, skeletal muscle, and abdominal adipose tissue to include C-reactive protein, interleukin-6, tumor necrosis factors  $\alpha$ , and  $\beta$  galactosidase staining.

**Results:** The study was funded in December 2021 with a projected data collection period from Spring 2023 through Summer 2024.

**Conclusion:** Despite the elevated risk of adverse changes in body composition, physical function, and biomarkers of aging among older adults with glucose intolerance and elevated adiposity, the benefits and risks of commonly prescribed anti-hyperglycemic or weight loss medications such as semaglutide are understudied. This study aims to fill this knowledge gap to inform clinicians about the potential for additional clinically meaningful, non-glycemic effects of semaglutide.

**Trial Registration:** The study is registered with ClinicalTrials.gov (NCT05786521).

**Keywords:** GLP1; lean body mass; physical function; biomarkers of aging

## Introduction

Glucose intolerance affects most older adults in the United States with 49% having prediabetes and an additional 29% having type 2 diabetes mellitus (T2DM) [1]. The mechanisms that lead to age-related glucose intolerance and T2DM are unknown, but age-dependent decreases in  $\beta$ -cell function and insulin sensitivity are thought to be important in the deterioration of glucose homeostasis with age [2]. Unfavorable changes in body composition observed with aging can contribute to insulin resistance, including redistribution of fat from peripheral subcutaneous areas to a central location [3] and loss of lean muscle mass [4]. Overall, changes observed with aging appear to lead to glucose intolerance and T2DM.

Although T2DM complications are primarily thought of as nephropathy, neuropathy, and retinopathy, T2DM is associated with detrimental changes in physical function and body composition, and biomarkers of aging. A recent meta-analysis reports a 50-80% increased risk of mobility disability; defined as walking speed, chair stand time, and balance test; for those with T2DM compared to people without T2DM [5]. In multiple longitudinal studies, older adults with T2DM have a greater decline in lean mass compared to those without T2DM [6-8]. Additionally, those with prediabetes compared to normoglycemia appear at increased functional risk [9]. These data suggest that insulin resistance, even in the early stages, is associated with adverse changes in body composition and physical function.

Historically, glycemic control has been the focus of T2DM management but the non-glycemic cardiovascular, renal, and weight benefits of some classes of anti-hyperglycemic agents lead to increased individualization of T2DM therapy regimen based on a patient's co-morbidities even after glycemic goals are reached. Semaglutide is a glucagon-like peptide 1 (GLP-1) receptor agonist. GLP-1 receptor agonists utilize the physiologic action of the incretin GLP-1 to increase insulin secretion and suppress glucagon release in response to meals. These drugs reduce hemoglobin A1c (HgbA1c) by 1.45-1.55%, depending on dose [10, 11]. Semaglutide additionally has non-glycemic benefits, including chronic weight management and reduction of cardiovascular events. A recent meta-analysis determined that semaglutide has an average weight loss of 12% with 54% of participants achieving at least 15% weight loss [12]. Currently, semaglutide is FDA-approved not only for T2DM management, to decrease cardiovascular events in adults with T2DM, and for chronic weight management for those with a body mass index (BMI)  $\geq 30$  kg/m<sup>2</sup> or  $\geq 27$  kg/m<sup>2</sup> with at least one weight-related comorbidity. Therefore, a significant portion of older adults maybe prescribed GLP-1 receptor agonists consideration 29% T2DM and 42.8% have a BMI  $\geq 30$  kg/m<sup>2</sup> [1, 13].

With the significant weight loss resulting from semaglutide use, concomitant lean body mass losses may occur. However, the results on lean mass appear mixed, with most trials indicating that semaglutide results in reductions in total fat mass and lean mass with most of the weight loss resulting from fat mass loss [14-17], while others revealed that lean body mass remained stable [18-20]. The effect of semaglutide on physical function has been infrequently studied in older adults. One study found the preservation of grip strength [14] but other validated measures of function in older adults such as short physical performance battery and six-minute walk distance have not been evaluated. Additional measures of muscle quality such as histology and mitochondrial respiration have not been completed in humans. Of note, these trials have an average studied population under 65 years of age, limiting the ability to extrapolate the results to older adults. These data suggest a gap in the literature related to the effects of semaglutide on body composition and physical function in older adults.

Since semaglutide has positive changes in glycemic control, cardiovascular system, and weight management; there are possible changes to the aging process as well. Although there is no standard set of biomarkers to obtain for evaluation of aging, systemic interleukin (IL)-6, tumor necrosis factors  $\alpha$ (TNF $\alpha$ ), C-reactive protein (CRP), and growth differentiating factor 15 (GDF15) are proposed biomarkers that are used in geroscience guided clinical trials[21]. A process that contributes



to aging is cellular senescence which is a stable cell cycle arrest triggered in normal cells that can lead to impaired tissue regeneration and chronic age-associated disease [22]. Cellular senescence can be detected by staining tissues for senescence-associated  $\beta$  galactosidase (SA- $\beta$ -gal) and senescence-associated secretory phenotype (SASP) [22]. Although semaglutide has not been widely studied, the class of GLP-1 receptor agonists shows evidence of improvement in CRP and TNF $\alpha$  [23] with one study indicating that CRP has a significant decline following 68 weeks of treatment with semaglutide [24]. Therefore, this underscores the necessity for further research in this field.

## Objectives

Our goals are to (1) ascertain the potential ramifications of semaglutide on the lean mass profiles and physical functionality of individuals with prediabetes or T2DM and (2) to assess the impact of semaglutide on molecular biomarkers of aging in blood, adipose, and skeletal muscle tissues. We hypothesize that semaglutide will induce favorable alterations in body composition, particularly for lean and adipose body mass quantified via Dual Energy X-ray Absorptiometry (DXA), and improve physical function, as assessed by metrics such as the Short Physical Performance Battery (SPPB), handgrip strength, and walking speed. Additionally, clinical changes will be associated with positive changes in systemic and tissue-specific biomarkers of aging.

## Methods

### Trial Design

This is a randomized, open-label pilot study. The institutional review board is notified of any deviations from the approved protocol or breaches of confidentiality.

As shown in Table 1, this study involves 8 visits which include screening and baseline assessments (visit 1 and visit 2), randomization (visit 2), intervention follow-up (visits 3-5), end-of-study assessments (visit 6a and 6b) and a post-intervention telemedicine follow up visit (visit 7). The intervention phase occurs over 20 weeks. Participants are contacted before in-person visits to ensure they are able to attend or reschedule if needed. All in-person visits are conducted at the Sam and Ann Barshop Institute for Longevity and Aging Studies, University of Texas Health Science Center at San Antonio.

Table 1. Schedule of Events

	Screening	Intervention start, T=0wk	T + 3wk ( $\pm$ 7d)	T + 7wk ( $\pm$ 7d)	T + 11wk ( $\pm$ 7d)	T + 20wk ( $\pm$ 7d)	T + 20wk ( $\pm$ 7d)	T + 21wk ( $\pm$ 7d)
Visit #	V1	V2	V3	V4	V5	V6a <sup>a</sup>	V6b <sup>a</sup>	V7
Medical History and physical exam	X					X		
Vitals, anthropometric measurements	X	X	X	X	X	X	X	
Clinical labs	X					X		
Randomization		X						
DXA scan	X					X		
ECG	X					X		
SPPB, grip strength, 6-minute walk	X					X		

Surveys (SF-12, CNAQ)	X					X		
Blood, muscle, and adipose biopsy		X					X	
Start of semaglutide for treatment group		X						
Semaglutide dispensed to treatment group		X	X	X	X			
Lifestyle education		X	X	X	X			
Assessment of adverse events		X	X	X	X	X	X	X

<sup>a</sup>Visits 6a and 6b are the end-of-study visits but are split into two visits. DXA: dual-energy x-ray absorptiometry, ECG: electrocardiogram, SPPB: short physical performance battery, SF-12: Short-Form 12, CNAQ: The Council on Nutrition appetite questionnaire.

## Participants and Inclusion and Exclusion Criteria

The study is recruiting 20 older adults from the San Antonio, Texas area. Overall, for this pilot our population is generally healthy older adults who are eligible to receive semaglutide for either indication of T2DM or chronic weight management. The inclusion criteria are as follows: (1) men and post-menopausal women; (2) age  $\geq 65$  years; (3) BMI - 27-40 kg/m<sup>2</sup>; (4) HgbA1c 5.7-7.5% or fasting blood glucose greater than 100 mg/dL; (5) community-dwelling; (6) willingness and ability to comply with protocol requirements. The exclusion criteria are as follows: (1) glomerular filtration rate  $\leq 29$  mL/min/1.73m [25]; (2) hematocrit  $\leq 33\%$ ; (3) thyroid stimulating hormone (TSH)  $> 7$  mIU/L with abnormal free thyroxine; (5) use of medications known to lower blood glucose besides metformin; (6) history of cardiovascular events; (7) systolic blood pressure  $\geq 170$  mmHg or diastolic blood pressure  $\geq 95$  mmHg; (8) active inflammatory, autoimmune, infectious, hepatic, gastrointestinal, malignant or uncontrolled psychiatric disease; (9)  $\geq 5\%$  change in weight during the 3 month period prior to screening; (10) personal or family history of medullary thyroid cancer[26]; (11) personal or family history of multiple endocrine neoplasia syndrome type 2[26]; (12) previous history of pancreatitis[26]; (13) electrocardiogram (ECG) with QTc prolongation (QTc  $> 470$ ms in men or  $> 480$  ms in women) or evidence of ischemic changes; (14) personal history of diabetic retinopathy[26]; (15) allergy to semaglutide; (16) type 1 diabetes or other insulin dependent diabetes. Individuals who initially fail their initial screening are eligible to be rescreened 12 weeks after their initial screening.

## Recruitment

Recruitment methods include electronic medical records and Barshop Call Center Registry queries, community engagement activities, and web-based advertisements (including social media). Potential participants undergo a prescreening questionnaire to ensure they meet the inclusion and exclusion criteria for enrollment. Those who qualify from the prescreening questions are scheduled for a full in-clinic screening evaluation and sent an advance copy of the consent form.

## Screening

We conduct an in-clinic evaluation consisting of medical and social history, current medications;

vital signs; anthropometric measurements; and physical exam. Additionally, a fasting blood profile is collected to assess a complete metabolic panel, complete blood count, HgbA1c, TSH, and coagulation panel. Body composition analysis, evaluation of physical function, and completion of patient-reported measures, as detailed below, also are carried out during this visit.

## Baseline and Post-Intervention Assessment Measures

Assessments are performed at baseline (screening or visit 1) and the end of the study (visit 6a 6b) for both groups (Textbox 1).

### Textbox 1. Study Assessment Measures

#### Laboratory tests

- Clinical labs: After an overnight fast, whole blood is collected for the assessment of complete blood count; comprehensive metabolic panel; hemoglobin A1c; lipid profile; thyroid stimulating hormone; prothrombin time, and partial thromboplastin time.
- Research labs: After an overnight fast, a credentialed research team member obtains blood, skeletal muscle, and abdominal adipose samples. Blood is drawn by staff well-versed in phlebotomy to reduce the number of attempts needed for venipuncture. Vastus lateralis muscle biopsy are obtained with a Bergstrom needle after location anesthesia is achieved. Assessments of inflammation and cellular senescence in muscle, adipose tissue, and blood samples are conducted within the clinical research unit. In adipose tissue, we perform staining for  $\beta$ -galactosidase [27]. For muscle tissue, histological processing and measurement of mitochondrial respiration are carried out immediately, followed by freezing additional tissue for the analysis of interleukin (IL)-6, tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), TNF $\alpha$  receptors 1 and 2, p12, and p16 concentrations. We also investigate changes in SIRT1 activity within muscle tissue. Plasma samples obtained from whole blood are used to measure concentrations of IL-6, TNF $\alpha$ , and C-reactive protein.

#### Anthropometric measurements

- Body mass index (BMI): Weight is measured on a standard, calibrated scale at baseline and throughout the study without shoes and with light clothing. Standing height (cm) is measured using a stadiometer at screening. BMI is calculated as weight (kg)/ height ( $m^2$ ).
- Waist circumference: Waist circumference is measured with a tape measure at the level of the iliac crest.
- Body composition: DXA scans are utilized to measure whole-body adipose and lean mass and total bone mineral density.

#### Physical function

- Short Physical Performance Battery: A standardized and validated assessment developed by the National Institute on Aging for use in the Established Population of the Epidemiologic Studies of the Elderly, which is used to evaluate the physical function of older adults independently from their self-reported abilities [28]. The tests fall under three categories: balance, gait speed, and chair stand test. The entire battery of tests can be finished within 10-15 minutes. The administrator begins by demonstrating all the activities to the participant. The balance assessment comprises side-by-side, semi-tandem, and tandem stands. The gait speed test involves walking along a 4-meter path at their usual pace, repeated twice. Finally, participants are assessed on their ability to rise from a chair without arm assistance, followed by completing five timed chair rises. The maximum score is 12. We will classify a score  $<10$  as at increased risk of all-cause mortality [29].
- Grip strength: Hand grip strength of both the dominant and non-dominant arms were

assessed using a handheld dynamometer[30]. The participant's hand grip strength is assessed and the average grip strength from three trials recorded. This test takes less than 5 minutes. Grip strength for women <16 kg and men >26 kg will be classified as at risk [31].

- 6-minute walk: Participants are asked to walk as quickly as they can, for as many times as they can over a 40-foot track for a total of 6 minutes. The total distance walked is obtained[32]. Values will be compared to normative values for health older adults: 367 m for women and 400 m for men [33].

#### Patient-reported outcomes

- The Council on Nutrition appetite questionnaire (CNAQ): Validated as an evaluation of appetite in older adults to predict anorexia before weight loss [34]. The CNAQ consists of eight items within a single domain, with responses rated on a 5-point scale. The total score is calculated by summing the scores of all eight items, ranging from 8 (indicating the poorest appetite) to 40 (indicating the best appetite).
- The Short- Form 12 : A validated tool for assessing health-related quality-of-life in older adults. A higher score indicated better health with a score of 50 indicating the average score in the general U.S. population [35].

## Intervention

### *Randomization*

Eligible participants are randomized to either the lifestyle counseling group (control) or the semaglutide with lifestyle counseling group (1:1) after their blood draw and muscle and adipose biopsies during visit two. Randomization is performed by the principal investigator at the end of the visit two through tables that were stratified by sex in sequential order at the time of randomization. Ten participants are to be enrolled in each group, with twenty overall enrolled. Due to cost restrictions, placebos are not used, and the principal investigator and the participants are unblinded.

### *Lifestyle Counseling*

All participants meet with a Registered Dietitian for an hour-long diet, exercise, and behavior modification sessions at visit 2-5. When in-person classes are not feasible, virtual sessions are made available. Behavior modification is based on the Diabetes Prevention Program Lifestyle Change Program[36]. Aligned with the Diabetes Prevention Program participants receive instructions for changing their dietary composition based on the U.S. Department of Agriculture's MyPlate guidelines which focus on fruit, vegetables, low-fat dairy, and whole grains. MyPlate reinforces small feasible and affordable changes that result in sustainable eating patterns that promote good health [37].

Participants are given detailed written and verbal instructions to achieve these guidelines. During these sessions, education materials are provided on budget-friendly nutrition for optimal aging: nutrition basics (i.e., healthy dietary patterns, calorie balance, food labels, hydration), recipe modification, and mindful eating. In addition, based on the American College of Sports Medicine physical activity recommendations (duration, intensity, mode) participants are encouraged to perform moderate-intensity physical activity at least 150 minutes per week [38]. Physical activity instruction also focuses on how to safely exercise in their home environments. We also cover topics based on participants' questions, which we anticipate may include managing common comorbidities (i.e., sleep behaviors, preventing and managing diabetes, avoiding excessive sitting, managing stress), and available community resources. At each follow-up lifestyle session, weight is obtained, and previously made goals are assessed to determine success or additional modifications required to achieve goals.

## Semaglutide

Participants are given verbal and written instructions on appropriate preparation and subcutaneous injection of semaglutide are self-administered once a week. At visit two, a semaglutide pen that administers 0.25-0.5 mg dose are provided and participants are monitored as they self-administer their first dose of 0.25mg. They continue semaglutide 0.25 mg for an additional 3 weeks, then increase as tolerated to 0.5 mg on visit 3 for four weeks, and 1 mg once a week on visit 4 for four weeks, and continue 1.0 mg on visit 5 for an additional 8 weeks. The titration planned is as follows: complete 4 weeks of 0.25 mg, 4 weeks of 0.5 mg, and 12 weeks of 1.0 mg. If titration is not tolerated, then the participant is kept on the tolerated dose for a longer period with a retry of the higher dose at the next visit.

At each visit, participants are given a semaglutide pen, along with a schedule for the date of the next planned self-administered injection. Participants are asked to bring their study medication to every visit to determine if the appropriate weekly dose has been administered. Participants are withdrawn from the study for noncompliance with semaglutide injections or inability to increase the weekly dose.

## Statistical Analysis

The primary outcomes include changes in lean mass, 6-minute walk distance, grip strength, and SPPB after the 20-week intervention between the semaglutide and lifestyle compared to lifestyle alone groups. We use intention-to-treat principles per the Consolidated Standard of Reporting Trials for analysis [39]. Experimental results are expressed as means  $\pm$  SEM. Comparisons of means between groups are performed by analysis of variance for repeated measures and general linear model are used to add covariates. Associations, within a group, between aging-related biomarkers vs. body composition and physical function outcomes are determined by Pearson's correlation. For tests of correlation coefficients between groups, we use Fisher's Z transformation. Multiple regression analysis are used to determine the relationship between aging-related biomarkers and body composition and physical function outcomes. Covariates include age, sex, ethnicity, and BMI. Alpha equal to 0.05 is used to determine significance. We plan for 20 participants, 10 per group, to accomplish this pilot study within the funding mechanism timeframe. The data generated through this study will be used are used to guide sample size and power calculations for further studies. The Biostatistics Core of the San Antonio Claude D. Pepper Older Americans Independence Center is assisting with statistical analyses. Planned software analysis program is SAS Institute [version 9.4].

## Monitoring Adverse Events

Adverse events are assessed during set interactions with the participants. This includes in-person visits at the research facility and a telemedicine encounter during visit 7. They will also be provided information on how to contact the research team if they have concerns about medication adverse reactions between visits. All adverse events are logged and reported to the Institutional Review Board. Additionally, adverse events are reviewed by a medical doctor to determine the severity, the event's relationship to the study intervention, and what action should be taken.

## Data Management

Each participant is assigned a study code not associated with their name. Data are saved and stored using REDCap (Research Electronic Data Capture; Vanderbilt University) electronic data management system, which is a secure, password-protected, local area network hosted at the University of Texas Health Science Center at San Antonio. Participants' charts, which will include all the visits' documents, are kept inside a locked cabinet located in the clinical research space.

## Ethical Approval and Considerations

The study is approved by the University of Texas Health Science Center San Antonio Institutional

Review Board (20220256HU) in accordance with the ethical standards of the responsible committees on humane experimentation and the Declaration of Helsinki. Eligible participants sign the informed consent after a research staff member has explained the study and answered all questions. Participants may choose to leave or opt out of the study at any time without penalty. Participants receive compensation for their time up to \$425 through a ClinCard if all visits are completed. Privacy and confidentiality is maintained by having only authorized research staff accessing patient health information (PHI), keeping participant files in a locked cabinet, and providing access to REDCap to the minimum authorized research staff. The study is registered with ClinicalTrials.gov (NCT05786521).

## RESULTS

The study was funded in December 2021, with a projected data collection period lasting from Spring 2023 through Summer 2024. Data analysis and results are aimed to be published in late 2024.

## Discussion

### Summary

The purpose of this study is to investigate the effect of semaglutide in addition to lifestyle counseling compared to lifestyle alone on body composition, physical function, and systemic and local biomarkers of aging. We hypothesize that semaglutide therapy will improve body composition and physical function and that these clinical improvements are supported by positive changes to biomarkers of aging.

GLP-1 receptor agonists are a relatively novel group of anti-hyperglycemic agents that are prescribed in increasing frequency [40]. Thus far, glycemic control, cardiovascular benefits, and whole-body weight reduction have been observed in people treated with semaglutide [10, 41, 42]. The majority of studies on GLP-1 receptor agonist use suggest a loss of lean mass, though this is not universally observed [18, 19]. With the popularity of these medications, the additional effects on health parameters especially important to older adults such as function and lean body mass need to be thoroughly evaluated.

Although there have been few studies examining the effect of semaglutide on measures of healthspan in older adults, several preclinical studies have been conducted. Semaglutide treatment in obese mice resulted in an increase of type I/II muscle fibers, muscle fiber density, mitochondrial mass, and decline of TNF $\alpha$ , IL6, and IL-1 $\beta$  compared to untreated obese mice, resulting in the semaglutide-treated mice having a muscle quality similar to the non-obese control group [43]. In a mouse model of sarcopenic obesity, where mice were fed a high-fat diet, treatment with semaglutide led to enhancements in muscle strength, muscle fiber characteristics, and markers of muscle atrophy when compared to mice that did not receive treatment. [44]. The improvements in muscle quality and strength seen with semaglutide may be through the Sirtuin1(SIRT1) pathway which is a key regulator of skeletal muscle biology and function. SIRT1 has also been associated with the ability of GLP-1 receptor agonists to improve hepatic steatosis and skeletal muscle insulin resistance [45, 46]. This study intends to add to this mechanistic research by providing an insight into the effects of semaglutide on biomarkers of aging. These findings may be especially relevant to older adults as clinical targets for future evaluation and potential intervention.

### Limitations

Although this is a randomized controlled trial, there are potential limitations. As this is an open-label trial, the lack of blinding can lead to ascertainment bias[47]. The small sample size for this pilot may lead to a type II error. Given the five-month duration of the trial, it is possible that maximum weight

loss may not be attained. Consequently, we lack a comprehensive understanding of the full impact of this class of medication under conditions of maximum weight loss.

## Conclusions

Based upon the known glycemic, cardiovascular, and body weight loss effects of GLP-1 receptor agonists in the treatment of T2DM and chronic weight management, we are conducting a clinical trial to expand this work by examining the effects of semaglutide on body composition, physical function and biomarkers of aging in older adults with prediabetes and well-controlled T2DM. This study fills a significant gap by focusing on the effect of frequently used medication in older adults and infrequently studied outcomes of clinical significance in this population. The findings will serve as valuable insights for future research endeavors, potentially shaping the treatment approaches utilizing GLP-1 receptor agonists among older adults.

## Acknowledgments

The research reported in this publication was supported by the National Institute on Aging (National Institutes of Health) through the Claude D. Pepper Older Americans Independence Center at the University of Texas Health Science Center at San Antonio (P30 AG044271), Diversity Supplement to Provide Diversity in Health-Related Research (R01AG052697-05S1), and the San Antonio VA Geriatric Research Education and Clinical Center.

## Data Availability Statement

A deidentified final data set underlying all publications resulting from the proposed research will be shared on request pursuant to a data use agreement, appropriately limiting the use of the data set and prohibiting the recipient from identifying or reidentifying (or taking steps to identify or reidentify) any individual whose data are included in the data set.

## Author's Contributions

TMC, LV, and RR wrote the initial draft of the manuscript. All authors have reviewed and edited the manuscript. All authors have approved the manuscript for publication.

## Abbreviations

BMI body mass index

CNAQ The Council on Nutrition appetite questionnaire

CRP C-reactive protein

DSMB data safety monitoring board

DXA dual x-ray absorptiometry

ECG electrocardiogram

GDF15 growth/differentiation factor-15

GLP-1 glucagon-like peptide 1

HgbA1c hemoglobin A1c

IL: interleukin

RD: Registered Dietitian

REDCap research electronic data capture; Vanderbilt University

SF-12 Short-Form 12

SPPB short physical performance battery

TSH thyroid stimulating hormone

T2DM type 2 diabetes mellitus

## References

1. Prevention CfDCA. National Diabetes Statistics Report. Centers for Disease Control and Prevention, US Dept of Health and Human Services;\. 2023.
2. Lee PG, Halter JB. The Pathophysiology of Hyperglycemia in Older Adults: Clinical Considerations. *Diabetes Care*. 2017 Apr;40(4):444-52. PMID: 28325795. doi: 10.2337/dc16-1732.
3. Batsis JA, Zagaria AB. Addressing Obesity in Aging Patients. *Med Clin North Am*. 2018 Jan;102(1):65-85. PMID: 29156188. doi: 10.1016/j.mcna.2017.08.007.
4. Volpi E, Nazemi R, Fujita S. Muscle tissue changes with aging. *Curr Opin Clin Nutr Metab Care*. 2004 Jul;7(4):405-10. PMID: 15192443. doi: 10.1097/01.mco.0000134362.76653.b2.
5. Wong E, Backholer K, Gearon E, Harding J, Freak-Poli R, Stevenson C, et al. Diabetes and risk of physical disability in adults: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol*. 2013 Oct;1(2):106-14. PMID: 24622316. doi: 10.1016/s2213-8587(13)70046-9.
6. Park SW, Goodpaster BH, Lee JS, Kuller LH, Boudreau R, de Rekeneire N, et al. Excessive loss of skeletal muscle mass in older adults with type 2 diabetes. *Diabetes Care*. 2009 Nov;32(11):1993-7. PMID: 19549734. doi: 10.2337/dc09-0264.
7. Park SW, Goodpaster BH, Strotmeyer ES, Kuller LH, Broudeau R, Kammerer C, et al. Accelerated loss of skeletal muscle strength in older adults with type 2 diabetes: the health, aging, and body composition study. *Diabetes Care*. 2007 Jun;30(6):1507-12. PMID: 17363749. doi: 10.2337/dc06-2537.
8. Lee JS, Auyeung TW, Leung J, Kwok T, Leung PC, Woo J. The effect of diabetes mellitus on age-associated lean mass loss in 3153 older adults. *Diabet Med*. 2010 Dec;27(12):1366-71. PMID: 21059088. doi: 10.1111/j.1464-5491.2010.03118.x.
9. Shang Y, Fratiglioni L, Vetrano DL, Dove A, Welmer AK, Xu W. Not Only Diabetes but Also Prediabetes Leads to Functional Decline and Disability in Older Adults. *Diabetes Care*. 2021 Mar;44(3):690-8. PMID: 33446522. doi: 10.2337/dc20-2232.
10. Sorli C, Harashima SI, Tsoukas GM, Unger J, Karsbøl JD, Hansen T, et al. Efficacy and safety of once-weekly semaglutide monotherapy versus placebo in patients with type 2 diabetes (SUSTAIN 1): a double-blind, randomised, placebo-controlled, parallel-group, multinational, multicentre phase 3a trial. *Lancet Diabetes Endocrinol*. 2017 Apr;5(4):251-60. PMID: 28110911. doi: 10.1016/s2213-8587(17)30013-x.
11. Wilding JPH, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, et al. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med*. 2021 Mar 18;384(11):989-1002. PMID: 33567185. doi: 10.1056/NEJMoa2032183.
12. Zhong P, Zeng H, Huang M, Fu W, Chen Z. Efficacy and safety of once-weekly semaglutide in adults with overweight or obesity: a meta-analysis. *Endocrine*. 2022 Mar;75(3):718-24. PMID: 34981419. doi: 10.1007/s12020-021-02945-1.
13. Hales CM CM, Fryar CD, Ogden CL. . Prevalence of obesity and severe obesity among adults: United States, 2017–2018. NCHS Data Brief [Internet]. 2020.
14. Volpe S, Lisco G, Racaniello D, Fanelli M, Colaianni V, Voza A, et al. Once-Weekly Semaglutide Induces an Early Improvement in Body Composition in Patients with Type 2 Diabetes: A 26-Week Prospective Real-Life Study. *Nutrients*. 2022 Jun 10;14(12). PMID: 35745144. doi: 10.3390/nu14122414.
15. Blundell J, Finlayson G, Axelsen M, Flint A, Gibbons C, Kvist T, et al. Effects of once-weekly semaglutide on appetite, energy intake, control of eating, food preference and body weight in subjects with obesity. *Diabetes Obes Metab*. 2017 Sep;19(9):1242-51. PMID: 28266779. doi: 10.1111/dom.12932.
16. Wilding JPH, Batterham RL, Calanna S, Van Gaal LF, McGowan BM, Rosenstock J, et al. Impact of Semaglutide on Body Composition in Adults With Overweight or Obesity: Exploratory Analysis of the STEP 1 Study. *J Endocr Soc*. 2021 May 3;5(Suppl 1):A16-7. PMID: . doi: 10.1210/jendso/bvab048.030.



17. McCrimmon RJ, Catarig A-M, Frias JP, Lausvig NL, le Roux CW, Thielke D, et al. Effects of once-weekly semaglutide vs once-daily canagliflozin on body composition in type 2 diabetes: a substudy of the SUSTAIN 8 randomised controlled clinical trial. *Diabetologia*. 2020/03/01;63(3):473-85. PMID: 31897524. doi: 10.1007/s00125-019-05065-8.
18. Volpe S, Lisco G, Fanelli M, Racaniello D, Colaianne V, Lavarra V, et al. Oral semaglutide improves body composition and preserves lean mass in patients with type 2 diabetes: a 26-week prospective real-life study. *Front Endocrinol (Lausanne)*. 2023;14:1240263. PMID: 37780624. doi: 10.3389/fendo.2023.1240263.
19. Gibbons C, Blundell J, Tetens Hoff S, Dahl K, Bauer R, Bækdal T. Effects of oral semaglutide on energy intake, food preference, appetite, control of eating and body weight in subjects with type 2 diabetes. *Diabetes, Obesity and Metabolism*. 2021;23(2):581-8. PMID: 33184979. doi: <https://doi.org/10.1111/dom.14255>.
20. Uchiyama S, Sada Y, Mihara S, Sasaki Y, Sone M, Tanaka Y. Oral Semaglutide Induces Loss of Body Fat Mass Without Affecting Muscle Mass in Patients With Type 2 Diabetes. *J Clin Med Res*. 2023 Jul;15(7):377-83. PMID: 37575352. doi: 10.14740/jocmr4987.
21. Justice JN, Ferrucci L, Newman AB, Aroda VR, Bahnson JL, Divers J, et al. A framework for selection of blood-based biomarkers for geroscience-guided clinical trials: report from the TAME Biomarkers Workgroup. *Geroscience*. 2018 Dec;40(5-6):419-36. PMID: 30151729. doi: 10.1007/s11357-018-0042-y.
22. Di Micco R, Krizhanovsky V, Baker D, d'Adda di Fagagna F. Cellular senescence in ageing: from mechanisms to therapeutic opportunities. *Nature Reviews Molecular Cell Biology*. 2021 2021/02/01;22(2):75-95. PMID: 33328614. doi: 10.1038/s41580-020-00314-w.
23. Bray JJH, Foster-Davies H, Salem A, Hoole AL, Obaid DR, Halcox JPJ, et al. Glucagon-like peptide-1 receptor agonists improve biomarkers of inflammation and oxidative stress: A systematic review and meta-analysis of randomised controlled trials. *Diabetes, Obesity and Metabolism*. 2021;23(8):1806-22. PMID: 33830637. doi: <https://doi.org/10.1111/dom.14399>.
24. Verma S, Bhatta M, Davies M, Deanfield JE, Garvey WT, Jensen C, et al. Effects of once-weekly semaglutide 2.4 mg on C-reactive protein in adults with overweight or obesity (STEP 1, 2, and 3): Exploratory analyses of three randomised, double-blind, placebo-controlled, phase 3 trials. *EClinicalMedicine*. 2023 Jan;55:101737. PMID: 36467859. doi: 10.1016/j.eclinm.2022.101737.
25. National Health and Nutrition Examination Survey 2017–March 2020 Prepandemic Data Files Development of Files and Prevalence Estimates for Selected Health Outcomes, (2021).
26. Romera I, Cebrián-Cuenca A, Álvarez-Guisasola F, Gomez-Peralta F, Reviriego J. A Review of Practical Issues on the Use of Glucagon-Like Peptide-1 Receptor Agonists for the Management of Type 2 Diabetes. *Diabetes Therapy*. 2019 2019/02/01;10(1):5-19. doi: 10.1007/s13300-018-0535-9.
27. Tchkonja T, Morbeck DE, Von Zglinicki T, Van Deursen J, Lustgarten J, Scrable H, et al. Fat tissue, aging, and cellular senescence. *Aging Cell*. 2010 Oct;9(5):667-84. PMID: 20701600. doi: 10.1111/j.1474-9726.2010.00608.x.
28. Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol*. 1994 Mar;49(2):M85-94. PMID: 8126356. doi: 10.1093/geronj/49.2.m85.
29. de Fátima Ribeiro Silva C, Ohara DG, Matos AP, Pinto A, Pegorari MS. Short Physical Performance Battery as a Measure of Physical Performance and Mortality Predictor in Older Adults: A Comprehensive Literature Review. *Int J Environ Res Public Health*. 2021 Oct 10;18(20). PMID: 34682359. doi: 10.3390/ijerph182010612.
30. Taekema DG, Gussekloo J, Maier AB, Westendorp RG, de Craen AJ. Handgrip strength as a predictor of functional, psychological and social health. A prospective population-based study among the oldest old. *Age Ageing*. 2010 May;39(3):331-7. PMID: 20219767. doi: 10.1093/ageing/afq022.
31. Studenski SA, Peters KW, Alley DE, Cawthon PM, McLean RR, Harris TB, et al. The FNIH

- sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci*. 2014 May;69(5):547-58. PMID: 24737557. doi: 10.1093/gerona/glu010.
32. ATS statement: guidelines for the six-minute walk test. *Am J Respir Crit Care Med*. 2002 Jul 1;166(1):111-7. PMID: 12091180. doi: 10.1164/ajrccm.166.1.at1102.
33. Enright PL, McBurnie MA, Bittner V, Tracy RP, McNamara R, Arnold A, et al. The 6-min walk test: a quick measure of functional status in elderly adults. *Chest*. 2003 Feb;123(2):387-98. PMID: 12576356. doi: 10.1378/chest.123.2.387.
34. Wilson MM, Thomas DR, Rubenstein LZ, Chibnall JT, Anderson S, Baxi A, et al. Appetite assessment: simple appetite questionnaire predicts weight loss in community-dwelling adults and nursing home residents. *Am J Clin Nutr*. 2005 Nov;82(5):1074-81. PMID: 16280441. doi: 10.1093/ajcn/82.5.1074.
35. Ware JE, Jr, Keller SD. SF-12: How to Score the SF-12 Physical and Mental Health Summary Scales. Second ed. Boston, MA: The Health Institute, New England Medical Center; 1995 1995.
36. The Diabetes Prevention Program. Design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care*. 1999 Apr;22(4):623-34. PMID: 10189543. doi: 10.2337/diacare.22.4.623.
37. Services USDoAaUSDoHaH. Dietary Guidelines for Americans, 2020-2025. 9th ed December 2020.
38. Medicine ACoS. ACSM's Guidelines for Exercise Testing and Prescription. 10th ed: Lippincott Williams & Wilkins; 2017.
39. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomized trials. *Ann Intern Med*. 2010 Jun 1;152(11):726-32. PMID: 20335313. doi: 10.7326/0003-4819-152-11-201006010-00232.
40. Adhikari R, Jha K, Dardari Z, Heyward J, Blumenthal RS, Eckel RH, et al. National Trends in Use of Sodium-Glucose Cotransporter-2 Inhibitors and Glucagon-like Peptide-1 Receptor Agonists by Cardiologists and Other Specialties, 2015 to 2020. *Journal of the American Heart Association*. 2022;11(9):e023811. PMID: 35475341. doi: 10.1161/JAHA.121.023811.
41. Marso SP, Bain SC, Consoli A, Eliaschewitz FG, Jódar E, Leiter LA, et al. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *New England Journal of Medicine*. 2016;375(19):1834-44. PMID: 27633186. doi: 10.1056/NEJMoa1607141.
42. Wilding JPH, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, et al. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *New England Journal of Medicine*. 2021;384(11):989-1002. PMID: 33567185. doi: 10.1056/NEJMoa2032183.
43. Ren Q, Chen S, Chen X, Niu S, Yue L, Pan X, et al. An Effective Glucagon-Like Peptide-1 Receptor Agonists, Semaglutide, Improves Sarcopenic Obesity in Obese Mice by Modulating Skeletal Muscle Metabolism. *Drug Des Devel Ther*. 2022;16:3723-35. PMID: 36304787. doi: 10.2147/dddt.S381546.
44. Xiang J, Qin L, Zhong J, Xia N, Liang Y. GLP-1RA Liraglutide and Semaglutide Improves Obesity-Induced Muscle Atrophy via SIRT1 Pathway. *Diabetes, Metabolic Syndrome and Obesity*. 2023 2023/12/31;16(null):2433-46. PMID: 37602204. doi: 10.2147/DMSO.S425642.
45. Jeon JY, Choi SE, Ha ES, Lee HB, Kim TH, Han SJ, et al. GLP-1 improves palmitate-induced insulin resistance in human skeletal muscle via SIRT1 activity. *Int J Mol Med*. 2019 2019/09/01;44(3):1161-71. PMID: 31524229. doi: 10.3892/ijmm.2019.4272.
46. Xu F, Li Z, Zheng X, Liu H, Liang H, Xu H, et al. SIRT1 Mediates the Effect of GLP-1 Receptor Agonist Exenatide on Ameliorating Hepatic Steatosis. *Diabetes*. 2014;63(11):3637-46. PMID: 24947350. doi: 10.2337/db14-0263.
47. Sedgwick P. What is an open label trial? *Bmj*. 2014 May 23;348:g3434. PMID: 24859904. doi: 10.1136/bmj.g3434.



## Supplementary Files

## Multimedia Appendixes

Peer review report from San Antonio Older Adult Independence Center .  
URL: <http://asset.jmir.pub/assets/60b1dc7c97d4bcaaf4463034408294ac.pdf>