

Assessing Social Interaction and Loneliness and Their Association with Frailty among Older Adults with Subjective Cognitive Decline or Mild Cognitive Impairment: An Ecological Momentary Assessment Approach

Bada Kang, Dahye Hong, Seolah Yoon, Chaeun Kang, Jennifer Ivy Kim

Submitted to: JMIR mHealth and uHealth
on: July 30, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript.....	5
Supplementary Files.....	37
Figures	38
Figure 1.....	39
Figure 2.....	40
Figure 3.....	41
Figure 4.....	42
Figure 5.....	43
Figure 6.....	44
Multimedia Appendixes	45
Multimedia Appendix 0.....	46
Multimedia Appendix 1.....	46
Multimedia Appendix 2.....	46
Multimedia Appendix 3.....	46
Multimedia Appendix 4.....	46
Multimedia Appendix 5.....	46

Assessing Social Interaction and Loneliness and Their Association with Frailty among Older Adults with Subjective Cognitive Decline or Mild Cognitive Impairment: An Ecological Momentary Assessment Approach

Bada Kang^{1,2} PhD; Dahye Hong^{1,3} MPH; Seolah Yoon^{1,3} BSN; Chaeun Kang⁴; Jennifer Ivy Kim¹ PhD

¹ Mo-Im Kim Nursing Research Institute Yonsei University College of Nursing Seoul KR

² Institute for Innovation in Digital Healthcare Yonsei University Seoul KR

³ College of Nursing and Brain Korea 21 FOUR Project Yonsei University Seoul KR

⁴ Department of Nursing Yonsei University College of Nursing Seoul KR

Corresponding Author:

Jennifer Ivy Kim PhD

Mo-Im Kim Nursing Research Institute
Yonsei University College of Nursing
50-1 Yonsei-Ro, Seodaemun-Gu
Seoul
KR

Abstract

Background: Frail older adults are at greater risk of adverse health-related outcomes such as falls, disability, and mortality. Mild behavioral impairment (MBI), which is characterized by neurobehavioral symptoms in individuals without dementia, is a crucial factor in identifying at-risk groups and implementing early interventions for frail older adults. However, the specific role of social functioning, which encompasses social interaction and loneliness levels, in relation to frailty within this group remains unclear.

Objective: This study investigated the association between frailty, social interaction, and loneliness levels, measured using an ecological momentary assessment approach, while adjusting for MBI symptoms in two contexts: the presence and severity of MBI symptoms among older adults with subjective cognitive decline (SCD) or mild cognitive impairment (MCI).

Methods: Older adults with SCD or MCI were recruited from an outpatient clinic specializing in the early diagnosis of dementia and care management of dementia at a community health center, as well as from a community service center in Seoul, South Korea. Participants reported their daily social interaction frequency and loneliness level via a mobile application, four times daily for two weeks. Frailty was evaluated using the Korean version of the Frailty Phenotype Questionnaire, and MBI was assessed using a 34-item MBI-Checklist covering five domains. Multivariate logistic regression adjusted for the presence or severity of MBI symptoms was used to analyze their association with frailty.

Results: Among the 101 participants analyzed, 42.6% were classified as frail. Higher average daily social interaction scores were consistently associated with lower odds of frailty. This was evident in the models adjusted for both the global (odds ratio [OR]: 0.51, $p = .01$) and domain-specific presence (OR: 0.42, $p = .01$) of MBI symptoms. Similar trends were observed in the models adjusted for global (OR: 0.52, $p = .04$) and domain-specific (OR: 0.45, $p = .02$) severity of MBI symptom.

Conclusions: Frequent social interaction was inversely associated with frailty, even after adjusting for the presence and severity of MBI symptoms. This study highlights the importance of integrating behavioral and social functional assessments for the early detection, prevention, and targeted intervention for frailty in older adults with cognitive decline. Further prospective studies using real-time measurements are needed to explore the associated risk factors and comprehensive functional status of this at-risk group.

(JMIR Preprints 30/07/2024:64853)

DOI: <https://doi.org/10.2196/preprints.64853>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

✓ **No, I do not wish to publish my submitted manuscript as a preprint.**

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/2018/12/e12345.html>

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Original Paper

Bada Kang, PhD

Mo-Im Kim Nursing Research Institute, Yonsei University College of Nursing, Seoul, Republic of Korea

Institute for Innovation in Digital Healthcare, Yonsei University, Seoul, Korea

Dahye Hong, MPH

Mo-Im Kim Nursing Research Institute, Yonsei University College of Nursing, Seoul, Republic of Korea

College of Nursing and Brain Korea 21 FOUR Project, Yonsei University, Seoul, Republic of Korea

Seolah Yoon, BSN

Mo-Im Kim Nursing Research Institute, Yonsei University College of Nursing, Seoul, Republic of Korea

College of Nursing and Brain Korea 21 FOUR Project, Yonsei University, Seoul, Republic of Korea

Chaeun Kang

Department of Nursing, Yonsei University College of Nursing, Seoul, Republic of Korea

Jennifer Ivy Kim, PhD*

Mo-Im Kim Nursing Research Institute, Yonsei University College of Nursing, Seoul, Republic of Korea

Corresponding Author

Jennifer Ivy Kim, PhD

Mo-Im Kim Nursing Research Institute, Yonsei University College of Nursing

50-1 Yonsei-Ro, Seodaemun-Gu, Seoul 03722, Republic of Korea

E-mail address: redyivy@gmail.com

Phone number: +82-10-3279-2750

Assessing Social Interaction and Loneliness and Their Association with Frailty among Older Adults with Subjective Cognitive Decline or Mild Cognitive Impairment: An Ecological Momentary Assessment Approach

Abstract

Background: Frail older adults are at greater risk of adverse health-related outcomes such as falls, disability, and mortality. Mild behavioral impairment (MBI), which is characterized by neurobehavioral symptoms in individuals without dementia, is a crucial factor in identifying at-risk groups and implementing early interventions for frail older adults. However, the specific role of social functioning, which encompasses social interaction and loneliness levels, in relation to frailty within this group remains unclear.

Objective: This study investigated the association between frailty status, social interaction frequency, and loneliness levels among older adults with subjective cognitive decline (SCD) or mild cognitive impairment (MCI), while adjusting for MBI symptoms in two contexts: the presence and severity of MBI symptoms.

Methods: Older adults with SCD or MCI were recruited from an outpatient clinic specializing in the early diagnosis and care management of dementia at a community health center, as well as from a community service center in Seoul, South Korea. Using an ecological momentary assessment approach, participants reported their daily social interaction frequency and loneliness level via a mobile application, four times daily for two weeks. Frailty status, the outcome variable, was assessed using the Korean version of the Frailty Phenotype Questionnaire. Additionally, MBI symptoms were assessed using the 34-item MBI-Checklist covering five domains. Multinomial logistic regression analyses were performed to investigate the association between frailty status (robust, pre-frail, and frail), the independent variables, adjusting for the presence or severity of MBI symptoms.

Results: Among the 101 participants analyzed, 29.7% were classified as pre-frail, and 12.8% as frail. Higher average daily social interaction scores were consistently associated with lower odds of frail compared to robust status. This was evident in the models adjusted for both the global presence (relative risk ratio [RRR]: 0.18, $p = .02$) and global severity (RRR: 0.20, $p = .02$) of MBI symptoms.

Conclusions: Frequent social interaction was inversely associated with frail status in older adults with SCD or MCI, even after adjusting for the presence and severity of MBI symptoms. These findings highlight the potential of social functioning as a modifiable factor for addressing frailty among older adults at risk of cognitive and functional decline. Future prospective studies using real-time measurements are needed to refine these findings and further investigate additional risk factors and functional outcomes in this group.

Keywords: aged; cognitive impairment; ecological momentary assessment; frailty; mild behavioral impairment

Introduction

The demographic shift toward an increasingly aging population presents significant public health challenges, with a prominent focus on aging-related adverse health outcomes such as frailty. Frailty is a multidimensional syndrome characterized by a decline in physiological reserve across multiple systems, leading to increased vulnerability to stressors and adverse outcomes. These outcomes include physical limitations, such as impaired mobility, muscle weakness, fatigue, cognitive decline, social isolation, and psychological distress [1-3]. Frailty manifests through physical, cognitive, and psychological components, often resulting in increased risks of falls, disability, hospitalization, and mortality [1,2,4]. Physically, frailty impairs an individual's ability to perform daily activities, while cognitively, it may exacerbate the risks of memory and attention deficits [1-6]. Social and psychological factors also play a role, as frailty is often associated with social isolation and depressive symptoms [3,7]. Frailty and pre-frailty incidences were estimated at 43 and 151 new cases per 1,000 person-years, respectively, indicating that approximately one in six community-dwelling older adults may be frail [8]. Frailty is more common in women, older individuals, those with a lower socioeconomic status, and individuals with comorbid chronic diseases and disabilities [1,8-10]. Subjective cognitive decline (SCD) refers to a self-perceived decline in cognitive performance without objective evidence of cognitive impairment [11,12]. Mild cognitive impairment (MCI) represents a state between normal cognition and dementia, characterized by a cognitive decline more pronounced than expected for an individual's age and educational level, but without significant interference in daily activities [13,14]. The co-occurrence of frailty and early-stage cognitive decline has been well documented, with studies showing an increased likelihood of frailty in older adults with SCD or MCI. Two systematic reviews and meta-analyses have offered solid evidence of this relationship [15,16]. Specifically, a cross-sectional study of 2,386 individuals with SCD revealed that they had a higher likelihood of being pre-frail or frail, compared with individuals with normal cognition, even after controlling for factors such as sociodemographic characteristics, physical functionality, psychosocial aspects, and biochemical influences [12]. The Victoria Longitudinal Study, which followed 632 individuals without dementia, also showed that increased frailty was associated with a more rapid memory decline [5]. Therefore, it is essential to identify individuals with SCD or MCI who are at risk of becoming pre-frail or frail, and to implement preventative interventions that focus on modifiable lifestyle and health factors.

Although not all cases of frailty and early cognitive impairment are reversible, evidence supports the potential reversibility of both conditions, including SCD and MCI, through the early detection of at-risk groups and application of tailored interventions [2,15,17,18]. In addition, factors such as age, sex, and activities of daily living are associated with frailty and cognitive decline [19-22], and mild behavioral impairment (MBI) is also correlated with higher levels of frailty and an increased risk of cognitive decline [2,3]. MBI, which emerges later in life, is a neurobehavioral syndrome characterized by sustained and impactful neuropsychiatric symptoms across five domains: decreased drive and motivation, affective dysregulation, impulse dyscontrol, social inappropriateness, and abnormal perception or thought content [23]. MBI is considered a precursor to, or concurrent with, MCI, affecting older adults with either normal cognition or SCD [23]. Significant associations have been observed between certain MBI domains and increased frailty risk in individuals with MCI [3]. Additionally, frailty was associated with both the presence and severity of MBI, especially in men [2]. These findings underscore the importance of assessing MBI to detect risks before the onset of frailty and dementia and warrant the application of interventions for its potential modification. Psychosocial functioning, which encompasses how individuals engage with society and adapt

to environmental challenges, is also an influential factor associated with frailty [24]. Infrequent social interactions, which reflect the quantitative aspects of an individual's limited contact with others [7], and loneliness or perceived social isolation are important factors in this context [25]. Although a definitive theory connecting limited social interaction, loneliness, and frailty has yet to be established, various studies have indicated that reducing social isolation and loneliness in later life can significantly improve overall well-being, underscoring the need for collective consideration of how these factors impact physical health [7]. The English Longitudinal Study of Aging demonstrated that social isolation and loneliness increased the risk of frailty in older adults [26]. Furthermore, given the significant impact of social isolation and loneliness, addressing even moderate levels of these factors is important, as they are associated with an increased risk of worsening frailty [27]. Understanding these psychosocial factors is vital in the broader aging context, as they affect not only physical health but also psychological well-being, thereby potentially influencing older adults' overall quality of life and healthcare needs [7]. Therefore, the interplay between psychosocial factors and health outcomes underscores the importance of holistic approaches in developing geriatric and preventive strategies.

The ecological momentary assessment (EMA) provides a dynamic and immediate record of personal experiences. It is particularly effective for older adults with cognitive decline, offers a comprehensive view of their everyday social lives, and yields more nuanced insights than conventional global measures [28]. By capturing patterns of daily social interaction or loneliness levels, the EMA addresses restrictions on time or recall bias [29]. However, the relationship between frailty, social interaction, and loneliness levels, particularly in the context of MBI symptomatology, remains underexplored, while few studies have used EMA to examine the real-time social functional status among older adults with cognitive decline. Therefore, this study aimed to investigate the cross-sectional relationships between social interaction frequency, loneliness levels, and frailty status among older adults with SCD or MCI in two contexts: 1) adjusting for the presence of MBI symptoms and 2) adjusting for the severity of MBI symptoms. We hypothesized that, similar to MBI symptoms, both the frequency of social interactions and level of loneliness would be significantly correlated with frailty. This exploration is anticipated to yield deeper insights into psychosocial factors related to frailty among individuals with SCD or MCI.

Methods

Design

Our study represents the first wave of a 3-year prospective longitudinal study aimed at developing functional prediction models based on behavioral and psychosocial indicators for older adults with SCD or MCI. The study protocol was published elsewhere [30]. The following is a detailed explanation of the variables considered in this study. We constructed the conceptual framework of our study, based on the social- ecological model for older adults [31] and successful aging model proposed by Rowe and Kahn [32], as shown in Figure 1.

Participants and setting

For the larger 3-year longitudinal study, we initially estimated the required sample size using R (generalized linear mixed model, effect size=0.30, k=2, power=0.80, α =0.05, df=30), yielding a minimum of 106 participants. The study protocol has been published elsewhere [30]. The effect size was selected based on prior studies examining the relationship between frailty and social behavioral factors, which reported moderate effect sizes in similar aging population [33,34]. Based on these findings, we adopted an effect size of 0.30 as a conservative estimate, to

reflect a moderate expected effect while ensuring robustness in our calculation. To account for potential attrition, we adjusted the sample size to 127 participants, anticipating a 20% dropout rate. This adjustment aligns with established guidelines that emphasize careful assessment of attrition rates in aging populations, particularly given the health-related challenges that contribute to higher dropout rates [35,36].

We initially recruited 145 individuals from a dementia relief center and a community service center in Seoul, South Korea (SCD = 91, MCI = 54) between October 2022 and November 2023. The dementia relief center houses an outpatient clinic that specializes in early diagnosis and care management of dementia, whereas the community service center serves as a place for older adults to engage in recreational and wellness programs, irrespective of a dementia diagnosis. Specific criteria were applied to recruit study participants from each center. The common inclusion criteria for the study participants were: 1) age over 65 years, 2) ability to use a smartphone, 3) ability to respond to momentary questionnaires via a mobile application, and 4) ability to provide written consent for participation.

Individuals who met the inclusion criteria were selected for the SCD group. Participants were either: 1) individuals from the dementia relief center who, according to a nurse's report, had neither been diagnosed with MCI nor dementia; or 2) individuals recruited from the community service center through convenience sampling, who self-reported having no history of MCI or dementia diagnosis and scored above 24 on the Korean Mini-Mental State Examination, second edition. Further, we confirmed SCD in individuals from both the dementia relief center and the community service center by ensuring that they provided an affirmative response to the question, "Do you think your memory has gotten worse, compared with the previous year?"

The MCI group consisted exclusively of participants clinically diagnosed with MCI by medical doctors at the dementia relief center. This group was screened to include only individuals who scored above 18 on the Korean Mini-Mental State Examination, second edition. Participants were excluded if they were 1) illiterate; 2) diagnosed with neurological diseases such as epilepsy, stroke, Parkinson's disease, or other types of brain damage; 3) diagnosed with psychiatric diseases such as schizophrenia or bipolar disorder; and 4) undergoing critical illness treatments such as chemotherapy, had severe cardiovascular disease, or had abused substances (including narcotics or alcohol) within the last three years.

After excluding individuals who withdrew their consent ($n = 19$), those who reported EMA responses for less than a week or reported no EMA responses per day ($n = 18$), and those with missing covariate data ($n = 7$), 101 individuals were included in the final sample. The participant selection process is illustrated in Figure 2.

Measures

Frailty

Frailty status was the primary outcome variable in this study. Frailty was assessed using the Korean version of the Frailty Phenotype Questionnaire (FPQ), which is specifically designed to screen for the Fried Frailty Phenotype in community-dwelling older adults [37]. While the Fried Frailty Phenotype requires direct measurements of gait speed, grip strength, and physical activity, the FPQ has been validated as an effective proxy for these assessments. Among Korean older adults, the FPQ demonstrated satisfactory diagnostic accuracy for the Fried Frailty Phenotype, with an area under the curve of 0.89, high sensitivity (81.7%), and specificity (82.5%) [37].

The FPQ is particularly well-suited for epidemiological research due to its low resource requirements and ease of administration, making it a practical screening tool for community-

dwelling older adults where detailed clinical assessments may not be feasible. The FPQ comprises five questions addressing the key aspects of frailty: fatigue (exhaustion), resistance (weakness), ambulation (slowness), inactivity, and weight loss. Scoring from 0 to 5, where 0 indicates robustness, 1–2 indicates pre-frail, and ≥ 3 indicates frail.

Daily Social Interaction Frequency and Level of Loneliness

Daily social interaction frequency and level of loneliness were the main independent variables. Daily social interaction frequency was measured to reflect quantifiable aspects of social contact, while the level of loneliness was assessed to capture one's perceived view of the consequences of social contact [7]. While the level of loneliness is often measured with a single directional question about one's loneliness status, a solid assessment method for social interaction has yet to be established [7]. Despite the complexity and heterogeneity of its indicators, social isolation can be quantitatively assessed by aggregating various factors into a composite index [7]. Building on a prior study that determined social activity frequency by inquiring about the frequency of participant engagement in socially interactive activities among older adults without clinical signs of dementia rated on a five-point scale [38], this study quantitatively assessed the daily frequency of social interaction.

Given the cognitive challenges faced by our participants with SCD or MCI, simplifying the data collection process was essential to reduce participant burden and ensure consistent and reliable responses. For this reason, we did not differentiate between types of social interactions (e.g., in-person, phone-based), as the study's primary goal was to explore the overall relationship between social engagement and frailty in older adults with cognitive decline.

The EMA method was particularly well-suited for this purpose because it allowed us to collect real-time data on both the frequency of social interactions and participants' perceived loneliness. This approach minimizes recall bias, particularly relevant for individuals with cognitive difficulties. A mobile EMA application was used. With smartphone ownership rates among older Korean adults in 2023 reaching 97.2% for those aged 60–69 and 72.9% for those aged 70 and above [39], the feasibility of this method was significantly enhanced.

The EMA method facilitated real-time data collection through a mobile phone application that set off alarms four times daily (upon waking and three times during mealtimes) over two weeks. Each alarm prompted participants to answer two questions. The first question assessed the frequency of social interactions since upon waking or since the last mealtime: "How many times have you had a social encounter?" Participants recorded the number of face-to-face meetings, phone calls, or video calls lasting more than five minutes using a five-point Likert scale (0 for "no contact," 1 for "once," 2 for "two times," 3 for "three times," and 4 for "four or more times"). The second question assessed loneliness: "At this moment, how lonely do you feel?" using a similar five-point Likert scale (0 = "very lonely" to 4 = "not lonely at all"). Both questions were asked four times a day at set intervals (upon waking and after breakfast, lunch, and dinner). Figure 3 shows the captured screens from the mobile application. The research staff provided follow-up contacts for troubleshooting as needed. Participants received gifts worth \$50 after completing the two-week data collection period.

We implemented four strategies to ensure accurate EMA data collection. First, we provided comprehensive training sessions, including repeated practice, to ensure participants could use the mobile application independently. Second, we provided detailed instructional manuals for completing the response process (see Multimedia Appendix 1 for details). Although the EMA questions are simple, we addressed potential challenges given that the participants were individuals with SCD or MCI. Third, we established a helpdesk for assistance with any application-related issues during the follow-up period. Last, we offered direct support as needed, with written contact information provided to ensure participants could reach

researchers via phone or in person. These measures aimed to enhance participants' understanding and proficiency, contributing to more accurate data collection. Approximately 15 participants requested assistance during the study, mainly due to issues with mobile EMA alarms not triggering due to internet connectivity issues or difficulties with smartphone lock functions. Researchers provided phone guidance and, when necessary, conducted in-person visits to resolve the issues and offered additional training.

The quantity of social interaction and the level of loneliness were indicated as daily mean scores based on previous studies [28,40,41]. Data processing involved calculating the maximum daily social interaction frequency as the final daily score. For instance, if a participant reported varying frequencies of social encounters throughout day 1, the highest frequency reported that day was recorded as the final score. The average daily social interaction score was then determined by summing the scores and dividing them by the total number of valid responses. Similarly, the average daily loneliness score was calculated by adding the total loneliness scores recorded each day (with at least one response) and dividing them by the total number of valid response days.

Presence and Severity of MBI Symptoms

The presence and severity of MBI symptoms were included as key covariates. The Mild Behavioral Impairment Checklist (MBI-C) was used to assess the onset of sustained and impactful neurocognitive symptoms in the pre-dementia stage [23]. The MBI-C systematically evaluates symptoms across the five domains of MBI, persisting for at least six months: 1) decreased motivation, 2) affective dysregulation, 3) impulse dyscontrol, 4) social inappropriateness, and 5) abnormal perception or thought content. Each symptom was initially assessed dichotomously (yes/no), followed by a severity rating of mild (1), moderate (2), or severe (3) [23].

In this study, the Korean version of the MBI-C was used to evaluate the presence and severity of MBI symptoms. The MBI-C, a brief screening tool previously validated, was implemented to detect MBI in line with established criteria [3]. The MBI-C has been translated into various languages including Korean. Psychometric evaluation of the Korean version demonstrated significant correlations with the Neuropsychiatric Inventory in individuals with amnesic MCI ($r = 0.25$, $p < .01$) and non-amnesic MCI ($r = 0.36$, $p < .05$) [42,43]. In this study, Cronbach's alpha for the MBI-C was 0.89. For presence assessment, we used a dichotomous method for both domain-specific and global MBI symptoms, with a cut-off score > 0 indicating the presence of MBI symptoms [2]. Severity was assessed as either (1) domain-specific, where the sum of scores from each individual domain was calculated, or (2) global, involving the cumulative sum of scores across all domains [2].

Covariates

Frailty encompasses numerous biological, physiological, and environmental changes that typically occur during the aging process [21]. Therefore, we considered sociodemographic and health-related characteristics, health behaviors, and functional limitations as covariates. The demographic characteristics of participants included gender (male/female), age group (< 70 , 70–74, 75–79, and ≥ 80 years), educational level ($< 9/\geq 9$ years) and living arrangement (living alone/residing with others).

For health-related characteristics, we considered the total number of chronic illnesses diagnosed, including hypertension, diabetes, cardiovascular diseases, cerebrovascular diseases, gastrointestinal diseases, chronic respiratory diseases, chronic kidney diseases, and musculoskeletal conditions. Participants' cognitive status was dichotomously categorized as SCD or MCI based on their responses, which were recorded to evaluate the inclusion criteria.

Health behavior aspects included drinking status (active drinker/non-drinker). Owing to the limited sample size of former or current smokers ($n = 2$), smoking status was excluded from the statistical analysis.

As demonstrated in previous studies, dependence on managing higher levels of functional performance is associated with frailty [21,22]. Therefore, functional limitations were assessed using the Korean Instrumental Activities of Daily Living (K-IADL) scale, which measures the ability to perform complex daily tasks such as cooking, grooming, housekeeping, shopping, financial management, transportation, laundry, short-distance walking, making phone calls, and managing medications [44]. Each item on the K-IADL scale is rated from 0 to 4, with higher scores indicating greater dependency. The K-IADL's validation, conducted with older adults aged over 65 years, demonstrated strong internal consistency (Cronbach's $\alpha = 0.94$), high inter-rater reliability (kappa value = 0.81–0.95), and test-retest reliability correlation (≥ 0.70) [44]. In this study, high K-IADL scores indicated diminished functional capabilities and were analyzed as continuous variables.

Statistical Analysis

Descriptive statistics were used to outline the general characteristics of the participants and compare differences between the robust, pre-frail, and frail groups. Categorical variables are presented as counts and percentages, while continuous variables are summarized as means with standard deviations.

To assess the normality of continuous variables, we performed the Shapiro-Wilk test. If the data was normally distributed and variance was equal, analysis of variance was applied to compare differences between groups. When normality was not met or variance were unequal, the Kruskal-Wallis test was used. A chi-square test was used to assess differences in categorical variables between groups, with Fisher's exact test applied when the expected cell count was less than 5.

We conducted multinomial logistic regression analyses to examine the association between frailty status, social interaction frequency, and loneliness levels among older adults with MCI or SCD. Frailty status was the primary outcome variable and was categorized into three groups: robust, pre-frail, and frail. Social interaction frequency and daily loneliness levels were the independent variables, assessed using EMA.

To account for potential confounding effects, MBI symptoms were included as a key covariate. Additionally, other confounders, including sex, age group, educational level, living arrangement, number of chronic illnesses, cognitive status, health behaviors, and functional limitations, were selected based on their established relevance to frailty and social engagement in previous studies [2,3,19-22].

Separate multinomial logistic regression models were performed to adjust for MBI symptom presence and severity. All independent variables were simultaneously entered into the model to ensure consistency and comparability across analyses. Multicollinearity was assessed using the variance inflation factor to confirm that the included variables were not highly correlated. Model fit was evaluated using the generalized Hosmer-Lemeshow test and model specification was checked for validity. Statistical significance was set at $P < .05$ and all analyses were performed using Stata software version 16 (Stata Corp., Texas, USA).

Ethics Considerations

This study was approved by the Institutional Review Board of Severance Hospital, Yonsei University Health System (No.4-2022-0637). In accordance with ethical guidelines, all participants provided their written informed consent prior to their inclusion in the study. To

ensure ethical integrity, individuals with severe cognitive decline who were assessed as incapable of making informed decisions regarding participation were excluded from the study.



Results

Participants' General Characteristics

Table 1 presents the general characteristics of participants. A total of 101 participants were included in the analysis; 58 (57.4%) were categorized as robust, 30 (29.7%) as pre-frail, and 13 (12.9%) as frail. Significant differences were observed between the groups in age and global MBI symptom severity. The mean age and standard deviations for each group was; robust (74.9 ± 5.1), pre-frail (75.7 ± 6.5), and frail (82.0 ± 4.9), with corresponding age ranges of 65-87, 65-89, and 73-89. respectively. For global MBI severity, the median score for all participants was 3 (Interquartile range (IQR): 1-9.5). The robust group had the lowest median severity score (2, IQR: 1-4.5), while the frail group exhibited the highest severity (9, IQR: 3-23.5).

Table 2 displays a descriptive summary of social interaction frequency and loneliness levels for all participants, stratified by frailty status. We summarized the average maximum daily social interaction score by calculating the average of each participant's daily maximum scores across the study period, resulting in a single, average score per participant. No significant differences were observed in the mean average maximum daily social interaction scores among the groups: robust (3.0 ± 0.8), pre-frail (2.9 ± 0.9), and frail (2.4 ± 0.9). Similarly, no significant differences were found in the mean average maximum daily loneliness levels, with scores of 3.4 ± 0.5 for the robust group, 3.3 ± 0.6 for the pre-frail group, and 3.1 ± 0.9 for the frail group.

Figure 4 illustrates the variations of the average daily social interaction scores and average daily loneliness levels categorized by frailty status. The robust group is characterized by the highest median values in daily social interactions and the lowest loneliness levels compared to pre-frail and frail groups. In contrast, the frail group demonstrates the lowest median social interaction frequency and the highest loneliness levels. Furthermore, the density distribution of mean maximum daily social interaction scores and mean maximum daily loneliness level by frailty status are illustrated in Appendix 2 and 3. While the overall distribution is right skewed across all participants, the frail group exhibits a relatively higher density in the lower frequency of social interactions compared to the robust or pre-frail group (Appendix 2). In addition, the frail group shows a relatively lower density of participants experiencing lower loneliness levels compared to robust and pre-frail groups, even though the overall distribution is right-skewed (Appendix 3).

Table 1. Baseline characteristics of participants by frailty status

Variables	Total (N = 101)	Robust (n = 58)	Pre-frail (n = 30)	Frail (n=13)	P-value
Sex					
Men	42 (41.6)	27 (46.5)	10 (33.3)	5 (38.5)	.47
Women	59 (58.4)	31 (53.5)	20 (66.7)	8 (61.5)	
Age					
Range	65-89	65-87	65-89	73-89	
Mean ± Standard deviation	76.1±5.9	74.9±5.1	75.7±6.5	82.0±4.9	<.001
Educational level					
< 9 years	30 (29.7)	12 (20.7)	12 (40.0)	6 (46.2)	.06
≥ 9 years	71 (70.3)	46 (79.3)	18 (60.0)	7 (53.8)	
Living arrangement^a					
Living alone	19 (18.8)	8 (13.8)	9 (30.0)	2 (15.4)	.22
Residing with others	82 (81.2)	50 (86.2)	21 (70.0)	11 (84.6)	
Cognitive status					
SCD^b	67 (66.3)	44 (75.9)	16 (53.3)	7 (53.9)	.06
MCI^c	34 (33.7)	14 (24.1)	14 (46.7)	6 (46.1)	
Drinking status^a					
Active drinker	25 (24.8)	17 (29.3)	5 (16.7)	3 (23.1)	.46
Non-drinker	76 (75.2)	41 (70.7)	25 (83.3)	10 (76.9)	
Number of chronic illnesses	2.01±1.11	1.96±1.07	1.93±1.14	2.46±1.19	.31
K-IADL^{d,e}					
Range	1-17	1-16	8-17	9-16	
Interquartile range	10-10	10-10	10-10	10-12.5	
Median	10	10	10	10	
Mean ± Standard deviation	10.4±1.9	10.4±2.1	10.4±1.7	11.0±1.9	.64
Global MBI^f symptom presence	81 (80.2)	45 (77.5)	24 (80.0)	12 (92.3)	.64
Global MBI^f symptom severity^e					
Range	0-41	0-25	0-40	0-41	
Interquartile range	1-9.5	1-4.5	1-12.3	3-23.5	
Median	3	2	8.5	9	
Mean ± Standard deviation	6.9±9.0	4.0±5.1	9.2±10.1	14.5±13.8	.002

^a Fisher's exact test was applied; ^b SCD: subjective cognitive decline; ^c MCI: mild cognitive impairment; ^d K-IADL: Korean Instrumental Activities of Daily Living scale; ^e Kruskal-Wallis test was applied; ^f MBI: mild behavioral impairment

Table 2. Descriptive statistics of daily social interaction frequency and loneliness levels by frailty status.

	Total (N = 101)	Robust (n = 58)	Pre-frail (n = 30)	Frail (n=13)	P-value
Social interaction frequency per day					
Range	0-4	0-4	0-4	0-4	
Mean $\pm$ Standard deviation	2.9 $\pm$ 0.8	3.0 $\pm$ 0.8	2.9 $\pm$ 0.9	2.4 $\pm$ 0.9	.08
Loneliness level per day					
Range	0-4	1-4	0-4	0-4	
Mean $\pm$ Standard deviation	3.4 $\pm$ 0.6	3.6 $\pm$ 0.5	3.3 $\pm$ 0.6	3.1 $\pm$ 0.9	.06




Figure 1. The proposed conceptual framework illustrates the relationship between frailty status, social interaction, and loneliness levels, while adjusting for the presence and severity of mild behavioral impairment symptoms among older adults with subjective cognitive decline or mild cognitive impairment. Based on the social-ecological model for older adults and the successful aging model proposed by Rowe and Kahn, individual determinants include sociodemographic characteristics, health status, and health behaviors. Social function, including daily social interaction frequency and daily loneliness level, is also considered.




Figure 2. Sequential steps involved in the recruitment and selection of study participants. We recruited 145 individuals aged ≥ 65 from a dementia relief center or a community service center in Seoul, South Korea. The figure depicts the subdivision of these individuals into three groups: 58 with robust, 30 with pre-frail, and 13 frail, based on specific inclusion and exclusion criteria.



Figure 3. The mobile application screens captured two questions that assessed daily social interaction frequency and loneliness levels. Daily social interaction frequency was measured by asking, "How many times did you interact with others, including face-to-face meetings, phone calls, or video calls lasting more than five minutes?" Responses were categorized into five options: no contact, once, twice, three times, and four or more times. Loneliness levels were measured by asking, "At this moment, how lonely do you feel?" Responses were also categorized into five options: very lonely, lonely, neutral, not lonely, and not lonely at all. Both questions were administered four times daily at specific intervals (upon waking and after breakfast, lunch, and dinner).




Figure 4. Distribution of average daily social interaction score and average daily loneliness level by frailty status. This figure displays the median levels and range variability of both average daily social interaction score and average daily loneliness levels among participants, categorized into robust, pre-frail and frail groups.

Primary Outcomes

Association Between Daily Social Interaction Frequency, Daily Loneliness Level, and Frailty Adjusting for the Presence of Global MBI Symptoms

Table 3 illustrates the relationship between frailty, daily social interaction frequency, and daily loneliness level after adjusting for the presence of global MBI symptoms. After adjusting for the presence of global MBI symptoms, neither the average maximum daily social interaction score (RRR: 0.74, $p=.42$) nor the average maximum daily loneliness level (RRR: 0.89, $p=.76$) were significantly associated with pre-frail status. Similarly, no significant relationship was observed between the average maximum daily loneliness level and frail status (RRR: 0.27, $p=.10$). However, participants with higher average maximum daily social interaction scores were significantly less likely to be frail (RRR: 0.18, $p=.02$). Sociodemographic and health-related characteristics also showed significant associations with frailty. Participants with SCD were less likely to be pre-frail compared to those with MCI (RRR: 0.31, $p=.03$), while older age was associated with a higher likelihood of being frail (RRR: 1.56, $p=.001$). Figure 5 displays the average predictive marginal effects of sociodemographic characteristics, health status, global MBI symptom presence, and social functioning on the probability of being classified as robust, pre-frail, or frail.

Association Between Daily Social Interaction Frequency, Daily Loneliness level, and Frailty Adjusting for the Severity of Global MBI Symptoms

Table 4 presents the associations between frailty, daily social interaction frequency, and daily loneliness levels after adjusting for the severity of global MBI symptoms. When adjusting for the severity of global MBI symptoms, no significant associations were observed between pre-frail status and either the average maximum daily social interaction score (RRR: 0.76, $p=.48$) or the average maximum daily loneliness level (RRR: 1.19, $p=.69$). Likewise, no significant relationship was identified between frail status and average maximum daily loneliness level (RRR: 0.27, $p=.14$). However, a significant inverse relationship was found between the average maximum daily social interaction scores and frail status (RRR: 0.20, $p=.02$). Additional sociodemographic and health-related factors were associated with pre-frail status. Participants with SCD were less likely to be pre-frail compared to those with MCI (RRR: 0.33, $p=.04$). Moreover, participants with more severe MBI symptoms were more likely to be pre-frail (RRR: 1.09, $p=.04$), and older age remained significantly associated with being frail (RRR: 1.53, $p=.002$). Figure 6 presents the average predictive marginal effects of sociodemographic characteristics, health status, global MBI symptom severity, and social functioning on the probability of being classified as robust, pre-frail, or frail.

Table 3. Multinomial logistic regression model for the association between frailty, global MBI-symptom presence, daily social interaction frequency, and level of loneliness (N=101).

Variables	Pre-frail (Base outcome: Robust)		Frail (Base outcome: Robust)	
	RRR ^a (95% CI ^b)	P-value	RRR (95% CI)	P-value
Women (Ref: men)	2.18 (0.57–8.26)	.25	3.54 (0.42–29.72)	.24
Age	1.01 (0.92–1.12)	.79	1.56 (1.20–2.01)	.001
Educational level ≥ 9 years (Ref: < 9 years)	0.46 (0.16–1.33)	.15	0.43 (0.06–2.89)	.38
Living alone (Ref: Residing with others)	2.44 (0.68–8.68)	.17	0.11 (0.003–3.27)	.20
SCD ^d (Ref: MCI ^c)	0.31 (0.11–0.87)	.03	0.35 (0.06–2.17)	.26
Active drinker (Ref: Non-drinker)	0.62 (0.15–2.51)	.50	9.10 (0.61–135.11)	.11
Number of chronic illnesses	0.78 (0.48–1.26)	.31	2.04 (0.73–5.66)	.17
K-IADL ^e	1.13 (0.86–1.49)	.37	1.38 (0.85–2.25)	.18
Global MBI ^f -symptom presence	0.89 (0.25–3.15)	.85	0.36 (0.01–8.91)	.53
Average maximum daily social interaction score	0.74 (0.36–1.53)	.42	0.18 (0.05–0.72)	.02
Average maximum daily loneliness level	0.89 (0.41–1.92)	.76	0.27 (0.05–1.31)	.10
Wald χ^2	52.04			

^a RRR: relative risk ratio; ^b CI: confidence interval; ^c MCI: mild cognitive impairment

^d SCD: subjective cognitive decline; ^e K-IADL: Korean Instrumental Activities of Daily Living scale; ^f MBI: mild behavioral impairment.




Figure 5. This figure shows the average predictive marginal effects of sociodemographic characteristics, health status, global mild behavioral impairment symptom presence, and social functioning on the probability of being classified as robust, pre-frail, or frail. The predictive margins were calculated using multinomial logistic regression, with error bars representing 95% confidence intervals for each effect.

Table 4. Regression models for the association between frailty, global MBI-symptom severity, daily social interaction frequency, and level of loneliness (N=101).

Variables	Pre-frail (Base outcome: Robust)		Frail (Base outcome: Robust)	
	RRR ^a (95% CI ^b)	P-value	RRR (95% CI)	P-value
Women (Ref: men)	2.22 (0.56–8.71)	.25	2.93 (0.38–22.34)	.30
Age	0.97 (0.88–1.10)	.63	1.53 (1.16–2.02)	.002
Educational level $\geq$ 9 years (Ref: < 9 years)	0.52 (0.17–1.59)	.26	0.54 (0.08–3.65)	.52
Living alone (Ref: Residing with others)	2.58 (0.69–9.69)	.16	0.08 (0.002–2.57)	.16
SCD ^d (Ref: MCI ^c)	0.33 (0.11–0.97)	.04	0.44 (0.07–2.57)	.36
Active drinker (Ref: Non-drinker)	0.65 (0.15–2.68)	.55	10.29 (0.61–174.13)	.11
Number of chronic illnesses	0.72 (0.44–1.18)	.20	2.14 (0.74–6.12)	.15
K-IADL ^e	1.10 (0.84–1.44)	.49	1.31 (0.81–2.13)	.27
Global MBI ^f -symptom severity	1.09 (1.01–1.18)	.04	1.06 (0.95–1.17)	.32
Average maximum daily social interaction score	0.76 (0.36–1.61)	.48	0.20 (0.05–0.78)	.02
Average maximum daily loneliness level	1.19 (0.51–2.78)	.69	0.27 (0.05–1.56)	.14
Wald χ^2	56.59			

^a RRR: relative risk ratio; ^b CI: confidence interval; ^c MCI: mild cognitive impairment

^d SCD: subjective cognitive decline; ^e K-IADL: Korean Instrumental Activities of Daily Living scale; ^f MBI: mild behavioral impairment.




Figure 6. This figure shows the average predictive marginal effects of sociodemographic characteristics, health status, global mild behavioral impairment symptom severity, and social functioning on the probability of being classified as robust, pre-frail, or frail. The predictive margins were calculated using multinomial logistic regression, with error bars representing 95% confidence intervals for each effect.

Discussion

Principal Findings

This study demonstrated a significant association between frailty and the frequency of daily social interaction in a cohort of 101 individuals at risk of dementia in South Korea. Specifically, higher average daily social interaction scores were significantly associated with lower relative risk ratios of being frail, even after adjusting for the presence and severity of global MBI symptoms. These findings underscore the potential role of daily social interaction frequency as a modifiable behavioral factor in individuals at the early stages of dementia and frailty risk. Our study found that frail older adults exhibited more severe MBI symptoms than robust or pre-frail groups. This observation aligns with prior research suggesting a potential link between frailty and the severity of MBI symptoms [2]. Furthermore, our findings indicate that frail older adults had lower average maximum daily social interaction frequency and higher loneliness levels compared to robust or pre-frail groups. Unlike previous studies that assessed social interaction or loneliness at a single time point [7], our study employed a two-week tracking period. This approach also builds on previous research exploring how frailty in older adults, characterized by decreased mobility and difficulty with daily activities, may contribute to reduced social engagement, increased loneliness, and further functional decline [7].

Our findings also align with existing evidence showing that both the presence and severity of global MBI symptoms are associated with frailty. This underscores the importance of assessing both cognitive and behavioral impairments when identifying individuals at risk of dementia [2]. However, given the relatively small sample size of frail individuals in this study, these findings should be interpreted with caution. Nonetheless, our results support prior research linking neuropsychiatric symptoms, such as psychosis, not only to an elevated risk of dementia but also frailty, which is often associated with accelerated cognitive decline [45]. To strengthen these observations, future research should consider analyzing larger cohorts with detailed assessments of domain-specific MBI symptoms and frailty. Such studies could yield more robust evidence, further elucidating the relationship between specific MBI domains and frailty and reinforcing their potential as targets for early intervention in individuals at risk of dementia and functional decline.

Our findings are consistent with those of a previous study that used data from the Korean Frailty and Aging Cohort Study, which found that less frequent contact with acquaintances was associated with a higher prevalence of frailty [46]. Similar patterns have been observed worldwide. For instance, a one-year prospective cohort study in Japan demonstrated that greater social isolation, measured using the Lubben Social Network Scale, increased the risk of developing pre-frail status [47]. Among Italian community-dwelling older adults, a significant relationship was found between frailty and higher levels of social isolation, measured using the Friendship Scale, and frailty [48]. Although the precise mechanisms through which social connections influence frailty are not yet fully understood, growing epidemiological evidence suggests that social interactions can affect health outcomes through psychosocial, behavioral, and biological pathways. A lack of social interactions, for instance, can amplify the sympathetic nervous system and the hypothalamus-pituitary-adrenal axis, thus leading to a chronic mild inflammatory state, which is characteristic of frailty syndrome [49-51]. Limited social interactions can also increase stress-related hormones such as glucocorticoids, impacting immune functions and potentially affecting the biological aspects of frailty [52]. Furthermore, frequent social interactions can motivate frail older adults to engage in cognitively and physically stimulating activities, providing daily support and health management resources [52,53].

Overall, our findings underscore the importance of social interaction correlates to frailty among older adults. This issue is particularly relevant in East Asian societies, where traditional family structures and intergenerational support are highly valued [54]. However, the rapid demographic shift toward an aging population and declining family size in developed East Asian countries, such as South Korea, Japan, and China, poses challenges to maintaining adequate social contact for older adults [54,55]. This phenomenon may increase the risk of frailty in older adults who lack sufficient social engagement due to changing family dynamics [56]. Furthermore, in developed countries, where healthcare systems are more established, promoting social functioning could be a cost-effective, non-pharmacological intervention to reduce frailty and its associated healthcare burden [57,58]. These findings suggest that public health policies aimed at enhancing social interactions, particularly for those at risk of cognitive decline, could reduce frailty and promote healthy aging in both East Asian and developed countries more broadly.

Our study did not establish a significant correlation between daily loneliness and frailty, in contrast to previous research. This discrepancy may be attributed to two primary reasons. First, our method of measuring loneliness was based on a single direct question covering a two-week period. While we aimed to reflect on the participants' subjective perceptions of their social interactions, a more in-depth, long-term study may be necessary to capture the full complexity of loneliness. A systematic review and meta-analysis of the relationship between loneliness and physical frailty in community-dwelling older adults indicated that baseline loneliness could predict worsening frailty, although longitudinal evidence is limited [59]. Therefore, future studies should focus on the cumulative effects of comprehensive potential factors prevalent in older adults, such as decreased physical activity, poor sleep quality, and malnutrition [59-61]. Additionally, these studies should examine these factors within the context of social vulnerability and frailty, while controlling for the effects of social networks [61].

Second, unlike the frequency of social interactions, loneliness is a subjective feeling and perceptions of loneliness can vary significantly among individuals [7]. Typically, loneliness is seen as a negative response to the gap between actual and desired relationships [62]. However, the threshold for feeling lonely often depends on sociocultural context or personal standards [63,64]. Therefore, employing more nuanced, context-specific questionnaires could lead to a deeper understanding of loneliness [7]. Nonetheless, given that our study participants were older adults at risk for dementia, using numerous detailed questionnaires might not be the most feasible approach. An alternative method could involve analyzing patterns among participants who provide similar responses to brief, directional questions, and then prospectively studying their perceptions of loneliness in response to specific events in a real-time setting. This approach could yield valuable insights into subjective experiences of loneliness in this population.

Comparison with Prior Work

A key strength of our study lies in its detailed exploration of the relationship between social interaction and frailty status among older adults with SCD or MCI. By categorizing frailty into robust, pre-frail, and frail groups, our findings build upon and extend previous research, indicating that older adults with higher levels of daily social interaction are less likely to exhibit frailty, even after accounting for cognitive and behavioral symptoms. While these findings align with prior research suggesting that protective effects of social interaction on health, our study adds nuance by examining these relationships specifically within older adults at risk of cognitive decline.

Our results support existing evidence that frail older adults exhibit lower social interaction

frequency and higher loneliness levels compared to their robust and pre-frail counterparts. These findings underscore the importance of social interaction as a critical behavioral dimension for health outcomes in older adults at risk of both cognitive and functional decline. While causation cannot be established, our findings suggest that fostering social connections may mitigate frailty risks and promote better health outcomes.

What distinguishes our study is its use of real-time data collection methods, which allowed us to capture day-to-day variations in social functioning with greater accuracy [40]. This approach offers a more granular understanding of how social interaction relates to frailty and cognitive status, building on prior retrospective studies. By applying real-time assessment within the context of frailty and cognitive decline, this study strengthens the evidence base for social interaction as a modifiable factor and highlights the need for future research to explore causative pathways.

Although the immediate practical application of these findings in clinical settings may be limited, this study contributes to the growing understanding of the relationship between social function, frailty, and cognitive decline. It highlights the potential for improving social engagement interventions that may reduce physical challenges targeted to older adults with SCD or MCI. Future research should focus on identifying the specific types, patterns, and contexts of social interaction that are most effective in reducing frailty and promoting health. Longitudinal and interventional studies will be crucial in establishing causation and designing tailored interventions to enhance social functioning in these at-risk populations.

Limitations

This study had several limitations. First, the sample size of 101 participants, particularly the small sample size of the frail group, may limit the generalizability of the findings. A larger and more diverse sample could yield more robust results and allow for detailed analyses across different subgroups. Second, the cross-sectional design of the study enabled the identification of associations but did not establish causation. Prospective longitudinal studies are necessary to determine the directionality of the relationships among frailty, MBI symptoms, and social functioning. Additionally, Alzheimer's disease pathology, hormonal changes, and chronic inflammation are potential mechanistic links between frailty and cognitive decline. While this study focuses on social interaction, physiological dimensions such as physical activity, metabolic health, and nutritional status were not assessed to minimize the participants' measurement burden during the two-week survey period. The primary aim of our study was to demonstrate how capturing temporal patterns in social functioning could inform daily life modifications to support health. Future research should integrate complementary physiological measures to enhance understanding and inform more comprehensive intervention strategies in longitudinal or experimental studies to better understand the directionality of these relationships. For example, studies using matching designs over extended timespans would be essential. Third, this study relied on self-reported data for loneliness levels and the tools used to measure daily social interaction frequency and loneliness levels, which may be susceptible to social desirability bias, which could influence how participants report their feelings and circumstances. Future analyses could incorporate digital devices, such as actigraphy, to log social interaction frequency and patterns, thereby integrating supplementary objective measures. Additionally qualitative aspects of loneliness, such as its depth and contextual nuances, may require more comprehensive assessment methodologies. Tools like daily diaries or note-taking functions are advisable for future EMA approaches to capture more detailed information on the types and quality of social interactions. Moreover, framing questions with both positive and negative perspectives could mitigate response patterns where participants consistently answer in one direction, enhancing the robustness of findings in larger studies.

Fourth, the use of a self-reported frailty assessment via the FPQ may introduce recall bias or inaccuracies in participants' self-perceptions. While the FPQ is a validated and practical screening tool for community-dwelling older adults, it may lack the granularity of more detailed clinical assessments. Future studies could consider integrating tools such as the Clinical Frailty Scale to capture a broader spectrum of frailty characteristics and enhance precision in frailty classification. Finally, while our study provided step-by-step guidance and a helpdesk for participants to navigate the EMA application, applying the EMA approach via mobile devices to older adults less familiar with mobile technology requires more comprehensive and detailed guidance. Ensuring that participants can use the mobile application independently is an essential step for successfully replicating this study methodology.

Conclusions

Our study suggests that daily social interaction is associated with frailty status among older adults with SCD or MCI, irrespective of the presence or severity of MBI symptoms. This finding builds on previous research, highlighting the positive potential of improving social interaction to reduce frailty risk among older adults with cognitive decline. To deepen understanding and enhance timely intervention strategies, future studies should explore the qualitative aspects of social interactions and frailty in greater depth. While the immediate clinical application of these findings may be limited, the use of real-time assessment in social functioning offers a promising path for further investigation. Integrating real-time social data with objective measures could provide a more personalized and comprehensive approach to care.

Acknowledgments

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government (Ministry of Science and ICT) (No. 2022R1C1C2003706), the Basic Science Research Program through the NRF, funded by the Ministry of Education (No. 2020R1A6A1A03041989), and the Brain Korea 21 Four Projects funded by the National Research Foundation.

Data Availability

The datasets used and/or analyzed in the current study are available from the first author upon reasonable request.

Conflict of Interest

The authors declare that they have no competing interests.

Authors' Contributions

BK supervised, conceptualized, and designed the study framework. JK designed the study framework, analyzed the data, drafted and edited the manuscript. DH, SY, and CK recruited the participants and collected the data. All authors contributed collectively to the data interpretation. The final draft of the manuscript has been thoroughly reviewed and approved by all the authors.

Abbreviations

CI: Confidence interval

EMA: Ecological momentary assessment

FPQ: Frailty Phenotype Questionnaire

K-IADL: Korean Instrumental Activities of Daily Living scale

MBI: Mild behavioral impairment

MCI: Mild cognitive impairment

OR: Adjusted odds ratio

SCD: Subjective cognitive decline



References

1. Hoogendijk EO, Afilalo J, Ensrud KE, Kowal P, Onder G, Fried LP. Frailty: implications for clinical practice and public health. *Lancet* 2019 Oct 12;394(10206):1365-1375. [doi: 10.1016/s0140-6736(19)31786-6] [Medline: 31609228]
2. Guan DX, Rockwood K, Smith EE, Ismail Z. Sex Moderates the Association between Frailty and Mild Behavioral Impairment. *J Prev Alzheimers Dis* 2022;9(4):692-700. [doi: 10.14283/jpad.2022.61] [Medline: 36281673]
3. Fan S, Liang X, Yun T, Pei Z, Hu B, Ismail Z, et al. Mild behavioral impairment is related to frailty in non-dementia older adults: a cross-sectional study. *BMC Geriatr* 2020 Nov 27;20(1):510. [doi: 10.1186/s12877-020-01903-2] [Medline: 33246409]
4. Markle-Reid M, Browne G. Conceptualizations of frailty in relation to older adults. *J Adv Nurs* 2003 Oct;44(1):58-68. [doi: 10.1046/j.1365-2648.2003.02767.x] [Medline: 12956670]
5. Thibaut S, McDermott K, McFall GP, Rockwood K, Dixon RA. Frailty effects on non-demented cognitive trajectories are moderated by sex and Alzheimer's genetic risk. *Alzheimer's Research & Therapy* 2019 2019/06/21;11(1):55. [doi: 10.1186/s13195-019-0509-9]
6. Wallace LMK, Theou O, Godin J, Andrew MK, Bennett DA, Rockwood K. Investigation of frailty as a moderator of the relationship between neuropathology and dementia in Alzheimer's disease: a cross-sectional analysis of data from the Rush Memory and Aging Project. *The Lancet Neurology* 2019;18(2):177-184. [doi: 10.1016/S1474-4422(18)30371-5]
7. Hajek A, Riedel-Heller SG, König H-H. Loneliness and Social Isolation in Old Age: Correlates and Implications: Taylor & Francis; 2023.
8. Ofori-Asenso R, Chin KL, Mazidi M, Zomer E, Ilomaki J, Zullo AR, et al. Global Incidence of Frailty and Prefrailty Among Community-Dwelling Older Adults: A Systematic Review and Meta-analysis. *JAMA Network Open* 2019;2(8):e198398-e198398. [doi: 10.1001/jamanetworkopen.2019.8398]
9. Buckinx F, Rolland Y, Reginster JY, Ricour C, Petermans J, Bruyère O. Burden of frailty in the elderly population: perspectives for a public health challenge. *Arch Public Health* 2015;73(1):19. [doi: 10.1186/s13690-015-0068-x] [Medline: 25866625]
10. Won CW, Lee S, Kim J, Chon D, Kim S, Kim C-O, et al. Korean frailty and aging cohort study (KFACS): cohort profile. *BMJ Open* 2020;10(4):e035573. [doi: 10.1136/bmjopen-2019-035573]
11. Jessen F, Amariglio RE, Buckley RF, van der Flier WM, Han Y, Molinuevo JL, et al. The characterisation of subjective cognitive decline. *Lancet Neurol* 2020 Mar;19(3):271-278. [doi: 10.1016/s1474-4422(19)30368-0] [Medline: 31958406]
12. Hsieh TJ, Chang HY, Wu IC, Chen CC, Tsai HJ, Chiu YF, et al. Independent association between subjective cognitive decline and frailty in the elderly. *PLoS One* 2018;13(8):e0201351. [doi: 10.1371/journal.pone.0201351] [Medline: 30071051]
13. Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, et al. Mild cognitive impairment. *Lancet* 2006 Apr 15;367(9518):1262-1270. [doi: 10.1016/s0140-6736(06)68542-5] [Medline: 16631882]
14. Panza F, Lozupone M, Solfrizzi V, Sardone R, Dibello V, Di Lena L, et al. Different Cognitive Frailty Models and Health- and Cognitive-related Outcomes in Older Age: From Epidemiology to Prevention. *J Alzheimers Dis* 2018;62(3):993-1012. [doi: 10.3233/jad-170963] [Medline: 29562543]
15. Borges MK, Canevelli M, Cesari M, Aprahamian I. Frailty as a Predictor of Cognitive Disorders: A Systematic Review and Meta-Analysis. *Front Med (Lausanne)* 2019;6:26. [doi: 10.3389/fmed.2019.00026] [Medline: 30838210]

16. Grande G, Haaksma ML, Rizzuto D, Melis RJF, Marengoni A, Onder G, et al. Co-occurrence of cognitive impairment and physical frailty, and incidence of dementia: Systematic review and meta-analysis. *Neurosci Biobehav Rev* 2019 Dec;107:96-103. [doi: 10.1016/j.neubiorev.2019.09.001] [Medline: 31491474]
17. Ruan Q, Yu Z, Chen M, Bao Z, Li J, He W. Cognitive frailty, a novel target for the prevention of elderly dependency. *Ageing Research Reviews* 2015 2015/03/01;20:1-10. [doi: 10.1016/j.arr.2014.12.004]
18. Solfrizzi V, Scafato E, Lozupone M, Seripa D, Giannini M, Sardone R, et al. Additive Role of a Potentially Reversible Cognitive Frailty Model and Inflammatory State on the Risk of Disability: The Italian Longitudinal Study on Aging. *Am J Geriatr Psychiatry* 2017 Nov;25(11):1236-1248. [doi: 10.1016/j.jagp.2017.05.018] [Medline: 28689645]
19. Panza F, Solfrizzi V, Frisardi V, Maggi S, Sancarolo D, Adante F, et al. Different models of frailty in predementia and dementia syndromes. *J Nutr Health Aging* 2011 Aug;15(8):711-719. [doi: 10.1007/s12603-011-0126-1] [Medline: 21968870]
20. Mulero J, Zafrilla P, Martinez-Cacha A. Oxidative stress, frailty and cognitive decline. *J Nutr Health Aging* 2011 Nov;15(9):756-760. [doi: 10.1007/s12603-011-0130-5] [Medline: 22089224]
21. Nourhashémi F, Andrieu S, Gillette-Guyonnet S, Vellas B, Albarède JL, Grandjean H. Instrumental activities of daily living as a potential marker of frailty: a study of 7364 community-dwelling elderly women (the EPIDOS study). *J Gerontol A Biol Sci Med Sci* 2001 Jul;56(7):M448-453. [doi: 10.1093/gerona/56.7.m448] [Medline: 11445604]
22. Nguyen TTH, Nguyen AT, Vu T-HT, Dau NT, Nguyen PQ, Nguyen TX, et al. Association of Frailty Status and Functional Disability among Community-Dwelling People Aged 80 and Older in Vietnam. *BioMed Research International* 2021 2021/07/19;2021:7109452. [doi: 10.1155/2021/7109452]
23. Ismail Z, Agüera-Ortiz L, Brodaty H, Cieslak A, Cummings J, Fischer CE, et al. The Mild Behavioral Impairment Checklist (MBI-C): A Rating Scale for Neuropsychiatric Symptoms in Pre-Dementia Populations. *J Alzheimers Dis* 2017;56(3):929-938. [doi: 10.3233/jad-160979] [Medline: 28059789]
24. Freitag S, Schmidt S. Psychosocial Correlates of Frailty in Older Adults. *Geriatrics (Basel)* 2016 Nov 1;1(4) [doi: 10.3390/geriatrics1040026] [Medline: 31022819]
25. Cacioppo JT, Fowler JH, Christakis NA. Alone in the crowd: the structure and spread of loneliness in a large social network. *J Pers Soc Psychol* 2009 Dec;97(6):977-991. [doi: 10.1037/a0016076] [Medline: 19968414]
26. Davies K, Maharani A, Chandola T, Todd C, Pendleton N. The longitudinal relationship between loneliness, social isolation, and frailty in older adults in England: a prospective analysis. *Lancet Healthy Longev* 2021 Feb;2(2):e70-e77. [doi: 10.1016/s2666-7568(20)30038-6] [Medline: 36098160]
27. Jarach CM, Tettamanti M, Nobili A, D'Avanzo B. Social isolation and loneliness as related to progression and reversion of frailty in the Survey of Health Aging Retirement in Europe (SHARE). *Age Ageing* 2021 Jan 8;50(1):258-262. [doi: 10.1093/ageing/afaa168] [Medline: 32915990]
28. Zhaoyang R, Sliwinski MJ, Martire LM, Katz MJ, Scott SB. Features of daily social interactions that discriminate between older adults with and without mild cognitive impairment. *J Gerontol B Psychol Sci Soc Sci* 2021 Feb 2 [doi: 10.1093/geronb/gbab019] [Medline: 33528558]
29. van Berkel N, Goncalves J, Lovén L, Ferreira D, Hosio S, Kostakos V. Effect of experience sampling schedules on response rate and recall accuracy of objective self-reports. *International Journal of Human-Computer Studies* 2019 2019/05/01;125:118-128. [doi: 10.1016/j.ijhcs.2018.12.002]

30. Kang B, Ma J, Jeong I, Yoon S, Kim JI, Heo SJ, et al. Behavioral marker-based predictive modeling of functional status for older adults with subjective cognitive decline and mild cognitive impairment: Study protocol. *DIGITAL HEALTH* 2024 2024/01/01;10:20552076241269555. [doi: 10.1177/20552076241269555]
31. Boulton ER, Horne M, Todd C. Multiple influences on participating in physical activity in older age: Developing a social ecological approach. *Health Expect* 2018 Feb;21(1):239-248. [doi: 10.1111/hex.12608] [Medline: 28768065]
32. Rowe JW, Kahn RL. Successful aging. *Gerontologist* 1997 Aug;37(4):433-440. [doi: 10.1093/geront/37.4.433] [Medline: 9279031]
33. Chen LK, Hwang AC, Lee WJ, Peng LN, Lin MH, Neil DL, et al. Efficacy of multidomain interventions to improve physical frailty, depression and cognition: data from cluster-randomized controlled trials. *J Cachexia Sarcopenia Muscle* 2020 Jun;11(3):650-662. [doi: 10.1002/jcsm.12534] [Medline: 32134208]
34. Zhang H, Hao M, Hu Z, Jiang S, Li Y, Wang X, et al. Social frailty and its association with cognitive trajectories in older adults: a prospective cohort study. *Alzheimers Res Ther* 2025 Feb 19;17(1):50. [doi: 10.1186/s13195-025-01687-w] [Medline: 39972471]
35. Fewtrell MS, Kennedy K, Singhal A, Martin RM, Ness A, Hadders-Algra M, et al. How much loss to follow-up is acceptable in long-term randomised trials and prospective studies? *Arch Dis Child* 2008 Jun;93(6):458-461. [doi: 10.1136/adc.2007.127316] [Medline: 18495909]
36. Beller J, Geyer S, Epping J. Health and study dropout: health aspects differentially predict attrition. *BMC Medical Research Methodology* 2022 2022/01/30;22(1):31. [doi: 10.1186/s12874-022-01508-w]
37. Kim S, Kim M, Jung HW, Won CW. Development of a Frailty Phenotype Questionnaire for Use in Screening Community-Dwelling Older Adults. *J Am Med Dir Assoc* 2020 May;21(5):660-664. [doi: 10.1016/j.jamda.2019.08.028] [Medline: 31672563]
38. Krueger KR, Wilson RS, Kamenetsky JM, Barnes LL, Bienias JL, Bennett DA. Social engagement and cognitive function in old age. *Exp Aging Res* 2009 Jan-Mar;35(1):45-60. [doi: 10.1080/03610730802545028] [Medline: 19173101]
39. Jung YC, Kim YH, Oh YS. Korea Media Panel Survey 2023. Korea Information Society Development Institute; 2023.
40. Liu H, Xie QW, Lou VWQ. Everyday social interactions and intra-individual variability in affect: A systematic review and meta-analysis of ecological momentary assessment studies. *Motivation and Emotion* 2019 2019/04/01;43(2):339-353. [doi: 10.1007/s11031-018-9735-x]
41. Kim H, Lee S, Lee S, Hong S, Kang H, Kim N. Depression Prediction by Using Ecological Momentary Assessment, Actiwatch Data, and Machine Learning: Observational Study on Older Adults Living Alone. *JMIR Mhealth Uhealth* 2019 Oct 16;7(10):e14149. [doi: 10.2196/14149] [Medline: 31621642]
42. Lin RS, Yu DS, Chau PH, Li PW, Ismail Z. Reliability and Validity of the Traditional Chinese Version of the Mild Behavioral Impairment – Checklist Among Persons With Mild Cognitive Impairment – A Validation Study. *Journal of Geriatric Psychiatry and Neurology* 2023;36(1):26-38. [doi: 10.1177/08919887221093363] [Medline: 35430911]
43. Kang Y, Chin J, Han N, Park J, Yeom J, Yang S-J, et al. P2-292: Mild Behavioral Impairment (MBI) in MCI, SCD, and Normal Elderly: A Pilot Study For Validation of the Mild Behavioral Impairment Checklist (MBI-C). *Alzheimer's & Dementia* 2018;14(7S_Part_15):P793-P793. [doi: 10.1016/j.jalz.2018.06.981]
44. Won CW, Rho YG, SunWoo D, Lee YS. The Validity and Reliability of Korean Instrumental Activities of Daily Living(K-IADL) Scale. *Ann Geriatr Med Res* 2002 12;6(4):273-280.
45. Ruthirakuhan M, Ismail Z, Herrmann N, Gallagher D, Lanctôt KL. Mild behavioral impairment is associated with progression to Alzheimer's disease: A clinicopathological


- study. *Alzheimers Dement* 2022 Nov;18(11):2199-2208. [doi: 10.1002/alz.12519] [Medline: 35103400]
46. Chon D, Lee Y, Kim J, Lee KE. The Association between Frequency of Social Contact and Frailty in Older People: Korean Frailty and Aging Cohort Study (KFACS). *J Korean Med Sci* 2018 Dec 17;33(51):e332. [doi: 10.3346/jkms.2018.33.e332] [Medline: 30546284]
 47. Uno C, Okada K, Matsushita E, Satake S, Kuzuya M. Friendship-related social isolation is a potential risk factor for the transition from robust to prefrailty among healthy older adults: a 1-year follow-up study. *Eur Geriatr Med* 2021 Apr;12(2):285-293. [doi: 10.1007/s41999-020-00422-y] [Medline: 33128225]
 48. Mulasso A, Roppolo M, Giannotta F, Rabaglietti E. Associations of frailty and psychosocial factors with autonomy in daily activities: a cross-sectional study in Italian community-dwelling older adults. *Clinical Interventions in Aging* 2016 2016/01/11;11(null):37-45. [doi: 10.2147/CIA.S95162]
 49. Wang J, Maxwell CA, Yu F. Biological Processes and Biomarkers Related to Frailty in Older Adults: A State-of-the-Science Literature Review. *Biol Res Nurs* 2019 Jan;21(1):80-106. [doi: 10.1177/1099800418798047] [Medline: 30198309]
 50. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001 Mar;56(3):M146-156. [doi: 10.1093/gerona/56.3.m146] [Medline: 11253156]
 51. Holt-Lunstad J. The Major Health Implications of Social Connection. *Current Directions in Psychological Science* 2021;30(3):251-259. [doi: 10.1177/0963721421999630]
 52. Devita M, Marescalco E, Bordignon A, Trevisan C, Sergi G, Coin A. Social interaction level modulates the impact of frailty on cognitive decline: a longitudinal study. *Aging Ment Health* 2023 Aug 21:1-6. [doi: 10.1080/13607863.2023.2247349] [Medline: 37602456]
 53. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet* 2020 Aug 8;396(10248):413-446. [doi: 10.1016/s0140-6736(20)30367-6] [Medline: 32738937]
 54. Nakagawa T, Cho J, Yeung DY. Successful Aging in East Asia: Comparison Among China, Korea, and Japan. *J Gerontol B Psychol Sci Soc Sci* 2021 Jun 8;76(Suppl 1):S17-s26. [doi: 10.1093/geronb/gbaa042] [Medline: 32324214]
 55. Alavi K, Sutan R, Shahar S, Manaf MRA, Jaafar MH, Abdul Maulud KN, et al. Connecting the dots between social care and healthcare for the sustainability development of older adult in Asia: a scoping review. *Sustainability* 2022;14(5):2573.
 56. Newmyer L, Verdery AM, Wang H, Margolis R. Population Aging, Demographic Metabolism, and the Rising Tide of Late Middle Age to Older Adult Loneliness around the World. *Popul Dev Rev* 2022 Sep;48(3):829-862. [doi: 10.1111/padr.12506] [Medline: 37649654]
 57. Seino S, Nishi M, Murayama H, Narita M, Yokoyama Y, Nofuji Y, et al. Effects of a multifactorial intervention comprising resistance exercise, nutritional and psychosocial programs on frailty and functional health in community-dwelling older adults: A randomized, controlled, cross-over trial. *Geriatr Gerontol Int* 2017 Nov;17(11):2034-2045. [doi: 10.1111/ggi.13016] [Medline: 28393440]
 58. Lestari SK, Eriksson M, de Luna X, Malmberg G, Ng N. Frailty and types of social relationships among older adults in 17 European countries: A latent class analysis. *Archives of Gerontology and Geriatrics* 2022 2022/07/01;101:104705. [doi: 10.1016/j.archger.2022.104705]
 59. Kojima G, Taniguchi Y, Aoyama R, Tanabe M. Associations between loneliness and physical frailty in community-dwelling older adults: A systematic review and meta-analysis. *Ageing Res Rev* 2022 Nov;81:101705. [doi: 10.1016/j.arr.2022.101705] [Medline: 35932978]
 60. Ong AD, Uchino BN, Wethington E. Loneliness and Health in Older Adults: A Mini-Review

- and Synthesis. *Gerontology* 2016;62(4):443-449. [doi: 10.1159/000441651] [Medline: 26539997]
61. Kojima G, Taniguchi Y, Kitamura A, Fujiwara Y. Is living alone a risk factor of frailty? A systematic review and meta-analysis. *Ageing Research Reviews* 2020/05/01/;59:101048. [doi: 10.1016/j.arr.2020.101048]
 62. Russell DW, Cutrona CE, McRae C, Gomez M. Is loneliness the same as being alone? *J Psychol* 2012 Jan-Apr;146(1-2):7-22. [doi: 10.1080/00223980.2011.589414] [Medline: 22303609]
 63. Rapolienè G, Aartsen M. Lonely societies: low trust societies? Further explanations for national variations in loneliness among older Europeans. *Eur J Ageing* 2022 Sep;19(3):485-494. [doi: 10.1007/s10433-021-00649-z] [Medline: 36052198]
 64. Swader CS. Loneliness in Europe: Personal and Societal Individualism-Collectivism and Their Connection to Social Isolation. *Social Forces* 2018;97(3):1307-1336. [doi: 10.1093/sf/soy088]


Supplementary Files

Figures

The proposed conceptual framework illustrates the relationship between frailty status, social interaction, and loneliness levels, while adjusting for the presence and severity of mild behavioral impairment symptoms among older adults with subjective cognitive decline or mild cognitive impairment. Based on the social-ecological model for older adults and the successful aging model proposed by Rowe and Kahn, individual determinants include sociodemographic characteristics, health status, and health behaviors. Social function, including daily social interaction frequency and daily loneliness level, is also considered.




Sequential steps involved in the recruitment and selection of study participants. We recruited 145 individuals aged ≥ 65 from a dementia relief center or a community service center in Seoul, South Korea. The figure depicts the subdivision of these individuals into three groups: 58 with robust, 30 with pre-frail, and 13 frail, based on specific inclusion and exclusion criteria.




The mobile application screens captured two questions that assessed daily social interaction frequency and loneliness levels. Daily social interaction frequency was measured by asking, "How many times did you interact with others, including face-to-face meetings, phone calls, or video calls lasting more than five minutes?" Responses were categorized into five options: no contact, once, twice, three times, and four or more times. Loneliness levels were measured by asking, "At this moment, how lonely do you feel?" Responses were also categorized into five options: very lonely, lonely, neutral, not lonely, and not lonely at all. Both questions were administered four times daily at specific intervals (upon waking and after breakfast, lunch, and dinner).

The screenshot displays a mobile application interface with a green header bar. The main content area is divided into two sections. The left section features a large green square with a white bell icon and the text "어울림 횟수와 외로움의 정도에 대해 응답하실 시간입니다." (It's time to respond to the frequency of social interaction and the degree of loneliness). Below this are two buttons: a blue "응답 할게요" (I will respond) button and a white "나중에 할게요" (I will do it later) button. The right section contains two questionnaires. The first questionnaire asks "기상 이후, 다른 사람과 몇 번이나 어울리셨습니까? (5분 이상의 직접 만남/전화/영상통화 포함)" (After waking up, how many times did you interact with others, including direct meetings/phone/video calls lasting more than 5 minutes?). It has five radio button options: "전혀" (None), "1회" (Once), "2회" (Twice), "3회" (Three times), and "4회 이상" (Four or more times). The second questionnaire asks "어르신은 현재 얼마나 외롭다고 느껴십니까?" (How lonely do you feel at the moment?). It has five radio button options: "전혀 외롭지 않다" (Not lonely at all), "외롭지 않다" (Not lonely), "보통이다" (Average), "외롭다" (Lonely), and "매우 외롭다" (Very lonely). At the bottom of the right section is a green "완료" (Complete) button. The bottom navigation bar shows three icons: a home icon, a circle icon, and a back icon.


Distribution of average daily social interaction score and average daily loneliness level by frailty status. This figure displays the median levels and range variability of both average daily social interaction score and average daily loneliness levels among participants, categorized into robust, pre-frail and frail groups.



This figure shows the average predictive marginal effects of sociodemographic characteristics, health status, global mild behavioral impairment symptom presence, and social functioning on the probability of being classified as robust, pre-frail, or frail. The predictive margins were calculated using multinomial logistic regression, with error bars representing 95% confidence intervals for each effect.



This figure shows the average predictive marginal effects of sociodemographic characteristics, health status, global mild behavioral impairment symptom severity, and social functioning on the probability of being classified as robust, pre-frail, or frail. The predictive margins were calculated using multinomial logistic regression, with error bars representing 95% confidence intervals for each effect.



Multimedia Appendixes

Compilation of Multimedia Appendices.

URL: <http://asset.jmir.pub/assets/4314f9a58d8c119a41593e890209530d.docx>

The timing of the four daily alarms.

URL: <http://asset.jmir.pub/assets/db1c6967e6dacc520f10ce9002171178.png>

First step to respond to the application.

URL: <http://asset.jmir.pub/assets/059d843bda6e02c2e27082721f2030e0.png>

Second step to respond to the application.

URL: <http://asset.jmir.pub/assets/cc097e7dff6796fb77521f3d7f79419.png>

Distribution of mean maximum daily social interaction scores by frailty status, based on multiple daily scores for each participant. The black histogram represents robust participants, the grey histogram represents pre-frail group, and the blue histogram represents frail group. Participants recorded the number of face-to-face meetings, phone calls, or video calls lasting more than five minutes using a five-point Likert scale (0 for “no contact,” 1 for “once,” 2 for “two times,” 3 for “three times,” and 4 for “four or more times”).

URL: <http://asset.jmir.pub/assets/16e81408a2493a8108c67c546505ad4c.png>

Distribution of mean maximum daily loneliness levels by frailty status, based on multiple daily scores for each participant. The black histogram represents robust participants, the grey histogram represents pre-frail group, and the red histogram represents frail group. Participants assessed loneliness using a five-point Likert scale (0 = “very lonely” to 4 = “not lonely at all”), with higher scores indicating lower levels of loneliness.

URL: <http://asset.jmir.pub/assets/6eecee2e2fa912604ac616f79f722593.png>