

Factors Influencing Information Distortion in Electronic Nursing Records: a Qualitative Study

Jianan Wang, Yihong Xu, Zhichao Yang, Jie Zhang, Xiaoxiao Zhang, Wen Li,
Yushu Sun, Hongying Pan

Submitted to: Journal of Medical Internet Research
on: September 27, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 4

Supplementary Files..... 21

 Figures 22

 Figure 1..... 23

Multimedia Appendixes 24

 Multimedia Appendix 1..... 25

 Multimedia Appendix 2..... 25

 Multimedia Appendix 3..... 25

Factors Influencing Information Distortion in Electronic Nursing Records: a Qualitative Study

Jianan Wang^{1*} BS; Yihong Xu^{1*} MS; Zhichao Yang¹ BS; Jie Zhang¹ BS; Xiaoxiao Zhang¹ BS; Wen Li¹ BS; Yushu Sun¹ MS; Hongying Pan¹ MS

¹Department of Nursing Sir Run Run Shaw Hospital Zhejiang University School of Medicine Hangzhou CN

*these authors contributed equally

Corresponding Author:

Hongying Pan MS

Department of Nursing

Sir Run Run Shaw Hospital

Zhejiang University School of Medicine

3 Qingchun East Road

Hangzhou

CN

Abstract

Background: Information distortion in nursing records pose significant risks to patient safety and impede the enhancement of care quality. The introduction of information technologies, such as decision support systems and predictive models, expands the possibilities for utilizing health data but also complicates the landscape of information distortion. Only through identifying influencing factors about information distortion, can care quality and patient safety be ensured.

Objective: This study aims to explore the factors influencing information distortion in electronic nursing records within the context of China's healthcare system, and provide appropriate recommendations to address these distortions.

Methods: This qualitative study utilized semi-structured interviews conducted with 14 nurses from a Class-A tertiary hospital. The data were analyzed through qualitative content analysis.

Results: The analysis identified 4 categories and 10 sub-categories: (1) Nurse-related factors—skills, awareness, and work habits; (2) Patient-related factors—willingness and ability; (3) Operational factors—work characteristics and system deficiencies; (4) Organizational factors—management system, organizational climate, and team collaboration.

Conclusions: While some factors influencing information distortion in electronic nursing records (ENRs) are similar to those observed in paper-based records, others are unique to the digital age. As healthcare continues to embrace digitalization, it is crucial to develop and implement strategies to mitigate information distortion. Regular training and education programs, robust systems and mechanisms, as well as optimized human resources and organizational practices, are strongly recommended.

(JMIR Preprints 27/09/2024:66959)

DOI: <https://doi.org/10.2196/preprints.66959>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

✓ **No, I do not wish to publish my submitted manuscript as a preprint.**

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#), I will be able to make the full manuscript visible to all users.

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Original Paper

Jianan Wang, BS, PhD candidate ^{1*}, Yihong Xu, MS ^{1*}, Zhichao Yang, BS ¹, Jie Zhang, BS ¹,
Xiaoxiao Zhang, BS ¹, Wen Li, BS ¹, Yushu Sun, MS ¹, Hongying Pan, MS ¹

1.Nursing Department, Sir Run Run Shaw Hospital, Zhejiang University School of medicine, Hangzhou, China

* These authors contributed equally to this work.

Corresponding Author:

Hongying Pan, MS

Nursing Department, Sir Run Run Shaw Hospital

Zhejiang University School of medicine

3 Qingchun East Road

Hangzhou, 310016

China

Phone: 86 13857188922

Email: 3191016@zju.edu.cn

Factors Influencing Information Distortion in Electronic Nursing Records: a Qualitative Study

Abstract

Background: Information distortion in nursing records pose significant risks to patient safety and impede the enhancement of care quality. The introduction of information technologies, such as decision support systems and predictive models, expands the possibilities for utilizing health data but also complicates the landscape of information distortion. Only through identifying influencing factors about information distortion, can care quality and patient safety be ensured.

Objective: This study aims to explore the factors influencing information distortion in electronic nursing records within the context of China's healthcare system, and provide appropriate recommendations to address these distortions.

Methods: This qualitative study utilized semi-structured interviews conducted with 14 nurses from a Class-A tertiary hospital. Participants were primarily asked about their experiences and observations of information distortion in clinical practice, as well as potential influencing factors and corresponding countermeasures. Data were analyzed using inductive content analysis, which involved initial preparation, line-by-line coding, the creation of categories, and abstraction.

Results: The analysis identified 4 categories and 10 sub-categories: (1) Nurse-related factors—skills, awareness, and work habits; (2) Patient-related factors—willingness and ability; (3) Operational factors—work characteristics and system deficiencies; (4) Organizational factors—management system, organizational climate, and team collaboration.

Conclusions: While some factors influencing information distortion in electronic nursing records (ENRs) are similar to those observed in paper-based records, others are unique to the digital age. As healthcare continues to embrace digitalization, it is crucial to develop and implement strategies to mitigate information distortion. Regular training and education programs, robust systems and mechanisms, as well as optimized human resources and organizational practices, are strongly recommended.

Keywords: information distortion; electronic health record; qualitative research; ethics

Introduction

The use of digital technologies in healthcare is improving human life quality, making medical services easier, safer, and more accessible [1]. However, this socio-technical transformation has also introduced more complex and diverse ethical concerns [2]. Examples of information ethics issues include information disclosure, distortion, alienation, and injustice, etc [3]. Among these, information distortion undermines data accuracy, posing immediate health risks and directly threatening patient safety, which demands significant attention [3]. While the definitions of information distortion vary across different research areas, all studies highlight its common theme: the alteration of information—whether through exaggeration, misinterpretation, or other means—making it a critical data quality issue [4-6]. There are a few research focusing on data quality, even though they don't pinpoint the word “distortion”, they refer to “accuracy” or “correctness”, which relate to distortion [7,8]. However, these research tend to provide only a general overview of information distortion, treating it as a subset of data quality issues without exploring it in depth. Compared to other quality issues like redundancy or inaccessibility, information distortion could have more severe implications. Inaccurate health data can hinder preventive and therapeutic interventions, potentially compromising patient health outcomes [9]. Research has shown that every 1% reduction in data completeness is associated with a 1.21% increase in missing events [10], the management of which may impose additional burdens in terms of manpower, materials, and finances [11,12]. As healthcare professionals increasingly rely on electronic medical records (EMRs) to support patient care, the consequences of information distortion become even more pronounced in the digital era. Technology-driven tools, such as clinical decision support systems and algorithms, depend on accurate EMR data to provide reliable guidance and improve health. However, inaccuracies in data can lead to inappropriate medical decisions being perpetuated across multiple patients [13,14]. Furthermore, predictive models built on flawed data may introduce biases, adversely affecting entire populations on a large scale [9]. High-quality data are also indispensable for policymakers, who rely on EMR data for informed decision-making and strategic planning [15].

Research has evaluated the quality of health data in medical documentation, highlighting issues such as accuracy, completeness, and readability, which often intersect with information distortion [16,17]. With similar functions, nursing documentation is also a crucial component of health data quality, serving as a record of a patient's health data and the care provided [18]. High-quality nursing documentation is essential not only as a tool for nurses to support decision-making and ensure continuity of care but also as a critical reference for doctors to understand the patient's condition and take appropriate medical actions [19]. Accurate and comprehensive documentation is a legal and ethical obligation, and also a professional requirement for nurses [20,21]. However, persistent information distortion impedes accurate record-keeping. Information distortion occurs when nursing records fail to accurately reflect nursing practices or the patient's true condition [3]. Several studies have found that paper-based nursing records often inadequately reflect nursing activities [22,23]. In the current information era, research from Italy still that only 37% of assessments and 45% of interventions are documented in nursing records [24]. Similarly, Korean scholars [25,26] found that certain nursing activities, such as nurse rounding and skin assessments, are often performed but not recorded in electronic nursing records (ENRs). The correlation score between nursing activities and corresponding records was 3.08 on a 4-point scale. Our study (currently under review) further investigates the manifestations of information distortion in ENRs through a qualitative approach. The findings reveal that information distortion permeates various aspects of ENRs, including assessments, care planning, interventions, incident management, billing, and more. This is particularly concerning, as certain decision-making processes cannot tolerate inaccurate information, which may lead to severe consequences. Given the widespread occurrence of information distortion, understanding the factors influencing this phenomenon is crucial for developing effective

interventions to mitigate it. Previous studies have explored contributors to information distortion in paper-based records [22,23], identifying issues such as time constraints and cumbersome charting formats. However, the advent of technology has introduced structural changes to documentation practices, like digital workflows, predefined templates, and integrated decision-support tools. These innovations may alter the factors affecting distortion in ENRs due to variations in processes such as data entry, storage, updating, analysis, and auditing. And while factors affecting the quality of both formats of nursing records—such as heavy workloads, nurse fatigue, and hardware shortages—have been studied [27,28], information distortion phenomena manifest in specific and concerted ways within clinical contexts. These phenomena are not merely errors or omissions; they may arise from complex psychological, organizational, or systemic factors that are difficult to fully uncover in broader quality studies. As no previous research has independently examined the factors associated with information distortion in ENRs, the aim of this study was to investigate the factors contributing to information distortion phenomena in ENRs in China. Our results will help identify potential risks associated with ENRs and contribute to the development of targeted preventive strategies to reduce information distortion.

Methods

Study Design

We chose a qualitative descriptive approach for this study, as it is well-suited for obtaining direct insights from participants about a poorly understood phenomenon [29,30]. Semi-structured interviews were conducted to facilitate in-depth discussions of participants' perspectives. The study adhered to the Consolidated Criteria for Reporting Qualitative Research guidelines [31](Multimedia Appendix 1).

Setting

This study was conducted in a Class-A tertiary hospital in Zhejiang Province, China. The hospital has been utilizing electronic documentation for over 20 years, and integrated nursing decision support system (NDSS) into its nursing information system (NIS) in 2016. The hospital's information system was independently developed in-house, tailored specifically to meet the institution's unique operational needs and workflows, ensuring full control over system customization, integration, and ongoing updates. The hospital's advanced level of digitalization has provided nurses with a comprehensive technological environment, giving them extensive experience with the system, which is invaluable for gaining insights into issues related to information distortion.

Ethics Approval

Approval was obtained from the Ethics Committee of Sir Run Run Shaw Hospital, Zhejiang University School of Medicine (approval number: 2024 Research No. 0270). Written informed consent was obtained from all participants by the author (JW) prior to data collection. No participants opted out of the study. All data transcripts for analysis were deidentified.

Participants and Recruitment

Both purposive and snowball sampling methods were employed. Purposive sampling is a common technique in qualitative descriptive studies, selecting individuals who have relevant experience with the phenomenon of interest [32]. To ensure interviewees provide valuable insights into the nuances of information distortion, we decided to select members of the Information Committee as participants. Each department has a representative in the committee. The committee members are nurses actively involved in clinical practice who also take on additional responsibilities, such as addressing nurses' concerns about system usage and sharing technical knowledge. Given their roles,

these nurses are more likely to have encountered challenges and ethical dilemmas related to electronic documentation. Their insights are invaluable for uncovering the underlying causes of information distortion, which may not be as evident to nurses with less experience in this area. Considering that the information systems vary across different specialties, we conducted maximum variation sampling based on department categories after confirming members' willingness to participate to ensure that the selected participants encompass as many aspects and dimensions of clinical nursing work as possible. Participants were not only asked about their own experiences but also about the experiences of their colleagues they had observed. This approach allowed for the inclusion and comparison of diverse perspectives, making the insights into factors influencing distortion in ENRs more representative of the broader nursing community. Snowball sampling involves participants recommending additional subjects from their acquaintances, thereby continuously introducing new potential participants. We invited information committee members to introduce nurses who had been involved in clinical informatics-related work or projects as interviewees. Targeted nurses were invited through the hospital's internal network. Inclusion criteria for nurses were: (1) Registered nurses employed at the hospital for at least one year and (2) voluntary participation with signed informed consent. Nurses who were absent during the study period (e.g., on leave, vacation, or away for training) were excluded. Nurses from other hospitals who were present for training were also excluded, but participant nurses were encouraged to share relevant experience from their previous employment at other hospitals. Recruitment continued until data saturation was achieved, meaning no new substantive themes emerged from additional data collection [33].

Data Collection

Face-to-face interviews were conducted in the coffee shop or the demonstration classroom in the hospital between January 9, 2024, and April 20, 2024, at times convenient for the participants. The interviews were conducted by one author (JW), a female PhD student trained in qualitative research methods. Only the participants and the interviewer were present during the interviews. The interviewer had no prior relationship with the participants, which helped minimize biases and address potential ethical concerns. Before the interviews began, the interviewer explained the concept of information distortion and the purpose of the study to ensure that participants had the necessary background to provide informed and accurate responses. Informed consent materials were provided, and written consent was obtained from all participants.

As part of the interview process, participants completed a short questionnaire about their personal characteristics, including gender, age, education, post, professional title, work experience, and work department. A topic guide with open-ended questions (see Multimedia Appendix 2) was used to ensure comprehensive coverage of relevant topics during the interviews. The interview questions were developed by reviewing existing literature and absorbing expert opinions. To ensure validity, the guide was pre-tested by two nurses and revised based on their feedback. Participants were initially asked to describe instances of information distortion they had encountered or observed in clinical settings. Subsequently, they were prompted to identify factors contributing to these phenomena. Follow-up questions were tailored to participants' responses to encourage deeper elaboration. The next section focused on coping strategies to mitigate information distortion. During this discussion, participants often provided additional insights into influencing factors. Finally, participants were invited to share any additional thoughts or address overlooked aspects before concluding the interview. The interviewer took field notes during the interviews to supplement the data and highlight key moments [34]. By the 14th interview, no new insights emerged, and data began to repeat, indicating that further data collection was unnecessary. Thus, data saturation was achieved after 14 interviews. Interviews lasted between 35 and 97 minutes. All interviews were audio-recorded with participants' permission, transcribed verbatim, and checked by participants. No repeat interviews were conducted.

Data Analysis

Inductive qualitative content analysis was used for data analysis, encompassing three phases: preparation, organizing, and reporting [35], in which codes and categories were directly drawn from the data [36]. The recorded interviews were transcribed by one author (JW) and then verified by the participants. The preparation phase starts with selecting the unit of analysis [35]. Since the aim of this study is to explore influencing factors of information distortion, we especially considered participants' words or a set of sentences and paragraphs addressing influencing factors as meaning units. Two researchers (JW and YX) read all relevant data repeatedly to achieve immersion and gain an overall understanding of the content. The organization phase includes open coding, creating categories and abstraction [35]. The same two researchers started open coding by independently reading transcripts line by line, extracting texts with similar meanings and grouping them into fewer content-related codes to identify influencing factors [37]. The researchers also made notes of their initial impressions and thoughts regarding the codes. The codes were collected to the coding sheets. Then, the two synthesized their coding sheets and independently sorted them into more formalized subcategories, using quotations derived from the interviews. Next, they compared each subcategories, merging those with same meanings and discussing those with differences. Discrepancies that could not be resolved through discussion were addressed within the research group until a consensus was reached. Rearranging subcategories also occurred in this process. Finally, based on the relationships between subcategories, researchers collaboratively grouped a larger number of subcategories into a smaller number of categories. All data management and analyses were conducted in NVivo 12 (QSR International, Melbourne, Australia). The interviews were conducted in Mandarin, and only the directly quoted passages were translated into English for inclusion in the paper. The translation was performed by a professional translator fluent in both languages. The translation was reviewed by the research team to ensure the meaning of the original Mandarin quotes was faithfully preserved in the English translation.

Rigor

The rigor of this study was ensured by credibility, transferability, dependability, and confirmability [38]. Credibility was ensured by having two researchers independently code the data, bracketing personal biases. Furthermore, member checks were conducted by inviting participants to review and provide feedback on generated categories and subcategories. Transferability was ensured by providing thick description of the participants' inclusion and exclusion criteria, data collection and analysis procedures, and participant characteristics, enabling readers to relate findings to their own contexts. Dependability was ensured through the detailed methodological documentation, where all headings, subcategories, and categories were documented at each step, allowing others to assess the process. Confirmability was ensured through maintaining a journal to track researchers' evolving thoughts, biases, and reflections. The final categories and subcategories were reviewed by the entire research team to ensure coherence, which also increases the objectivity and confirms the accuracy of the findings.

Results

Participants

The demographic characteristics of the participants are presented in Table 1. We interviewed a total of 14 nurses, including clinical nurses, advanced practice nurses, and nurse educators. In terms of professional title, 3 participants are primary nurses, while 11 are charge nurses. All participants provide direct patient care, with charge nurses also responsible for overseeing junior nurses, mentoring interns, trainees, and new colleagues. Their working departments varies from intensive care unit, surgery, and internal medicine. All interviewees were female, aged 26-37 years, with work

experience ranging from 4 to 14 years. Most of them (n=12) have bachelor's degree, and 2 of them have master's degree.


Table1. Characteristics of participants (N=14).

Characteristic	Values, N (%)
Gender	
female	14 (100)
Age (years)	
25-29	3 (21)
30-34	7 (50)
35-39	4 (29)
Education	
Bachelor's degree	12 (86)
Master's degree	2 (14)
Post	
Clinical Nurse	8 (57)
Advanced Practice Nurse	3 (21)
Nurse Educator	3 (21)
Professional title	
Primary Nurse	3 (21)
Charge Nurse	11 (79)
Work experience (years)	
1-5	2 (14)
6-10	8 (57)
11-15	4 (29)
Work department	
Intensive care unit	1 (7)
General Surgery	3 (21)
Medical Oncology	1 (7)
Respiratory Medicine	1 (7)
General Internal Medicine	1 (7)
Endocrinology	1 (7)
General Ward	1 (7)
Cardiac Surgery	1 (7)
Rheumatology and Immunology	1 (7)
Neurosurgery	1 (7)
Neurology	2 (14)

Factors Influencing Information Distortion Behaviors

We generated 4 categories and 10 sub-categories about factors influencing information distortion phenomena in ENRs through the interviews(see Figure 1) . Illustrative quotes from participants are provided in the Multimedia Appendix 3.

Figure 1. Categories of influencing factors.



Nurse-related Factors: Skills

The lack of certain skills may contribute to nurses' information distortion phenomena. With the advancement of technology, such as NDSS and the copy-and-paste function (CPF), nurses may rely too heavily on system-generated advice or records from colleagues without critical evaluation, increasing the likelihood of distorted records. Additionally, a lack of multitasking and time management skills, particularly among junior nurses, can make them "busier and slower at handling issues", leading to omissions or errors in their ENRs that fail to accurately reflect patients' conditions. Nurses also indicated that job familiarity influences these phenomena. New nurses, unfamiliar with their duties, are more prone to making mistakes in their records, while experienced nurses, who may be overconfident in their abilities, might neglect to recheck their entries, resulting in inaccuracies. Furthermore, the introduction of new system designs can also lead to errors in records due to unfamiliarity.

Nurse-related Factors: Awareness

Work attitude plays a significant role in influencing information distortion phenomena. Nurses who are meticulous in their work tend to produce more accurate records, while those who are more casual and indifferent are more likely to write distorted records. The perception of information importance also affects the accuracy of their ENRs. Nurses reported that they pay more attention to critical aspects related to patient safety, frequently audited items, and department-specific issues. These areas are documented more carefully, resulting in fewer inconsistencies. However, minor issues not directly related to a patient's illness, such as a small skin condition, may go unrecorded even if noticed. Additionally, perceived threats, such as fear of medical disputes or judgment from others, can influence the occurrence of distortion phenomena. In terms of medical disputes, nurses indicated two possible reactions: they might document all nursing practices in ENRs to verify their workload and reduce disputes, which is a great phenomenon, or they might falsify records to appear perfect, given that ENRs serve as the primary evidence of their work. Concerns about others' judgment, particularly regarding incident reporting, also play a role. Nurses may worry about negative repercussions, thinking, "Will the head nurse blame me?" or "How could you make such a basic mistake?" As a result, they might cover up mistakes to protect their image. Furthermore, due to easily accessible entries to their colleagues' records given by digital technologies, nurses may give in to the professionalism or authority, leading to potential information distortion. Nurses might feel less confident in recording their true observations if more experienced colleagues have documented

differently, or they might question their own judgments when comparing their records to those of others so that they might heal their real thoughts.

Nurse-related Factors: Work Habits

Specific work habits can contribute to information distortion. Many nurses use CPF to enhance work efficiency. Some nurses prefer to copy records from the previous shift's and then adjust them after completing their nursing activities. However, this habit can lead to forgetting to make the necessary adjustments, resulting in identical records across shifts. Additionally, when nurses do not take the time to review their records, errors may go unnoticed. Nurses also mentioned that fixed work habits can pose a problem. When unexpected events occur, they might be omitted from the records simply because they fall outside of the usual routine.

Patient-related Factors: Willingness

Patients are a primary source of information, and much of the content in ENRs is based on their complaints. In clinical practice, patient-reported information is typically reflected accurately in ENRs. However, when patients are uncooperative, such as by intentionally withholding information or being unwilling to share, nurses may face challenges in accurately assessing their condition.

Patient-related Factors: Ability

Even when patients are highly willing to cooperate, their lack of certain objective abilities can still hinder the accuracy of information. Nurses mentioned difficulties with patients who speak dialects or foreign languages, where communication often becomes challenging and prone to misunderstandings, leading to more frequent instances of distorted records. In addition, cognitive and comprehension abilities can also affect the accuracy of ENRs. Cognitive decline is common among elderly patients and hinders effective communication. Under this circumstance, if family members are not well-informed about the patient's condition, nurses may face challenges in obtaining accurate information, potentially resulting in distorted records.

Operational Factors: Work Characteristics

Nurses face a well-documented heavy workload. The implementation of information systems and electronic records is designed to streamline clinical workflows and enhance efficiency [39]. However, interviews revealed that nurses continue to feel burdened by both documentation and clinical duties. As a result, they often complete their documentation hastily, leading to potential omissions and inaccuracies. Due to their heavy workload, nurses also admitted to using shortcuts in documentation further compromising the accuracy of their records. Certain characteristics of nursing records also contribute to information distortion. For instance, forms assessed frequently, such as critical care forms updated hourly, tend to be more accurate than those reviewed only once daily, as frequent access facilitates the identification and correction of mistakes. Moreover, assessments with subjective criteria are more prone to discrepancies, as individual nurses may have differing interpretations, leading to a lack of consensus on the patient's condition. Frequent interruptions also exacerbate information distortion, as nurses must address these issues before returning to their documentation, potentially resulting in inaccuracies in both timing and content.

Operational Factors: System Deficiencies

Nurses reported that inadequate information system design hinders accurate documentation. Limited system options and insufficient space for remarks often prevent them from fully capturing their observations. Even with remark sections, redundant entries increase the risk of distortion. Additionally, poor integration between the clinical and nursing information system, and between systems in the emergency room and wards, hampers the properly displayed and submission of information, making it difficult for ENRs to reflect the actual situation accurately.

Organizational Factors: Management System

Nursing leadership strongly influences staff behaviors [40]. Nurses noted that their leaders' focus impacts how carefully they document. When leaders stress the importance of accurate ENRs, nurses become more meticulous, reducing distortion. Direct leaders, who "focus on specific areas," were seen as more influential than higher-level leaders. Hospital regulations also impact documentation. Nurses reported that mismatch between regulatory requirements and clinical realities often lead them to adjust records. Areas without strict rules or audits receive less attention, as inconsistencies are seen as acceptable. In addition, the audit system has a dual impact on information distortion. The National Healthcare Security Administration enforces regulations to ensure the proper use of medical insurance funds and conducts irregularly electronic audits of records. Through digital technologies, these audits are conducted more frequently, discreetly, and cost-effectively. While this system encourages detailed and accurate documentation to meet compliance standards, it can also lead some nurses to falsify records to secure higher reimbursements.

Organizational Factors: Organizational Climate

The organizational atmosphere strongly influence nurses' behaviors, particularly through conformity. Nurses interviewed indicated that even if they recognized information distortion, they would gradually take it less seriously if the organization did not prioritize the issue. Conversely, in organizations where accuracy is emphasized, nurses collectively focus more on precise ENRs. Additionally, some nurses admitted to distorting records to protect the organization's reputation, either by avoiding incident reports or shifting blame to other departments to lower the rate of adverse events. Group pressure also plays a role, when confronted with numerous colleagues pointing out issues in their ENRs, they were more likely to yield and alter their records, even if they had originally documented the truth.

Organizational Factors: Team Collaboration

Patient care is definitely a teamwork across healthcare professionals [41]. And the collaboration between multiple labor could also impact information distortion. Nurses reported that their communication with doctors plays a role in affecting the accuracy of ENRs. They indicated that doctors often communicate directly with patients rather than with them, leaving nurses unaware of the patients' true condition. And when nurses ask patients about what the doctor said, it may lead patients to question the nurses' professionalism, preventing nurses from obtaining accurate information at that moment. Additionally, nurses were often hesitant to address questionable orders from doctors, either due to concerns about overstepping boundaries or for inconvenience. Consequently, nurses might record inaccurate ENRs based on these orders or ignore them entirely, leading to inconsistencies with reality. Furthermore, doctors' occasionally review nursing records and point out documentation issues, can act as a deterrent, prompting nurses to prioritize accuracy in their documentation. Nurses have noted that electronic formats make these reviews occur more frequently. Some nurses also acknowledged intentionally underrating patients' conditions to draw more attention from doctors, thereby enhancing patient safety.

Similar patterns were observed in collaboration with nursing colleagues; for instance, failing to mark a care plan as completed was sometimes used as a strategy to maintain colleagues' focus. Additionally, interactions with nursing assistants can also affect information distortion. If nursing assistants lack adequate expertise or effective communication with nurses, misunderstandings about the patient's condition may arise, resulting in inaccurate ENRs.

Discussion

Principal Findings

Using a qualitative approach, we identified various factors contributing to information distortion in ENRs. It is acknowledged that information distortion is not limited in ENRs, similar issues have been observed in general EMRs [16,17]. However, our findings reveal specific contents in nursing context. Some distortions have existed in nursing records long before the advent of information technologies [22,23], and some influencing factors remain constant across paper-based and digital records. However, with the introduction of digital technologies, the phenomenon has different manifestations, and some factors influencing are unique to the digital era.

First, while technological advancements provide conveniences, like the application of NDSS enhances and accelerates professional decision-making, they can also increase nurses' laziness, leading nurses to rely more on system-generated advice rather than their own judgment. Noting that data-driven tools cannot be accurate all the time, the over reliance of these can increase more distortion. Second, technologies also bring changes to nurses' work habits, such as copy and paste, although it is time-saving, it can exacerbate record distortion if nurses do not monitor closely. Third, compared with paper-based records, the writing of ENRs happens through information systems, which means that the design of information system play a key role in documentation. A poorly designed one may limit nurses' options, introduce redundant entries, or obstructing information exchange, which all potentially hindering nurses recording accurately and worsening information distortion. Fourth, digital records have made entries more accessible, with just a few clicks, users could get targeted information, not only their own records but others. With the increasing interaction or comparison with others' records, the accuracy of records might be doubted especially for those who are less confident or give way to authority. However, electronic formats also improve records quality in some ways. This kind of operation promises consistent check, and makes records auditing easier, when utilized effectively, this can enhance supervision and reduce information distortion.

Therefore, digital technologies can be viewed as a double-edged sword in relation to information distortion. Given that digitalization is inevitable, it is crucial to develop strategies to address these issues. We propose several measures to enhance the quality of ENRs.

First, regular training should focus on critical thinking, multitasking skills, new procedures, and onboarding. This study shows that skill gaps in this areas lead to record distortion, task delays, and poor performance [42,43]. Targeted improvement initiatives are crucial. Moreover, communication training, emphasizing empathy and nonverbal skills, should be conducted to build trust and improve the accuracy of patients information. Teamwork training across healthcare professionals, is equally important for maintaining accurate records and fostering a collaborative environment. Educational programs should also highlight the consequences of medical disputes and the importance of accurate records. Continuing education should extend to both nurses and nursing assistants to enhance knowledge, confidence, and collaboration.

Second, certain mechanisms should be improved. For example, routine file inspections conducted by administrators, along with penalties for falsifying records, can serve as deterrents against misconduct [44]. Providing patients access to their records (e.g., via Open Notes [45]) could also enhance the accuracy of documentation, as patients can help identify errors. Additionally, establishing communication and feedback system could address usability issues in NIS and resolve any regulatory concerns. Research has shown that a positive work attitude influences the quality of nurses' documentation [46]. Therefore, implementing recognition programs to highlight exemplary practices is recommended to improve nurses' attitudes toward record-keeping and reduce distortion.

Third, given the pivotal role of information technologies in nursing, tools like translation services and wearable devices can help minimize patients communication barriers during data collection[38]. Artificial intelligence can improve ENR quality by using machine learning to detect record errors and large language models to analyze clinical text, ensuring data completeness and accuracy [47-49]. Additionally, information systems must be updated to align with clinical requirements. Enhancing interoperability will facilitate cross-professional checks, reducing uncertainties and omissions in ENRs [50]. Informatics nurses should promote collaboration between technicians and clinical staff to enhance the usability of intelligent devices and information systems, mitigating technology-related challenges [51].

Fourth, heavy workload significantly impacts the quality of ENRs. High patient-to-nurse ratios compel nurses to complete tasks, including their documentation, more hastily [52]. Clinical nurses spend 17-27% of their time on ENR-related tasks, surpassing the time allocated to direct patient care or communication [53,54]. Therefore, increasing nurse staffing and simplifying documentation processes are essential to reduce their burden [55,56]. This would facilitate more thorough documentation and decrease the likelihood of information distortion. In addition, the use of medical scribes in the USA—unlicensed individuals who enter data into EMRs under clinicians' supervision—has been growing [57]. These scribes can also assist with ENRs, helping to reduce nurses' documentation workload and allowing them to focus more on maintaining the accuracy of records.

Finally, management practices are closely associated with organizational procedures, and effective leadership serves as a potential catalyst for fostering positive team behaviors [58,59]. Therefore, it is essential that leaders place greater emphasis on addressing information distortion issues, as this will influence staff attitudes. Leaders should also promote an open and positive organizational climate by establishing core values and behavioral guidelines that emphasize trust, learning, and accountability, while reducing feelings of blame or fear [60]. The role of direct leaders, as identified in this study, is essential to the whole team. Therefore, they are responsible for not only conveying information from upper management to the nursing staff but also for leading by example, embodying the values and behaviors expected within the team.

Limitations

This study has several limitations. First, the numbers of interviewees may appear small. However, saturation was reached, and this sample size aligns with the typical saturation range of 12–20 interviews, as suggested by Guest et al. [61]. Second, interviewees were female from one hospital, and only those with specific training in electronic notes or information technology were invited, this may have contributed to bias as findings from this study may not apply to other settings. However, given the predominance of female nurses in the Chinese workforce, the limitation of gender was inevitable. We used maximum variation to gain perspectives from various departments, which may help increase transferability. In addition, participants from Information Committee engage in more frequent discussions with other nurses about information systems, so their perspectives represent those commonly mentioned by most nurses in their departments. Furthermore, during the interviews, we also asked about their colleagues' experiences, allowing for the sharing and comparison of other nurses' opinions, which helps improve the generalizability of the findings. Third, although we discussed management strategies and leadership styles, we did not include nurse leaders in our interviews, which may have limited our understanding of how leadership affects information distortion. However, the aim of this study was to explore clinical nurses' perspectives about influencing factors of information distortion in ENRs, since nurse leaders in China do not participate in direct care, they are primarily concerned with macro-level management and policy-making rather than the detailed aspects of documentation behaviors, which may not fit for this study. Future research could incorporate the perspectives of male nurses and nurse leaders from more hospitals for

a more comprehensive view of influencing factors in information distortion.

Conclusions

We conducted a qualitative study using semi-structured interviews with Chinese nurses to explore the influencing factors of information distortion in ENRs. The analysis identified that these phenomena are driven by nurse-related, patient-related, operational, and organizational factors. To reduce the likelihood of information distortion, further development is needed in training, mechanisms, techniques, human resources, and organizational practices.

Acknowledgements

The authors would like to thank all participants for their time and contributions. This work was supported by Health Commission of Zhejiang Province (2024KY1142) and Science Technology Department of Zhejiang Province (2023C03191).

Data availability

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Conflicts of Interest

None declared.

Authors' Contributions

JW and YX conceptualized this research and designed the protocol; JW collected the data; JW and YX analyzed the data under HP's guidance and supervision; JW drafted the manuscript; All authors provided important intellectual content, reviewed the manuscript, and approved the final version of the manuscript.

Abbreviations

CPF: copy-and-paste function

EMR: electronic medical record

ENR: electronic nursing record

NDSS: nursing decision support system

NIS: nursing information system

Multimedia Appendix 1: [Consolidated Criteria for Reporting Qualitative studies (COREQ) checklist]

Multimedia Appendix 2: [Topic guide]

Multimedia Appendix 3: [Illustrative quotes from participants]

References

1. Ali O, Abdelbaki W, Shrestha A, Elbasi E, Alryalat MAA, Dwivedi YK. A systematic literature review of artificial intelligence in the healthcare sector: Benefits, challenges, methodologies, and functionalities. *J Innov Knowl* 2023 Jan;8(1):100333. [doi: 10.1016/j.jik.2023.100333]
2. Nittari G, Khuman R, Baldoni S, Pallotta G, Battineni G, Sirignano A, et al. Telemedicine Practice: Review of the Current Ethical and Legal Challenges. *Telemed J E Health* 2020 Dec;26(12):1427-1437. [doi: 10.1089/tmj.2019.0158]
3. Wang J, Xu Y, Zhang X, Pan H. Ethical predicaments and countermeasures in nursing informatics. *Nurs Ethics* 2024 Sep;31(6):1050-1064. [doi: 10.1177/09697330231215962]

4. Zhou Y, Fu Y, Wu X. Value analysis with blockchain-based information transparency system to eliminate information distortion. *Int J Prod Econ* 2023 Nov;265: 109008. [doi: 10.1016/j.ijpe.2023.109008]
5. Boyle PJ, Purdon M. The information distortion bias: implications for medical decisions. *Med Educ* 2019 Nov;53(11):1077-1086. [doi: 10.1111/medu.13919]
6. Jun SY, Ju TW, Park HK, Lee JC, Kim TM. Information distortion in word-of-mouth retransmission: the effects of retransmitter intention and source expertise. *Asian Bus Manag* 2023;22(5):1848-1876. [doi: 10.1057/s41291-023-00221-w]
7. Makeleni N, Cilliers L. Critical success factors to improve data quality of electronic medical records in public healthcare institutions. *S Afr J Inf Manag* 2021;23(1):1-8. [doi: 10.4102/sajim.v23i1.1230]
8. Syed R, Eden R, Makasi T, Chukwudi I, Mamudu A, Kamalpour M, et al. Digital Health Data Quality Issues: Systematic Review. *J Med Internet Res* 2023 Mar;25:e42615. [doi: 10.2196/42615]
9. Juhn YJ, Ryu E, Wi CI, King KS, Malik M, Romero-Brufau S, et al. Assessing socioeconomic bias in machine learning algorithms in health care: a case study of the HOUSES index. *J Am Med Inform Assoc* 2022 Jun;29(7):1142-1151. [doi: 10.1093/jamia/ocac052]
10. Johnson SG, Speedie S, Simon G, Kumar V, Westra BL. Quantifying the Effect of Data Quality on the Validity of an eMeasure. *Appl Clin Inform* 2017 Oct;8(4):1012-1021. [doi: 10.4338/ACI-2017-03-RA-0042]
11. Sood PD, Liu S, Lehmann H, Kharrazi H. Assessing the Effect of Electronic Health Record Data Quality on Identifying Patients With Type 2 Diabetes: Cross-Sectional Study. *JMIR Med Inform* 2024 Aug;12:e56734. [doi: 10.2196/56734]
12. Xu D, Sheng JQ, Hu PJ, Huang TS, Hsu CC. A Deep Learning-Based Unsupervised Method to Impute Missing Values in Patient Records for Improved Management of Cardiovascular Patients. *IEEE J Biomed Health Inform* 2021 Jun;25(6):2260-2272. [doi: 10.1109/JBHI.2020.3033323]
13. Hasan, S, Padman R. Analyzing the effect of data quality on the accuracy of clinical decision support systems: a computer simulation approach. *AMIA Annu Symp Proc*. 2006 Jan;2006:324-328. PMID: 17238356
14. Graber ML, Byrne C, Johnston D. The impact of electronic health records on diagnosis. *Diagnosis* 2017 Nov;4(4):211-223. [doi: 10.1515/dx-2017-0012]
15. Terry AL, Stewart M, Cejic S, Marshall JN, Lusignan SD, Chesworth BM, et al. A basic model for assessing primary health care electronic medical record data quality. *BMC Med Inform Decis Mak* 2019 Feb;19(1):30. [doi: 10.1186/s12911-019-0740-0]
16. Endriyas M, Kawza A, Alano A, Lemango F. Quality of medical records in public health facilities: A case of Southern Ethiopia, resource limited setting. *Health Informatics J* 2022 Jul-Sep;28(3):14604582221112853. [doi: 10.1177/14604582221112853]
17. Wu KA, Boccaccio K, Buckles D, Hartwig MG, Klapper JA. Efforts to improve the billing accuracy of robotic-assisted thoracic surgery through education, updated procedure cards, and electronic medical record system changes. *BMJ Open Qual* 2024 Apr;13(2): e002710. [doi: 10.1136/bmjopen-2023-002710]
18. McCarthy B, Fitzgerald S, O'Shea M, Condon C, Hartnett-Collins G, Clancy M, et al. Electronic nursing documentation interventions to promote or improve patient safety and quality care: A systematic review. *J Nurs Manag* 2019 Apr;27(3):491-501. [doi: 10.1111/jonm.12727]
19. Kebede M, Endris Y, Zegeye DT. Nursing care documentation practice: The unfinished task of nursing care in the University of Gondar Hospital. *Inform Health Soc Care* 2017 Sep;42(3):290-302. [doi: 10.1080/17538157.2016.1252766]
20. Goldsmith J. The NMC code: conduct, performance and ethics. *Nursing times*. 2011 Sep;107(37):12-14. PMID: 22010552
21. American Nurses Association. ANA's Principles for Nursing Documentation. Maryland. 2010.

- URL: <https://www.nursingworld.org/~4af4f2/globalassets/docs/ana/ethics/principles-of-nursing-documentation.pdf> [assessed 2025-01-15]
22. Inan NK, Dinc L. Evaluation of nursing documentation on patient hygienic care. *Int J Nurs Pract*. 2013 Feb;19(1):81-87. [doi: 10.1111/ijn.12030]
 23. Brooks JT. An analysis of nursing documentation as a reflection of actual nurse work. *Medsurg Nurs* 1998 Aug;7(4):189-198. PMID: 9782888
 24. De Marinis MG, Piredda M, Pascarella MC, Vincenzi B, Spiga F, Tartaglini D, et al. 'If it is not recorded, it has not been done!': consistency between nursing records and observed nursing care in an Italian hospital. *J Clin Nurs* 2010 Jun;19(11-12):1544-1552. [doi: 10.1111/j.1365-2702.2009.03012.x]
 25. Lee S, Yu S, Park H, Kim TG. Assessing How Well Electronic Nursing Records Reflect Changes in the Nursing Profession's Scope of Practice in Korea. *Comput Inform Nurs* 2020 Jun;38(6):312-319. [doi: 10.1097/CIN.0000000000000585]
 26. Park H, Yu S, Lee SH. Evaluating the Correlation Between Nursing Practice and Electronic Nursing Records Using Importance-Performance Analysis. *Comput Inform Nurs* 2021 Apr;39(9):492-498. [doi: 10.1097/CIN.0000000000000737]
 27. Ahn M, Choi M, Kim Y. Factors Associated with the Timeliness of Electronic Nursing Documentation. *Healthc Inform* 2016 Oct;22(4):270-276. PMID: 27895958
 28. Cheevakasemsook A, Chapman Y, Francis K, Davies C. The study of nursing documentation complexities. *Int J Nurs Pract* 2006 Dec;12(6):366-74. [doi: 10.1111/j.1440-172X.2006.00596.x]
 29. Sandelowski M. Whatever happened to qualitative description? *Res Nurs Health* 2000 Aug;23(4):334-340. [doi: 10.1002/1098-240x(200008)23:4<334::aid-nur9>3.0.co;2-g]
 30. Kim H, Sefcik JS, Bradway C. Characteristics of Qualitative Descriptive Studies: A Systematic Review. *Res Nurs Health* 2017 Feb;40(1):23-42. [doi: 10.1002/nur.21768]
 31. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007 Dec;19(6):349-57. [doi: 10.1093/intqhc/mzm042]
 32. Palinkas LA, Horwitz SM, Green CA, Wisdom JP, Duan N, Hoagwood K. Purposeful Sampling for Qualitative Data Collection and Analysis in Mixed Method Implementation Research. *Adm Policy Ment Health* 2015 Sep;42(5):533-544. [doi: 10.1007/s10488-013-0528-y]
 33. Morse JM. Designing funded qualitative research. *Handbook of qualitative research*. California: Sage Publications 1994:220-235.
 34. Phillippi J, Lauderdale J. A Guide to Field Notes for Qualitative Research: Context and Conversation. *Qual Health Res* 2018 Feb;28(3):381-388. [doi: 10.1177/1049732317697102]
 35. Elo S, Kyngas H. The qualitative content analysis process. *J Adv Nurs* 2008 Apr;62(1):107-15. [doi: 10.1111/j.1365-2648.2007.04569.x]
 36. Cho J, Lee E-H. Reducing Confusion about Grounded Theory and Qualitative Content Analysis: Similarities and Differences. *Qual Rep* 2014;19(64):1-20. [doi: 10.46743/2160-3715/2014.1028]
 37. Hsieh HF, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res* 2005 Nov;15(9):1277-1288. [doi: 10.1177/1049732305276687]
 38. Ahmed SK. The pillars of trustworthiness in qualitative research. *J Med Surg Public Health* 2024 Apr;2:100051. [doi: 10.31219/osf.io/se58y]
 39. Kelley TF, Brandon DH, Docherty SL. Electronic nursing documentation as a strategy to improve quality of patient care. *J Nurs Scholarsh* 2011 Jun;43(2):154-62. [doi: 10.1111/j.1547-5069.2011.01397.x]
 40. Zhou X, Wu Z, Liang D, Jia R, Wang M, Chen C, et al. Nurses' voice behaviour: The influence of humble leadership, affective commitment and job embeddedness in China. *J Nurs Manag* 2021 Sep;29(6):1603-12. [doi: 10.1111/jonm.13306]
 41. Schmutz JB, Meier LL, Manser T. How effective is teamwork really? The relationship between teamwork and performance in healthcare teams: a systematic review and meta-analysis. *BMJ*

- Open 2019 Sep 12;9(9):e028280. [doi: 10.1136/bmjopen-2018-028280]
42. Ates N, Erdal N, Harmanci Seren AK. The relationship between critical thinking and job performance among nurses: A descriptive survey study. *Int J Nurs Pract* 2023 Oct;29(5):e13173. [doi: 10.1111/ijn.13173]
 43. Westbrook JI, Woods A, Rob MI, Dunsmuir WT, Day RO. Association of interruptions with an increased risk and severity of medication administration errors. *Arch Intern Med* 2010 Apr 26;170(8):683-90. [doi: 10.1001/archinternmed.2010.65]
 44. Gibbs JP. Deterrence theory and research. *Nebr Symp Motiv* 1985;33:87-130. PMID: 3836360
 45. Bell SK, Folcarelli PH, Anselmo MK, Crotty BH, Flier LA, Walker J. Connecting Patients and Clinicians: The Anticipated Effects of Open Notes on Patient Safety and Quality of Care. *Jt Comm J Qual Patient Saf* 2015 Aug;41(8):378-384. [doi: 10.1016/s1553-7250(15)41049-9]
 46. Udaksana I, Supartha IWG, Ketut IB, Authority IMA. Influence Of Empowerment, Work Environment and Work Attitude on The Performance of Emergency Room Nurses. *Quali-Access Success* 2022 Apr;23(187):150-60. [doi: 10.47750/qas/23.187.19]
 47. Zimolzak AJ, Wei L, Mir U, Gupta A, Vaghani V, Subramanian D, et al. Machine Learning to Enhance Electronic Detection of Diagnostic Errors. *JAMA Netw Open* 2024 Sep 3;7(9):e2431982. [doi: 10.1001/jamanetworkopen.2024.31982]
 48. Ross A, McGrow K, Zhi D, Rasmy L. Foundation Models, Generative AI, and Large Language Models: Essentials for Nursing. *Comput Inform Nurs* 2024 May;42(5):377-387. [doi: 10.1097/CIN.0000000000001149]
 49. Xu Y, Ding S, Liu X, Pan H. The limitations of nursing decision support system and countermeasures. *Chin J Nurs* 2020;55(3):405-409. [doi: 10.3761/j.issn.0254-1769.2020.03.018]
 50. Bernardi FA, Alves D, Crepaldi N, Yamada DB, Lima VC, Rijo R. Data Quality in Health Research: Integrative Literature Review. *J Med Internet Res* 2023 Oct;25:e41446. [doi: 10.2196/41446]
 51. Kim L. Cybercrime, ransomware, and the role of the informatics nurse. *Nurs* 2020 Mar;50(3):63-65. [doi: 10.1097/01.NURSE.0000654064.67531.c5]
 52. Cho H, Nguyen O, Weaver M, Marcelle C, Pruitt J, Lee T, et al. Use of Electronic Health Records and Documentation Burden Among Nurses in Acute and Critical Care. *Stud Health Technol Inform* 2024 Jul 24;315:777-778. [doi: 10.3233/SHTI240325]
 53. Khan AR, Rosenthal CD, Ternes K, Sing RF, Sachdev G. Time Spent by Intensive Care Unit Nurses on the Electronic Health Record. *Crit Care Nurse* 2022 Oct 1;42(5):44-50. [doi: 10.4037/ccn2022518]
 54. Bakhoun N, Gerhart C, Schremp E, Jeffrey AD, Anders S, France D, et al. A Time and Motion Analysis of Nursing Workload and Electronic Health Record Use in the Emergency Department. *J Emerg Nurs* 2021 Sep;47(5):733-741. [doi: 10.1016/j.jen.2021.03.007]
 55. Kim Y, Lee MJ, Choi M, Cho E, Ryu GW. Exploring nurses' multitasking in clinical settings using a multimethod study. *Sci Rep* 2023 Apr 7;13(1):5704. [doi: 10.1038/s41598-023-32350-9]
 56. Cooper AL, Kelly MS, Brown JA. Co-design of an intervention to reduce the burden of clinical documentation: A clinician-researcher collaboration. *Appl Nurs Res* 2023 Oct;73:151730. [doi: 10.1016/j.apnr.2023.151730]
 57. Campbell LL, Case D, Crocker JE, Foster M, Johnson M, Lee CA, et al. Using medical scribes in a physician practice. *J AHIMA* 2012 Nov-Dec;83(11):64-69. PMID: 23210302
 58. Dys S, Tunalilar O, Hasworth S, Winfree J, White DL. Person-centered care practices in nursing homes: Staff perceptions and the organizational environment. *Geriatr Nurs* 2022 Jan-Feb;43:188-196. [doi: 10.1016/j.gerinurse.2021.11.018]
 59. Masood M, Afsar B. Transformational leadership and innovative work behavior among nursing staff. *Nurs Inq* 2017 Oct;24(4). [doi: 10.1111/nin.12188]
 60. Hamed MMM, Konstantinidis S. Barriers to Incident Reporting among Nurses: A Qualitative Systematic Review. *West J Nurs Res* 2022 May;44(5):506-523. [doi:

10.1177/0193945921999449]


61. Guest G, Bunce A, Johnson L. How Many Interviews Are Enough? Field Methods 2006 Feb;18(1):59-82. [doi: 10.1177/1525822x05279903]



Supplementary Files

Figures

Categories of influencing factors.



Multimedia Appendixes

Consolidated Criteria for Reporting Qualitative studies (COREQ) checklist.

URL: <http://asset.jmir.pub/assets/f9afe67310d19ea7412828e0cd81a0ee.docx>

Topic guide.

URL: <http://asset.jmir.pub/assets/8f6dca034179fa3fb11f95979655fd90.docx>

Illustrative quotes from participants.

URL: <http://asset.jmir.pub/assets/9a5b7c5b72f62127aab7290aedf0c8cf.docx>

