

Whole-body electromyostimulation (WB-EMS) training on glycemic control in people with prediabetes: a study protocol for a randomized controlled pilot-trial

Mahdieh Shojaa, Katharina Knaub, Norbert Schmitz, Andreas Michael Nieß, Barbara Munz, Sarah Rau, Viktoria Feit, Wallen Mphepo, Rahel Dingler, Wolfgang Kemmler

Submitted to: JMIR Research Protocols
on: November 13, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 20

 Figures 21

 Figure 1..... 22

 Figure 2..... 23

CONSORT (or other) checklists..... 24

 CONSORT (or other) checklist 0..... 25

 CONSORT (or other) checklist 0..... 25

Whole-body electromyostimulation (WB-EMS) training on glycemic control in people with prediabetes: a study protocol for a randomized controlled pilot-trial

Mahdieh Shojaa¹ PhD; Katharina Knaub¹ MSc; Norbert Schmitz¹ PhD; Andreas Michael Nieß² PhD; Barbara Munz² PhD; Sarah Rau¹ MPH; Viktoria Feit¹ MSc; Wallen Mphepo³ PhD; Rahel Dingler¹; Wolfgang Kemmler⁴ PhD

¹ Department of Population-Based Medicine University Hospital Tübingen Tübingen DE

² Department of Sports Medicine University Hospital Tübingen Tübingen DE

³ Evidence-Based Healthcare Programme University of Oxford Oxfordshire GB

⁴ Institute of Radiology Friedrich-Alexander-University of Erlangen-Nürnberg Erlangen DE

Corresponding Author:

Mahdieh Shojaa PhD

Department of Population-Based Medicine
University Hospital Tübingen
Röntgenweg 9
Tübingen
DE

Abstract

Background: Diabetes prevention programs focus on people with prediabetes because they have a greater risk of developing type 2 diabetes mellitus (T2DM) than people with normal blood glucose levels. Weight management can reduce this risk. However, in our largely sedentary society, there is less enthusiasm for regular exercise. Whole-body electromyostimulation (WB-EMS) is a training technology that provides exercise-like effects by inducing muscle contractions using electrical currents. There is evidence that local EMS can improve glucose metabolism. Several studies investigated the effect on WB-EMS on cardiometabolic risk factors including blood glucose control in a population of subjects with metabolic syndrome. However, to the best of our knowledge, there is no randomized controlled trial examining the preliminary efficacy of WB-EMS on hemoglobin A1c (HbA1c) levels in individuals with prediabetes.

Objective: The objective of this randomized controlled trial is to pilot procedures for a randomized controlled trial testing WB-EMS training on glycemic changes in sedentary adults with prediabetes.

Methods: Sixty community-dwelling sedentary adults aged 40-65 years with prediabetes will be randomized to one of three arms: WB-EMS + an activity tracker and a lifestyle education program (LEP) focusing on diabetes prevention, an activity tracker and LEP, or LEP only, with 20 subjects in each arm. The WB-EMS training will consist of 1.5×20 min per week. The intervention will last 16 weeks. As a pilot study, our main outcomes concern the number of participants who will be recruited, comply with intervention, and follow up. The primary efficacy outcome of interest includes HbA1c. The intention-to-treat analysis will be conducted with the objective of providing confidence interval estimation of treatment effects.

Results: The recruitment of study participants started in February 2024. As of October 2024, the recruitment was still ongoing. So far, 42 participants were allocated to the study groups. The anticipated date of recruitment completion is April 2025.

Conclusions: The results of this trial will provide valuable evidence for future investigations comparing the efficacy of the WB-EMS intervention with traditional exercise training to improve glycemic control in this population. Clinical Trial: Clinicaltrials.gov ID NCT06188481, registered December 7, 2023, <https://clinicaltrials.gov/study/NCT06188481>

(JMIR Preprints 13/11/2024:68761)

DOI: <https://doi.org/10.2196/preprints.68761>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <https://www.jmir.org/preprint/68761>

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in <https://www.jmir.org/preprint/68761>



Original Manuscript

Protocol for a randomized controlled trial

Whole-body electromyostimulation (WB-EMS) training on glycemic control in people with prediabetes: a study protocol for a randomized controlled pilot-trial

Mahdieh Shojaa ^{a,*}, Katharina Knaub ^a, Norbert Schmitz ^a, Andreas Michael Nieß ^b, Barbara Munz ^b, Sarah Rau ^a, Viktoria Feit ^a, Wallen Mphepo ^c, Rahel Dingler ^a, Wolfgang Kemmler ^d

Author affiliation:

^a Department of Population-Based Medicine, Institute of Health Sciences, University Hospital Tübingen, Tübingen, Germany

^b Department of Sports Medicine, University Hospital Tübingen, Tübingen, Germany

^c Evidence-Based Healthcare Programme, University of Oxford, Oxfordshire, England

^d Institute of Radiology, Friedrich-Alexander-University of Erlangen-Nürnberg, Erlangen, Germany

*Corresponding author:

Mahdieh Shojaa

University Hospital Tübingen, Institute of Health Sciences, Department of Population-Based Medicine

Röntgenweg 9

72076 Tübingen

Germany

E-mail: mahdieh.shojaa@med.uni-tuebingen.de

Tel.: 07071 29-88815

Abstract

Background: Diabetes prevention programs focus on people with prediabetes because they have a greater risk of developing type 2 diabetes mellitus (T2DM) than people with normal

blood glucose levels. Weight management can reduce this risk. However, in our largely sedentary society, there is less enthusiasm for regular exercise. Whole-body electromyostimulation (WB-EMS) is a training technology that provides exercise-like effects by inducing muscle contractions using electrical currents. There is evidence that local EMS can improve glucose metabolism. Several studies investigated the effect on WB-EMS on cardiometabolic risk factors including blood glucose control in a population of subjects with metabolic syndrome. However, to the best of our knowledge, there is no randomized controlled trial examining the preliminary efficacy of WB-EMS on hemoglobin A1c (HbA1c) levels in individuals with prediabetes.

Objectives: The objective of this randomized controlled trial is to pilot procedures for a randomized controlled trial testing WB-EMS training on glycemic changes in sedentary adults with prediabetes.

Methods: Sixty community-dwelling sedentary adults aged 40-65 years with prediabetes will be randomized to one of three arms: WB-EMS + an activity tracker and a lifestyle education program (LEP) focusing on diabetes prevention, an activity tracker and LEP, or LEP only, with 20 subjects in each arm. The WB-EMS training will consist of 1.5×20 min per week. The intervention will last 16 weeks. As a pilot study, our main outcomes concern the number of participants who will be recruited, comply with intervention, and follow up. The primary efficacy outcome of interest includes HbA1c. The intention-to-treat analysis will be conducted with the objective of providing confidence interval estimation of treatment effects.

Results: The recruitment of study participants started in February 2024. At the time of submission of this protocol for publication, the recruitment was still ongoing. As of October 2024, 42 participants were allocated to the study groups. The anticipated date of recruitment completion is April 2025.

Conclusion: The results of this trial will provide valuable evidence for future investigations comparing the efficacy of the WB-EMS intervention with traditional exercise training to improve glycemic control in this population.

Trial registration: Clinicaltrials.gov ID NCT06188481, registered December 7, 2023, <https://clinicaltrials.gov/study/NCT06188481>

Key words

Prediabetes, Whole-body electromyostimulation, Type 2 diabetes mellitus, Prevention

Introduction

Type 2 diabetes mellitus (T2DM) is a serious and widespread disease, and is recognized as a major public health challenge worldwide. According to the International Diabetes Federation (IDF), the current estimate of adults aged 20-79 years living with diabetes equals 9.3 % (463 million), and is projected to increase to 10.9% (700 million) by 2045 [1]. As the prevalence of T2DM increases, effective prevention measures are increasingly important. Focusing on those who are at high risk of developing T2DM is a pivotal starting point.

A state of glucose metabolism is called prediabetes when diabetes mellitus is not yet present but elevated fasting plasma glucose levels or impaired glucose tolerance are known [2]. HbA1c levels between 5.7% and 6.4% are indicative of prediabetes. The prevalence of prediabetes in Germany is estimated to be 26% in people aged 45 to 64 years and 31% in people aged 65 to 69 years [3]. Similarly, in the US, 21.7 % of people have prediabetes according to the HbA1c diagnostic criteria [4]. The clinical relevance of prediabetes is that there is a significantly increased risk of developing T2DM compared to the normoglycemic population, which in turn is associated with an increased risk of developing cardiovascular disease, kidney disease, cancer, depression and dementia [2]. The results from the population-based Rotterdam Study

suggest that up to 74% of 45-year-olds with prediabetes will develop T2DM during their lifetime [5].

The results from the KORA-F4 study (Cooperative Health Research in the Region of Augsburg) and the SHIP-TREND study (Study of Health in Pomerania) suggest that up to one third of people with prediabetes can be classified as physically inactive [6]. Lifestyle changes, such as physical activity and weight management, are of critical importance in preventing the development and progression of T2DM [2,7].

Although, physical activity (PA) interventions have shown beneficial effects on oral glucose tolerance, fasting glucose and HbA1c levels in people with prediabetes [8], insufficient PA in people with T2DM and prediabetes remains an issue in these populations [9]. There are several barriers associated with participation and long-term adherence to exercise programs among physically inactive people with diabetes, including lack of time, lack of motivation, poorer (self-rated) health, risk of injury, and social and environmental barriers [10]. Therefore, PA programs with an attractive profile are needed to reduce sedentary behavior and increase adherence to physical activity. A potential approach to achieve this goal is an innovative training technology called whole-body electromyostimulation (WB-EMS). WB-EMS provides exercise-like effects by inducing muscle contractions using electrical currents from an external source. WB-EMS simultaneously stimulates up to 8-12 major muscle groups with up to 2,800 cm² of electrode area. Electrodes are placed on the gluteus muscle, thighs, lower back, upper back, latissimus dorsi, abdomen, chest and upper arms, and each body region is stimulated with an individualized pulse intensity while individuals perform low-intensity functional exercises in a standing position [11].

In recent years, WB-EMS has found its way not only into therapeutic training, but also into rapid and widespread distribution, particularly in Europe, with more than 2,000 commercial WB-EMS providers serving approximately 250,000 clients in Germany alone [12]. Due to its increasing popularity, time efficiency, ease of implementation, joint-friendliness and personalized application, WB-EMS is increasingly the subject of scientific research. In fact, several studies have demonstrated positive effects of WB-EMS on muscle hypertrophy and body composition [13,14]. Additionally, high adherence to the intervention and low dropout rates were reported previously [15]. Therefore, this training technology can be considered a promising approach when aiming to increase PA in a population of sedentary adults.

A non-randomized study of WB-EMS showed benefits to HbA1c levels and fasting glucose in middle-aged and older men with T2DM [16]. In addition, several small studies have shown that local EMS has beneficial effects on glycemic control [17]. In contrast, a recent randomized crossover study by Holzer et al. found no significant short-term difference in the acute postprandial glucose response between exercise with and without WB-EMS. However, due to the small sample size (6 T2DM patients) and short intervention duration (three 20-minute sessions), the results of this study cannot be generalized [18]. Some studies further investigated the effect on WB-EMS in a population of subjects with metabolic syndrome. While prediabetes and metabolic syndrome share some common characteristics, people with metabolic syndrome have heterogeneous cardiometabolic profiles, where prediabetes or elevated blood sugar levels might be present, but does not necessarily occur. A study showed that 12 weeks of regular WB-EMS in combination with nutritional counseling had beneficial effect on the body weight and composition, and positively affected blood cholesterol levels, but not HbA1c levels [19].

A recent randomized controlled study investigated the effect of WB-EMS in comparison to resistance training on cardiometabolic risk factors. The study found that there was no significant effect on variables including HbA1c [20].

Our intention is to use WB-EMS focusing on HbA1c in people with prediabetes and then

examine its preliminary efficacy. Therefore, an appropriate design is needed to compare the HbA1c of the intervention group with that of the comparison groups.

Study objectives

Our main objective is to pilot feasibility for a randomized trial testing WB-EMS for HbA1c level in people with prediabetes. The primary objective is to provide an estimation of the preliminary efficacy of the intervention on glycemic control in a sedentary group of adults with prediabetes compared with control groups.

The secondary objectives are to assess the preliminary efficacy of

- 1) WB-EMS training on individual's biomarkers (e.g. triglycerides, high-density lipoprotein (HDL), low-density lipoprotein (LDL)),
- 2) WB-EMS training on body composition, blood pressure, and heart rate.
- 3) WB-EMS on other outcomes, such as quality of life and stress

Methods

Study design and setting

The trial is designed as a randomized controlled pilot trial in a parallel design with three arms. This trial protocol version 1 from August 2023 was approved by the local Ethics Committee at the Medical Faculty of the Eberhard Karls University and at the University Hospital of Tübingen (reference: 525/2023B02) and registered on the Clinicaltrials.gov with the ID NCT06188481. The trial will be conducted at the Department of Population-Based Medicine at the University Hospital of Tübingen, Germany.

Study population

The study population will be selected on the basis of the following inclusion and exclusion criteria.

Inclusion criteria:

- (1) community-dwelling sedentary men and women aged 40-65 years without T2DM,
- (2) elevated HbA1c levels (5.7%-6.4%),
- (3) not functionally impaired (Short Physical Performance Battery (SPPB) ≥ 10),
- (4) signed informed consent,
- (5) consent to use the WB-EMS and activity tracker.

Exclusion criteria:

- (1) high-grade arrhythmia, cardiac pacemaker carriers, heart failure, nephropathy,
- (2) cognitive impairment,
- (3) T2DM.

Study procedure

Figure 1 illustrates the study procedure. Screening of participants will take place in two stages (telephone pre-screening, on-site screening). During the telephone pre-screening, the study staff will check key study eligibility criteria and inform participants about the study activities. If they are interested, they will be invited to the on-site screening. During the on-site screening, the study staff will perform the HbA1c measurement. Furthermore, physical examination (e.g. blood pressure) and detailed medical history will be collected to account for any possible contraindications and to ensure safe participation in the training program.

All participants will receive a written information letter and informed consent form for participation. All participants will be informed of the voluntary nature of the trial and that they may withdraw from the trial at any time without giving a reason. Only those who confirm their willingness to participate in the trial by signing the informed consent form will be enrolled. The final determination of eligibility to participate in the trial will take place after the screening. The participants will be randomized to one intervention group (IG) and two control groups (CGs). The duration of the intervention will be 16 weeks. At the end of the intervention period, subjects will return to the study site for a follow-up assessment. The participants of the IG will

have a second follow-up assessment after 32 weeks.

Sample size

The study population will consist of 60 individuals with 20 subjects in each study group. As this is a pilot study, no formal power analysis has been carried out. However, the target sample size was based on recruitment for 15 months at an estimated rate of four patients per month (ie, 60 patients), which was considered adequate to obtain reasonably reliable sample size estimates [21–23].

Recruitment

Participants will be recruited through newspaper and social media advertisements. Flyers and posters will also be distributed to local doctors' offices, pharmacies, hospitals, community centers and self-help groups.

Randomization, blinding, and concealment of allocation

Eligible participants will be allocated to the study groups by computerized block randomization. This will be performed by an independent trial statistician. Neither the participants nor the researchers will know their allocation in advance. The participants will be strictly separated after randomization. The study staff will prevent participants from different study groups from arriving at the study site at the same time by planning study visits accordingly.

The blinding strategy focuses on study assistants and outcome assessors who will be blinded to the participants' group status. All data analysts will also be blinded to group allocation.

Intervention

The main intervention will be WB-EMS using equipment that has EU medical device approval (miha bodytec®, Type II, Gersthofen, Germany). This equipment allows the simultaneous stimulation of up to 12 muscle groups (e.g. thigh and upper arm, gluteal region, abdomen, chest, lower back, upper back) with a total stimulation area of approximately 2800 cm². The participants will receive the functional training underwear recommended by the manufacturer (water-absorbent cotton/elastane mix), over which a vest (upper body) and belts (upper arms, thigh and gluteus muscle) are individually adjusted before each session. The system allows the intensity to be adjusted for each region (see Figure 2). We will use an impulse protocol and exercise setting that has been evaluated in recent studies focusing on total and regional muscle mass [13], body fat [24] and physical function [13] in older cohorts. Furthermore, we designed low-intensity movements/exercises to minimize the effect of voluntary movements themselves as low as possible.

A bipolar electric current with a frequency of 85 Hz and an impulse width of 350 µs is used in an interval approach with 6 s of EMS stimulation with a direct impulse boost and 4 s of rest. During the 6-s stimulation period, low-effort movements are performed in a standing position. The intensity of the EMS will be regulated on the basis of the individual's reported Rating of Perceived Exertion (RPE) and the Borg Category Ratio Scale (Borg-CR10) [15]. We will use RPE to generate and maintain a sufficient but tolerable intensity of EMS application. After 4 weeks of conditioning at a lower pulse intensity, participants will be encouraged to increase the intensity. The impulse intensity will be adjusted individually for each body region in close interaction with the participant. During the session, the instructors will increase the intensity slightly every 3 minutes in close cooperation with the participants in order to maintain the target RPE (6-7) during the session. We will use a personal training setting with one certified and experienced instructor responsible for a maximum of two participants. The training will adhere to the requirements of the international EMS guidelines [25].

Study groups

Intervention group

The participants in the IG will train following a supervised and guided WB-EMS program 1.5 times per week (e.g., every Monday or Tuesday and every other Thursday or Friday) for 16 weeks. The duration of each training session will be 20 minutes. Each session will include a 3-minute warm-up, a 15-minute main workout, and a 2-minute cool-down. Additionally, participants will be asked to wear an activity tracker on their wrist for the entire 16-week study period and to participate in the lifestyle education program.

Control groups

Two CGs are planned for the study. One CG will receive both the activity tracker to measure daily steps and the evidence-based lifestyle education program. The other CG will receive the lifestyle education program only.

Activity tracker

The activity tracker (vivosmart 5, Garmin) will be used for self-monitoring of daily steps. Activity trackers may improve physical activity independent of the intervention (e.g., people can monitor their daily steps). [26]. Therefore, it is important to include two control groups: one with the activity tracker to isolate the net effect of the WB-EMS intervention and one without the activity tracker.

Lifestyle education program

The evidence-based lifestyle education program consists of six 20-minute sessions that focus on lifestyle factors and diabetes prevention. The aim of the program is to provide education, information and advice to prevent disease progression and improve quality of life and mobility. The education program includes topics such as basics of healthy nutrition pattern, meal planning, importance of physical activity and stress management. Participants will receive recorded videos on each topic to ensure standardization of the content. To assess the compliance, the participants will be followed up by the study staff, who will ask, whether the program was completed, or remind to complete the program, in case of non-compliance.

Outcome measures

The assessments described below will be performed at baseline (day 1) and after the 16-week intervention period. IG participants will also be assessed at 32 weeks. The timetable of the assessments is presented in Table 1. All measurements will be performed by the same research assistant, using identically calibrated equipment, in the exact same environment and at approximately the same time of day (± 90 minutes).

Table 1: Timetable of study procedure and assessments

	STUDY PERIOD				
	Enrollment	Allocation	Intervention	Follow-ups	
Timepoint		Day 0	Week 1 to 16	Week 16	Week 32
Eligibility screening	X				
Informed consent	X				
Review inclusion and exclusion criteria	X				
Allocation		X			

Assessments					
HbA1c assessment	X			X	X
Lipid and cholesterol profile		X		X	X
Blood pressure		X		X	X
Hand grip strength		X		X	X
Body composition		X		X	X
Waist circumference		X		X	X
Questionnaires (WHOQOL-BREF, SFI, PSS-10, PHQ-9)		X		X	X
Interventions					
WB-EMS, activity tracker and evidence-based lifestyle education program			WB-EMS 1.5/week		
			activity tracker		
			6 x 20 min LEP		
Activity tracker and evidence-based lifestyle education program			6 x 20 min LEP		
			activity tracker		
Evidence-based lifestyle education program			6 x 20 min LEP		

Abbreviations: WB-EMS: whole-body electromyostimulation, IG: intervention group, PSS: Perceived Stress Scale, WHOQOL-BREF: WHO Health-Related Quality of Life, SFI: Secure Flourish Index, PHQ: Patient Health Questionnaire, LEP: lifestyle education program

Primary outcome

As a pilot study, our main outcome is to assess the feasibility of the proposed intervention, with specific attention to the recruitment, adherence, and completion rates of the follow-up assessments. The prespecified primary outcome measure will be HbA1c levels. A blood sample (capillary blood) will be collected via the finger stick technique. There is no need for the participants to fast prior to blood measurement. The sample will be analyzed according to the manufacturer's instructions using cobas b 101 system 2.0 (Roche Diagnostics, Mannheim, Germany).

Secondary outcomes

Cardiometabolic parameters: Changes from baseline in the levels of triglycerides, total cholesterol as well as HDL and LDL cholesterol will be assessed in capillary blood. A blood sample will be collected with finger stick technique. The sample will be analyzed according to manufacturer's instructions using cobas b 101 system 2.0 (Roche Diagnostics, Mannheim, Germany).

Waist circumference will be measured with a standard tape measure around the abdomen between the distal end of the rib cage and the top of the iliac crest along the horizontal plane (WHO guidelines).

Additionally, changes in systolic and diastolic blood pressure as well as in heart rate will be

investigated (boso medicux X, BOSCH + SOHN, Jungingen, Germany). The measurements will be performed in the sitting position after a short resting period of at least 5 minutes.

Body composition: Body mass and composition will be measured by direct segmental, multi-frequency bio-impedance technique (InBody 770, Seoul, South Korea). This device measures impedance of the trunk, arms and legs separately using a tetrapolar eight-point tactile electrode system that applies six frequencies (1, 5, 50, 250, 500 and 1000 kHz).

Muscle strength: Hand grip strength will be measured using a hydraulic hand dynamometer SAEHAN SH5001 (SAEHAN Corporation, Tisselt, Belgium). The dynamometer grip width will be adjusted individually to the participant's hand size. Tests will be performed in an upright standing position, arms down by the side. The standardized instruction to the participants will be consistently "squeeze as strongly as possible". Three tests intermitted by 20 s of rest will be performed for the dominant hand, the average value will be used for the evaluation.

Stress: A self-reported questionnaire Perceived Stress Scale consisting of 10 questions about stress (PSS-10) will be used [27]. In each question, subjects are asked how often they feel a certain way on a 5-point scale from 1 for 'never' to 5 for 'very often'. The PSS score indicates levels of perceived stress, whereby higher scores indicate higher stress levels.

Depressive symptoms: The Patient Health Questionnaire which consists of 9 questions on depressive symptoms (PHQ-9) is a standardized and validated self-reported questionnaire. It will be used to assess for changes from baseline regarding the presence and severity of such symptoms. Possible score ranges from 0 (no depression) to 27 (severe depression) [28].

Health-related quality of life: The standardized and validated questionnaire WHO Health-Related Quality of Life (WHOQOL-BREF) consisting of 26 questions on general quality of life will be used [29]. This self-reported tool includes four domains: physical health, psychological health, social relationships, and environmental health. It also contains items about quality of life and general health. Each item of the WHOQOL-BREF is scored from 1 to 5 on a 5-point Likert scale. The scores are then transformed linearly to a 0 - 100 scale. A higher score indicates a higher quality of life.

Well-being: Changes from baseline in self-reported well-being will be evaluated with the standardized and validated questionnaire Secure Flourish Index (SFI), which consists of 12 questions on general well-being. Each of the questions is assessed on a scale of 0 – 10. The SFI score is obtained by summing the scores from the 12 questions and results in a score from 0 – 120 [30].

Baseline characteristics

Table 1 describes the type of data and variables that are collected at various stages of the project. Furthermore, the following demographic and descriptive data are collected at the baseline assessment: gender, age, height (cm), medical history (diseases, surgery, medicine consumption), physical activity (type, frequency, duration), educational level and employment status.

Trainers will monitor adverse events by asking participants if they have experienced any health problems (e.g., physical injury, muscle soreness) since the last session. The research team will monitor any adverse events that may occur during participation in this study.

Data management and analysis

Data collection and storage

The personal data of the subjects will be treated confidentially and will only be accessible to the persons directly involved in the study. For the scientific analysis, the data will be stored electronically in a pseudonymized form by assigning a subject ID to each study participant, which does not allow any conclusions to be drawn about the individual. The list of subject codes and participants will be kept in a locked cabinet and will be accessible only to authorized personnel. Further processing, analysis and publication of the data for scientific purposes will

be done anonymously, so that no conclusions can be drawn about individual study participants.

Activity tracker data

The tracker can be set to measure a customizable set of matrices, as needed for the project to avoid unnecessary data collection. This pilot study will collect and process step counts. The Fitrockr Health Data Research & Analytics Platform provides a platform for research and clinical trials to collect and analyze data from wearable devices. Authorized members of the research team will access the device data via Bluetooth or a USB cable using the Fitrockr Hub App when the study participants are on-site. All data will be securely stored on a server in Germany. The participants' usernames will be generated according to the subject ID.

Statistical analysis

The data will be analyzed using the intention-to-treat (ITT) principle which includes all randomized participants. Descriptive statistics, such as mean and standard deviation (SD) will be performed for continuous variables, and frequencies and percentages will be used for categorical variables. Diagnostic tests will be conducted for normality of residuals, linearity, multicollinearity, functional relation, influence and/or outliers in the dataset. Assuming data normality, two-way repeated measures ANOVA and 95% CI will be used to assess outcomes at baseline and at 4-month follow-up. The analysis plan will not include reporting of p-values, in accordance with the CONSORT 2010 statement: extension to randomized pilot and feasibility trials that "any estimates of effect using participant outcomes as they are likely to be measured in the future definitive RCT would be reported as estimates with 95% confidence intervals without P values—because pilot trials are not powered for testing hypotheses about effectiveness." [31]

Analysis of covariance will be used to calculate the effect estimate (study outcomes) of the intervention. Effect size calculation for between-group intervention effects and within-group effects will also be performed. All statistical analyses will be performed using SPSS Statistics (version 28, IBM).

Results

Following the approval of the ethics committee on November 6, 2023, preparations for the advertisement and implementation of the project, including the registration of the clinical trial on ClinicalTrials.gov (ID: NCT06188481), were conducted in early January 2024.

The recruitment of participants and on-site screening started at the beginning of February 2024. At the time of submission of this protocol for publication, the recruitment was still ongoing. As of October 2024, 169 individuals have registered for the study. Of these, 76 individuals were invited to attend the on-site screening and the rest was excluded during the telephone pre-screening process. Finally, 47 participants were eligible, and 42 from those were randomized and allocated to the study groups. The 16-week intervention phase starts in groups of at least six participants (block randomization) in order to ensure randomized allocation to the three study groups. The use of sealed opaque envelopes ensured that neither participants nor researchers were aware of the allocation prior to the study (i.e., allocation concealment).

The study is still in progress, and thus far, no adverse events have been recorded. The anticipated date of recruitment completion is April 2025.

Discussion

While exercise is regarded as one of the key components in T2DM management and prevention [32], the adherence to regular exercise still remains low in people with T2DM and prediabetes [9]. Barriers to exercise adherence include the lack of time, lack of motivation/enthusiasm, physical discomfort, and social and environmental barriers [10]. Due to the ease of implementation, time efficiency and personalized application of WB-EMS, it is

increasingly the subject of scientific research.

This randomized clinical trial is designed to evaluate the preliminary efficacy of 16 weeks of WB-EMS training on changes in HbA1c levels. The results of the current study will not only give new insights into the effect on WB-EMS in prediabetes, but also serve as a foundation for future studies in this population. This may make a big contribution to the development of preventive strategies specifically for type 2 diabetes.

A meta-analysis of 35 studies reported that local EMS intervention lowered fasting blood glucose (FBS) (SMD: 0.48; 95% CI: 0.17 to 0.78; $p=0.002$; $I^2=0\%$) [17]. In a recent 16-week non-RCT in the Netherlands, WB-EMS was used to reduce visceral fat in elderly patients with non-insulin-dependent diabetes mellitus (NIDDM). The exercise program consisted of 20 minutes of WB-EMS twice a week, with HbA1c measured at baseline and at the end of the intervention. However, the control group consisted of healthy participants without diabetes, which led to a non-HbA1c comparison between groups. Only male patients showed a significant change in HbA1c (55.7 ± 12.7 mmol/mol vs. 52.7 ± 13.8 mmol/mol, $p<0.05$), while female patients did not experience a significant change [33]. To the best of our knowledge, there is currently no other randomized controlled trial evaluating the effect of WB-EMS on glycemic control in people with prediabetes.

HbA1c is considered the gold standard for long-term blood glucose monitoring as it reflects average levels over a period of 2-3 months. Studies have shown that exercise programs typically take longer than 12 weeks to impact HbA1c levels, which is similar to the lifespan of a red blood cell. Most interventional exercise studies that demonstrate improved HbA1c levels have a duration of approximately 13 weeks [34]. Additionally, previous studies using WB-EMS have shown significant improvement in muscle mass after a 16-week intervention with similar training frequency and impulse protocol as those used in our study [13,24]. Accordingly, the WB-EMS intervention will be conducted over a period of 16-weeks in this study.

To monitor the level of PA, participants in the IG will be asked to wear an activity tracker during the intervention. However, there is evidence of positive effect of activity trackers on PA behavior through personalized feedback on daily steps [26]. Therefore, some of the participants will be randomized to an additional control group with an activity tracker in order to isolate the net effect of the WB-EMS intervention.

The results of the current study may provide evidence of the preliminary efficacy of WB-EMS for a future definitive RCT as an alternative adoptive exercise approach to motivate sedentary people with prediabetes who are unable or choose not to follow a traditional exercise program. As this is the first RCT with WB-EMS training to focus on individuals with prediabetes and to address HbA1c as the primary outcome, the results of the trial will inform the sample size calculation for a definitive trial. Moreover, we believe that this research study will provide additional evidence for the health and performance aspects of the WB-EMS application.

Competing interests

The authors declare that they have no competing interests.

List of abbreviations

BIA: bioimpedance analysis; Borg-CR10: Borg Category Ratio Scale; CG: control group; IG: intervention group; ITT: intention-to-treat; FBS: fasting blood sugar; HbA1c: hemoglobin A1c; HDL: high-density lipoprotein; IDF: International Diabetes Federation; LEP: lifestyle education program; LDL: low-density lipoprotein; NIDDM: non-insulin-dependent diabetes mellitus; T2DM: type 2 diabetes mellitus; SPPB: Short Physical Performance Battery; PA: physical activity; PHQ-9: Patient Health Questionnaire; RCT: randomized controlled trial; RPE: Rating of Perceived Exertion; SD: standard deviation; SFI: Secure Flourish Index; WB-EMS: whole-body electromyostimulation; WHOQOL-BREF: WHO Health-Related Quality of Life

References

1. International Diabetes Federation. IDF Diabetes Atlas. 9th edition. Brussels, Belgium: The International Diabetes Federation; 2019. Available from: www.diabetesatlas.org ISBN:978-2-930229-87-4
4. Herman WH. Prediabetes diagnosis and management. *JAMA* 2023;329(14):1157. doi: 10.1001/jama.2023.4406
5. Ligthart S, Van Herpt TTW, Leening MJG, Kavousi M, Hofman A, Stricker BHC, Van Hoek M, Sijbrands EJG, Franco OH, Dehghan A. Lifetime risk of developing impaired glucose metabolism and eventual progression from prediabetes to type 2 diabetes: a prospective cohort study. *Lancet Diabetes Endocrinol* 2016;4(1):44–51. doi: 10.1016/S2213-8587(15)00362-9
6. Stöckl D, Rückert-Eheberg I-M, Heier M, Peters A, Schipf S, Krabbe C, Völzke H, Tamayo T, Rathmann W, Meisinger C. Regional variability of lifestyle factors and hypertension with prediabetes and newly diagnosed type 2 diabetes mellitus: the population-based KORA-F4 and SHIP-TREND studies in Germany. *PLoS ONE* 2016 Jun 8;11(6):e0156736. doi: 10.1371/journal.pone.0156736
7. Campbell TJ, Alberga A, Rosella LC. The impact of access to health services on prediabetes awareness: a population-based study. *Prev Med* 2016;93:7–13. doi: 10.1016/j.ypmed.2016.09.008
8. Rossen J, Von Rosen P, Johansson U-B, Brismar K, Hagströmer M. Associations of physical activity and sedentary behavior with cardiometabolic biomarkers in prediabetes and type 2 diabetes: a compositional data analysis. *Phys Sportsmed* 2020;48(2):222–228. doi: 10.1080/00913847.2019.1684811
9. Mortensen SR, Skou ST, Brønd JC, Ried-Larsen M, Petersen TL, Jørgensen LB, Jepsen R, Tang LH, Bruun-Rasmussen NE, Grøntved A. Detailed descriptions of physical activity patterns among individuals with diabetes and prediabetes: the Lolland-Falster Health Study. *BMJ Open Diab Res Care* 2023;11(5):e003493. doi: 10.1136/bmjdr-2023-003493
10. Thielen SC, Reusch JEB, Regensteiner JG. A narrative review of exercise participation among adults with prediabetes or type 2 diabetes: barriers and solutions. *Front Clin Diabetes Healthc* 2023;4:1218692. doi: 10.3389/fcdhc.2023.1218692
11. Kemmler W, Von Stengel S, Kohl M, Rohleder N, Bertsch T, Sieber CC, Freiburger E, Kob R. Safety of a combined WB-EMS and high-protein diet intervention in sarcopenic obese elderly men. *Clin Interv Aging* 2020;15:953–967. doi: 10.2147/CIA.S248868
12. Kemmler W, Kleinöder H, Fröhlich M. Editorial: Whole-body electromyostimulation: a training technology to improve health and performance in humans? *Front Physiol* 2020;11:523. doi: 10.3389/fphys.2020.00523
13. Kemmler W, Grimm A, Bebenek M, Kohl M, Von Stengel S. Effects of combined whole-body

- electromyostimulation and protein supplementation on local and overall muscle/fat distribution in older men with sarcopenic obesity: the randomized controlled Franconia Sarcopenic Obesity (FranSO) study. *Calcif Tissue Int* 2018;103(3):266–277. doi: 10.1007/s00223-018-0424-2
14. Von Stengel S, Bebenek M, Engelke K, Kemmler W. Whole-body electromyostimulation to fight osteopenia in elderly females: the randomized controlled Training and Electrostimulation Trial (TEST-III). *J Osteoporos* 2015;2015:1–7. doi: 10.1155/2015/643520
 15. Kemmler W, Weissenfels A, Teschler M, Willert S, Bebenek M, Shojaa M, Kohl M, Freiburger E, Sieber C, von Stengel S. Whole-body electromyostimulation and protein supplementation favorably affect sarcopenic obesity in community-dwelling older men at risk: the randomized controlled FranSO study. *Clin Interv Aging* 2017;12:1503–1513. PMID:28989278
 16. Van Buuren F, Horstkotte D, Mellwig KP, Fründ A, Vlachojannis M, Bogunovic N, Dimitriadis Z, Vortherms J, Humphrey R, Niebauer J. Electrical Myostimulation (EMS) improves glucose metabolism and oxygen uptake in type 2 diabetes mellitus patients-results from the EMS Study. *Diabetes Technol Ther* 2015;17(6):413–419. PMID:25734937
 17. Sanchez MJ, Mossayebi A, Sigaroodi S, Apaflor JN, Galvan MJ, Min K, Agullo FJ, Wagler A, Bajpeyi S. Effects of neuromuscular electrical stimulation on glycemic control: a systematic review and meta-analysis. *Front Endocrinol* 2023;14:1222532. doi: 10.3389/fendo.2023.1222532
 18. Holzer R, Schulte-Körne B, Seidler J, Predel H-G, Brinkmann C. Effects of acute resistance exercise with and without whole-body electromyostimulation and endurance exercise on the postprandial glucose regulation in patients with type 2 diabetes mellitus: a randomized crossover study. *Nutrients* 2021;13(12):4322. PMID:34959874
 19. Reljic D, Konturek P, Herrmann H, Neurath M, Zopf Y. Effects of whole-body electromyostimulation exercise and caloric restriction on cardiometabolic risk profile and muscle strength in obese women with the metabolic syndrome: a pilot study. *J Physiol Pharmacol* 2020;71(1):89–98. doi: 10.26402/jpp.2020.1.08
 20. Reljic D, Herrmann HJ, Neurath MF, Zopf Y. Iron Beats Electricity: Resistance Training but Not Whole-Body Electromyostimulation Improves Cardiometabolic Health in Obese Metabolic Syndrome Patients during Caloric Restriction—A Randomized-Controlled Study. *Nutrients* 2021;13(5):1640. doi: 10.3390/nu13051640
 21. Kunselman AR. A brief overview of pilot studies and their sample size justification. *Fertility and Sterility* 2024;121(6):899–901. doi: 10.1016/j.fertnstert.2024.01.040
 22. Julious SA. Sample size of 12 per group rule of thumb for a pilot study. *Pharmaceutical Statistics* 2005;4(4):287–291. doi: 10.1002/pst.185
 23. Van Belle G. Statistical Rules of Thumb, 2nd edition. Hoboken, NJ, USA: John Wiley & Sons, Inc.; 2008. ISBN:978-0-470-37796-3


24. Kemmler W, Teschler M, Weißenfels A, Bebenek M, Fröhlich M, Kohl M, Von Stengel S. Effects of whole-body electromyostimulation versus high-intensity resistance exercise on body composition and strength: a randomized controlled study. *Evid Based Complement Alternat Med* 2016;2016:1–9. doi: 10.1155/2016/9236809
25. Kemmler W, Fröhlich M, Ludwig O, Eifler C, von Stengel S, Willert S, Teschler M, Weissenfels A, Kleinöder H, Micke F, Wirtz N, Zinner C, Filipovic A, Wegener B, Berger J, Evangelista A, D'Ottavio S, Sara JDS, Lerman A, Perez de Arrilucea Le Floch UA, Carle-Calo A, Guitierrez A, Amaro-Gahete FJ. Position statement and updated international guideline for safe and effective whole-body electromyostimulation training-the need for common sense in WB-EMS application. *Front Physiol* 2023;14:1174103. PMID:37035684
26. Ellingson LD, Lansing JE, DeShaw KJ, Peyer KL, Bai Y, Perez M, Phillips LA, Welk GJ. Evaluating motivational interviewing and habit formation to enhance the effect of activity trackers on healthy adults' activity levels: randomized intervention. *JMIR Mhealth Uhealth* 2019;7(2):e10988. doi: 10.2196/10988
27. Cohen S, Williamson GM. Perceived stress in a probability sample of the United States. *The Social Psychology of Health* 1988;31–67.
28. Kroenke K, Spitzer RL, Williams JBW. The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med* 2001;16(9):606–613. doi: 10.1046/j.1525-1497.2001.016009606.x
29. Kim S. World Health Organization Quality of Life (WHOQOL) Assessment. In: Michalos AC, editor. *Encyclopedia of Quality of Life and Well-Being Research*. Dordrecht: Springer Netherlands; 2014. p. 7578–7880. ISBN:978-94-007-0752-8
30. VanderWeele TJ. On the promotion of human flourishing. *Proc Natl Acad Sci USA* 2017;114(31):8148–8156. doi: 10.1073/pnas.1702996114
31. Eldridge SM, Chan CL, Campbell MJ, Bond CM, Hopewell S, Thabane L, Lancaster GA, PAFS consensus group. CONSORT 2010 statement: extension to randomised pilot and feasibility trials. *BMJ* 2016;355:i5239. PMID:27777223
32. Kanaley JA, Colberg SR, Corcoran MH, Malin SK, Rodriguez NR, Crespo CJ, Kirwan JP, Zierath JR. Exercise/Physical activity in individuals with type 2 diabetes: a consensus statement from the american college of sports medicine. *Med Scin Sports & Exerc* 2022;54(2):353–368. doi: 10.1249/MSS.0000000000002800
33. Houdijk APJ, Bos NFJME, Verduin WM, Hijdendaal MM, Zwartkruis MAL. Visceral fat loss by whole-body electromyostimulation is attenuated in male and absent in female older non-insulin-dependent diabetes patients. *Endocrino Diabet Metab* 2022;5(6):e377. doi: 10.1002/edm2.377
34. Grace A, Chan E, Giallauria F, Graham PL, Smart NA. Clinical outcomes and glycaemic responses to different aerobic exercise training intensities in type II diabetes: a systematic review and meta-analysis. *Cardiovasc Diabetol* 2017;16(1):37. doi: 10.1186/s12933-017-0518-6



Supplementary Files

Figures

Flow chart of the study procedure. Abbreviations: WB-EMS: whole body electromyostimulation, T2DM: type2 diabetes.



Whole-body electromyostimulation (WB-EMS).



CONSORT (or other) checklists

SPIRIT checklist.

URL: <http://asset.jmir.pub/assets/334cd60f4eb37322574103cdba428447.pdf>

Model consent form (addition to the SPIRIT checklist).

URL: <http://asset.jmir.pub/assets/04da04086d1d7078971485750fa9959f.pdf>

