

Empowering Patients and Caregivers to Use AI and Computer Vision for Wound Monitoring: A Feasibility Study

Rose Raizmam, José Luis Ramírez-GarcíaLuna, Tanmoy Newaz, Sheila C Wang, Gregory K Berry, Ling Yaun Kong, Heba Tallah Mohammed, Robert Douglas John Fraser

Submitted to: Journal of Participatory Medicine
on: December 01, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript.....	5
Supplementary Files.....	21
0.....	21
Figures	22
Figure 1.....	23
Figure 2.....	24
Figure 3.....	25
Multimedia Appendixes	26
Multimedia Appendix 0.....	27
CONSORT (or other) checklists.....	28
CONSORT (or other) checklist 0.....	28

Empowering Patients and Caregivers to Use AI and Computer Vision for Wound Monitoring: A Feasibility Study

Rose Raizmam^{1, 2} NP, MN; José Luis Ramírez-GarcíaLuna³ MD, PhD; Tanmoy Newaz⁴ BSC; Sheila C Wang^{5, 6, 7} MD, PhD; Gregory K Berry³ MD; Ling Yaun Kong^{8, 9} MD; Heba Tallah Mohammed⁵ MD, PhD; Robert Douglas John Fraser^{5, 10} BSCN, MN

¹ Scarborough Health Network Scarborough CA

² University of Toronto Lawrence S Bloomberg Faculty of Nursing Toronto CA

³ McGill University Health Centre Department of Surgery Montreal CA

⁴ University of Calgary Calgary CA

⁵ Swift Medical Inc. Toronto CA

⁶ University of Toronto Faculty of Medicine Toronto CA

⁷ Women's College Hospital Toronto CA

⁸ McGill University Montreal CA

⁹ Jewish General Hospital Montreal CA

¹⁰ Western University Arthur Labatt Family School of Nursing London CA

Corresponding Author:

Robert Douglas John Fraser BSCN, MN

Swift Medical Inc.

1 King S W

Suite 4800

Toronto

CA

Abstract

Background: Chronic wounds affect 1-2% of the global population, and pose significant health and quality-of-life challenges for patients and caregivers. Advances in artificial intelligence (AI) and computer vision (CV) technologies present new opportunities for enhancing wound care, particularly through remote monitoring and patient engagement. A Digital Wound Care Solution (DWCS) that objective wound tracking using AI/CV was redesigned as a patient-facing mobile application to empower patients and caregivers to actively participate in wound monitoring and management.

Objective: This study aimed to evaluate the feasibility, usability, and preliminary clinical outcomes of the Patient Connect application in enabling patients and caregivers to remotely capture and share wound data with healthcare providers.

Methods: A feasibility study was conducted at two outpatient clinics in Canada between May 2020 and February 2021. Twenty-eight patients with chronic wounds were recruited and trained to use the Patient Connect app for wound imaging and secure data sharing with their care teams. Wound images and data were analyzed using AI/CV models integrated into the app. Clinicians reviewed the data to inform treatment decisions during follow-up visits or remotely. Key metrics included app usage frequency, patient engagement, and wound closure rates.

Results: Participants captured a median of 13 wound images per wound, with images submitted every 8 days on average. The study cohort included patients with diabetic ulcers, venous ulcers, pressure injuries, and post-surgical wounds. A median wound closure rate of 80% was achieved across all patients, demonstrating the app's clinical potential. Feedback from patients and clinicians highlighted the app's usability, data security features, and ability to enhance remote monitoring.

Conclusions: The Patient Connect application effectively engaged patients and caregivers in chronic wound care, demonstrating feasibility and promising clinical outcomes. By enabling secure, remote wound monitoring through AI/CV technology, the app has the potential to improve patient adherence, enhance care accessibility, and optimize clinical workflows. Future studies should focus on evaluating its scalability, cost-effectiveness, and broader applicability in diverse healthcare settings.

(JMIR Preprints 01/12/2024:69470)

DOI: <https://doi.org/10.2196/preprints.69470>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in a JMIR publication, my title and abstract will remain visible to all users.

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Empowering Patients and Caregivers to Use AI and Computer Vision for Wound Monitoring: A Feasibility Study

Background:

Chronic wounds affect 1-2% of the global population, and pose significant health and quality-of-life challenges for patients and caregivers. Advances in artificial intelligence (AI) and computer vision (CV) technologies present new opportunities for enhancing wound care, particularly through remote monitoring and patient engagement. A Digital Wound Care Solution (DWCS) that facilitates wound tracking using AI/CV was redesigned as a patient-facing mobile application to empower patients and caregivers to actively participate in wound monitoring and management.

Objective:

This study aimed to evaluate the feasibility, usability, and preliminary clinical outcomes of the Patient Connect application in enabling patients and caregivers to remotely capture and share wound data with healthcare providers.

Methods:

A feasibility study was conducted at two outpatient clinics in Canada between May 2020 and February 2021. Twenty-eight patients with chronic wounds were recruited and trained to use the Patient Connect app for wound imaging and secure data sharing with their care teams. Wound images and data were analyzed using AI/CV models integrated into the app. Clinicians reviewed the data to inform treatment decisions during follow-up visits or remotely. Key metrics included app usage frequency, patient engagement, and wound closure rates.

Results:

Participants captured a median of 13 wound images per wound, with images submitted every 8 days on average. The study cohort included patients with diabetic ulcers, venous ulcers, pressure injuries, and post-surgical wounds. A median wound closure rate of 80% was achieved across all patients, demonstrating the app's clinical potential. Feedback from patients and clinicians highlighted during the feasibility testing support insight into the app's usability, data security features, and ability to enhance remote monitoring that need to be explored in further qualitative research.

Conclusion:

The Patient Connect application effectively engaged patients and caregivers in chronic wound care, demonstrating feasibility and promising clinical outcomes. By enabling secure, remote wound monitoring through AI/CV technology, the app has the potential to improve patient adherence, enhance care accessibility, and optimize clinical workflows. Future studies should focus on evaluating its scalability, cost-effectiveness, and broader applicability in diverse healthcare settings.

Introduction

Chronic wounds are commonly defined as wounds that fail to heal within 4- 12 weeks through normal, timely and orderly stages [1]. These wounds pose a major public health challenge, with 1-2% of the global population estimated to experience a chronic wound during their lifetimes [2]. Diabetic ulcers (DU), venous ulcers (VU), and pressure injuries (PI) are especially prevalent, making up over 90% of all chronic wounds [3] and often requiring significant wound care management and resources. However, due to their low rate of complete healing, chronic wounds have major impacts on both the health and quality of life of patients and their families, leading to significant issues such as severe and prolonged pain, loss of function and mobility, amputation, mental health deterioration, social isolation and embarrassment, financial burden, and chronic morbidity or death [4]. Recently, there has been a significant transformation in healthcare delivery, focusing on remote access through telemedicine that leverages the widespread availability of smartphones and their applications. Technologies that facilitate telemedicine and ensure continuity of care for chronic wound patients are urgently needed, as high risk of wound-related complications exist for those without access to consistent follow-ups [5].

The rise of artificial intelligence (AI) has shown great promise, particularly in the field of wound care. These technologies provide healthcare professionals with novel tools that contribute towards many improvements in treatment efficiency and efficacy, including early detection, risk factor analysis, prediction, diagnosis, intelligent treatment, outcome prediction, and prognostic evaluation [6]. Additionally, AI-powered tools have been shown to empower patients to take control of their own health and wellbeing. For instance, AI tools can provide patients with information regarding their conditions and treatment options, thereby enabling them to make informed decisions while also strengthening patient-healthcare provider relationships through trust-building [7]. Computer vision (CV) is a particular form of AI that extracts information from digital images or videos in order to recognize content from visual data [8]. These technologies are especially promising in the field of wound care, as they can help classify wound severity, provide accurate predictions of wound healing, and track changes in wounds over time through image analysis [9-10]. CV technologies have previously been shown to provide significant time savings during wound assessments [11], decrease costs and days needed for wound healing [12], and improve data capture reproducibility and accuracy [13]. Notably, patients have also been found to exhibit positive perceptions towards the use of wound photography in their treatment journeys by helping them track their wound progress or increasing their involvement within their own care [14].

Swift Medical™ Skin & Wound (hereafter referred to as Digital Wound Care Solution (DWCS)) developed a mobile application and dashboard, specifically designed to accurately and reliably measure and document wound characteristics. The system, which is already available and is a privacy-compliant (HIPAA and PHIPA), Health Canada registered and FDA Class I medical device, uses CV technology to automatically focus and calculate wound dimensions from images acquired by the mobile device's camera, allowing users to obtain precise and consistent measurements. These capabilities have been demonstrated to reduce the time needed to assess the wounds of patients in a more accurate manner [11,15]. In addition to viewing a wound's image series over time, additional information such as healing-associated metrics, wound-bed information, anatomical location and patient identifiers are captured. While the app has provided doctors and wound-care specialists with a powerful assessment solution and a dashboard to remotely monitor and collaborate for an effective wound management strategy, in order to fully realize the system's potential, patients themselves will need to be able to acquire and securely share images and other relevant information with their care-providers. By actively engaging patients in their own wound care journeys through a patient-centric application, individuals may feel empowered to be more active in the treatment process.

Understanding the importance of innovative technologies in improving health outcomes for chronic

wound patients, the DWCS have recently developed a stream-lined, patient-facing version of the AI-powered application called Patient Connect. Patient Connect is designed for easy use by patients and/or their care-providers using their own personal smartphones, ensuring a more patient-centric approach to wound management (Figure 1). The user interface (UI) was designed with differences in technology and clinical literacy in mind. The DWCS has detailed clinical documentation fields an advanced reporting included. The Patient Connect interface had language changes to be grade three literacy level accessible. Educational materials including instructional videos and simply language guides for basic wound dressings were included within the app to attempt to improve engagement.

Patients are authorized directly by their healthcare provider and can only access their own records through their personal device. This requires a 2-step verification via email or a cell-phone number and their date of birth. Like the standard version of the app, it automatically focuses and calculates wound dimensions from the images acquired. Images and other measurements are not stored on the phone camera roll of the patient's personal devices, instead they are encrypted within the application and securely transmitted to healthcare providers on the same secure, web-based servers from the DWCS. The patient's healthcare provider can access the patient's generated images and patient-reported data using their application or the web dashboard; thereby, enabling the remote monitoring of wound progression.

The objective of this report is to present results of a feasibility study of early adopters of our patient-centric AI-powered wound assessment technology to image their wound to be included in their medical record and for self-monitoring, within two outpatient clinics in a university-affiliated hospital and a community hospital to determine overall feasibility, usability, and preliminary outcomes of the Patient Connect application.

Methods

A non-randomized, single arm-feasibility study was conducted between May 2020 and Feb 2021. The study received multisite ethics approval provided by the Scarborough Health Network Research Ethics Board (SUR-21-007). A nurse practitioner at Scarborough Health Network and two physicians at Montreal Jewish General Hospital were the primary clinicians engaged in the project, and both had previous experience using AI/CV-enabled wound care documentation in clinical practice. Standardized training was provided on enrolling patients, enabling access, and reviewing patient-submitted wound images and information in the clinician application and dashboard. Training materials were provided to support patient onboarding to use the service. This included multimedia content (videos on how to download and access the app) that was shared via SMS message when the patient was enrolled and content embedded within the app (e.g., how to capture wound images). Paper hand out material including instructions were also provided (see sample in supplemental files). Clinicians had access to review images submitted through the dashboard on a weekly basis and during follow-up visits.

A purposive sampling technique was employed to recruit patients or caregivers from the Montreal Jewish General Hospital and the Rouge Valley Scarborough Hospital for early testing of the Patient Connect app. A sample size between 20-30 participants was determined based on feasibility study design considerations. According to established feasibility study guidelines, sample sizes of 30 or fewer participants may be appropriate for qualitative feasibility studies [16]. This sample size allowed for evaluating the usability, engagement, and feasibility of the intervention while balancing recruitment and resource constraints. Patients were the primary focus for the inclusion criteria, with patient caregivers acting as an inclusion alternative if the patient consented. Inclusion criteria to the cohort were (1) patients' attending staff were already a user of the DWCS, (2) the patient or a close

relative possessed and was familiar with a smartphone device, and (3) the patient had a stable wound, as assessed by their healthcare provider. Caregivers were considered as an inclusion alternate if the patient consented. Caregivers were suitable alternatives if the wound was in an area that was difficult to image (e.g., sacrum, back) or the patient had limitations that made them unable to use the application (e.g., mobility, technology literacy). Exclusion criteria were Android phone users as the Patient Connect app currently only runs on iOS devices. In addition, the study excluded patients who did not consent and who did not approve their caregiver to act as an alternate, since, for these patients, caregiver participation was essential for independent app usage. No changes were made to the study methods after the commencement of the study, including eligibility criteria and assessment measurements. All prespecified metrics and inclusion criteria remained unchanged throughout the study period.

Enrolled participants were encouraged to use the application when their dressing was being changed by themselves, caregivers or other healthcare professionals outside of the participating organizations (e.g., home health). Two case series displaying the measurement and progress tracking of patient-captured and caregiver-captured wound images on the Patient Connect app are shown in Figures 2 and 3, respectively. Due to the variation in wound-changing protocols and the feasibility design, there was no set requirement for imaging completion by the patients per week. However, patients were encouraged to take at least one picture during each wound-changed session. The clinicians collected additional feedback during follow-up appointments. User experience, facilitators, and barriers were documented and shared with the project manager and software development team to support quality improvement and ensure app performance and stability.

Usability metrics were collected to assess feedback on the engagement, consistency, and effectiveness of the tool. These include the frequency app utilization (i.e., the number of wound images uploaded per patient), submission intervals, completion rates of imaging sessions, and tracking adherence rates concerning continued use during the study period. The mean was used to report on continuous or normally distributed variables, and median was used for data with outliers or skewed distribution (e.g., wound size, number of images) to minimize influence of extreme values (Table 1). The application has embedded monitoring software (Mixpanel, California, US) for debugging that enabled logging of successful logins, progress through the imaging workflow and de-identified summaries were available to the research team to see counts and frequency of image submission. These features are common practice in mobile and cloud based-software development to identify software issues and iteratively improve user workflows.

Additionally, qualitative feedback was collected about ease of use, technical difficulties, general user experience, satisfaction with the tool that was collected during follow-up visits, as well as barriers like light, clarity of images, and comfort level using the app alone. The degree of clinician engagement was assessed by tracking the frequency of image review, using the AI-assisted assessments into treatment decisions, and feedback on patient-submitted data.

The patients were followed until the closure of their wounds or February 2021, whichever occurred first. Wound closure was defined as a wound measurement of 0 cm². All data included in this report was obtained from the solution's deidentified servers, allowing for data retrieval while maintaining the confidentiality of patients' personal information.

Results

A total of 28 patients adopted the Patient Connect App as early users. The cohort included patients with varied wound types, including diabetic foot ulcer (DFU), venous leg ulcer (VLU), PI, and surgical wounds. The characteristics of the

wounds are presented on Table 1.

Approximately half of the patients were diabetics with plantar ulcers (52%). There was a balanced gender mix in this study, with 52% of patients (14) reporting as males and 48% (13) as females. The sample population had a range of wound sizes from 0.48 cm² to 27.91 cm² and a median size of 3.71cm². Wound measurement was captured from photographs using AI/CV models, so wounds outside of the photograph (i.e., circumferential) had limitations to their data. This suggested that single-surface wounds were optimal for patient and caregiver imaging and automated AI/CI analysis of the wound. Wound imaging was found to be ideally suited for patients with images on a single surface. However, it was possible to upload multiple images if wounds were circumferential.

The median follow-up was 3 months, with a median of 13 images acquired by the patient or caregiver per wound. Images were captured on average every 8 days. Interestingly, despite a general infrequency of in-person follow-up visits, the median wound closure rate recorded in the app was 80% (range 15-100%). No adverse events or unintended harms were reported among participants.

Projected Cost Savings:

The Patient Connect app enables remote monitoring of the wounds and reduces the need for in-person visits and related costs. With patients documenting a median of 13 images per wound over 3 months, this assessment could replace several visits to the clinic. Assuming that each time a picture is submitted, one trip is saved, that could mean there is the possibility of eliminating up to 13 trips per patient, representing savings anywhere between \$216 and \$432 in travel costs per patient (with an average travel cost of \$16.64 per visit)[17,18]. As for the sample of this study consisting of 28 patients, this would mean \$6048 to \$12096 in total travel savings over the three months. Savings could amount to \$216K-\$432K with 1,000 users in a year.

Additionally, fewer trips would equate to fewer hours lost at work for both patients and caregivers. Assuming 2 hours off work per visit at an average hourly wage of \$36.64 CAD , with 13 visits avoided, a direct saving of \$595/missed trip or \$16,674 could be achieved for the study cohort. A scale of 1000 users would mean savings of \$595K/year in workforce productivity.

User Experience and Quality Improvement Insights

Patient feedback on Patient Connect was useful in determining usability, engagement in wound care, and areas for improvement. Many participants noted that remote wound image capture and sharing opened their eyes to changes in the wound that made them more active in the wound care process and compliant with treatment. Some patients reported that taking pictures regularly helped monitor their healing and increase their motivation to adhere to wound care protocols such as the frequency of dressing changes, hygiene practices, and alleviating pressure techniques.

Although Patient Connect appeared useful in many aspects, several issues came to light. Literacy and accessibility problems were felt, particularly among older adults or other patients unfamiliar with smartphone apps, who sometimes required caregiver assistance to capture and submit images of their wounds. Patients had difficulty taking clear pictures if the wounds were in hard-to-reach areas (e.g., sacrum, back, heels) and tended to submit images erratically. Lighting posed challenges since some patients had difficulty ensuring adequate exposure for accurate AI analysis. While many people found the app helpful, some users experienced fatigue with engagement and became less consistent in taking images, especially if slow healing of the wound was involved. A few participants expressed common data privacy concerns about sharing images digitally, while continued education on encryption and security protocols was offered to help provide reassurance.

Discussion

In this report, we demonstrate that the Patient Connect's regular use by a group of selected patients allowed the remote monitoring of their wounds, successfully capturing medical-grade images that were subsequently used by clinicians for treatment decisions. This capability is not only crucial for maintaining continuity of care but also for enhancing patient engagement and treatment adherence, as evidenced by the increase in image sharing and self-monitoring behavior. The application facilitated the collection and analysis of data, which was instrumental in improving patient behavior and health outcomes by providing real-time feedback and enabling timely communication through wound status updates with healthcare professionals.

Patients using the Patient Connect application exhibited a high frequency of engagement with the AI software, submitting an average of 13 pictures, or one image every 8 days to clinicians throughout the duration of their wound care. Additionally, a median wound closure rate of 80% was observed across all patients and wound types. These findings suggest that the use of the Patient Connect application for participants may have supported engagement with monitoring wound healing, which may have influenced better healing outcomes across the diverse wound types. It is recognized, however, that factors such as standard wound care practices, clinical interventions, and individual patient conditions may have influenced the results. Clinical decisions within wound care may be delayed without adequate history. Patients in the study enabled a better record of the wound's response or lack of response to treatment that may support more timeline adjustments in care, which could be better understood through future research.

Interestingly, our results align with findings from other smartphone-based AI treatment platforms. For instance, Labovitz and colleagues demonstrated that, among patients with recently diagnosed ischemic strokes receiving anticoagulants, real-time monitoring via a smartphone-based AI application led to significantly improved medication adherence [19]. This intervention resulted in a 50% increase in adherence rates compared to the standard care control

group, as measured by plasma drug concentration levels.

Our findings also align with previously published results demonstrating the potential of the patient-centered digital wound care technology for remote wound monitoring. For example, a case study by Kong et al. (2021) highlighted the successful application of the DWCS technology in the management of a male patient with type 1 diabetes and multiple comorbidities, including chronic kidney disease and a prior toe amputation [20]. Initially managed for osteomyelitis of a chronic foot ulcer via text and email, the patient transitioned to using the DWCS Patient Connect app for monitoring and management between June 2020 and January 2021. Over seven months, the patient submitted 39 wound images—a nearly twenty-fold increase in the sharing of wound-related data compared to the situation before using the application—enabling the tracking of accurate measurements of two additional wounds. The app fostered patient engagement through weekly assessments, promoting self-examination and preventive behaviors such as infection and trauma monitoring and off-loading of wound pressure through orthotics. Remote follow-ups reduced healthcare visits, alleviating patient anxiety by minimizing direct contact and enhancing physicians' confidence to deliver effective care remotely. Streamlined workflows and the use of images captured during dressing changes further saved time and costs, demonstrating the app's potential to optimize wound management and expand care capacity. The patient also found the app "educational and empowering", highlighting the ability of patient-centred technology to improve patient sentiment and better engage individuals with their wound care treatments.

In Kong and colleagues' case study [20], the assessed patient expressed concerns about sharing wound images via standard messaging platforms, highlighting a common issue with smartphone-based remote care strategies: the security of patient data [21]. Before transitioning to the app, the patient, despite having direct access to their physician, felt that sending images could impose on the physician's time. Additionally, the patient was uncomfortable with the idea that the images would be transmitted through standard messaging and stored on the physician's smartphone, raising privacy and data security concerns. In contrast, by storing images captured using the app on secure cloud-based servers, this reduced the patient's anxiety towards sharing images and facilitated the physician's ability to rapidly and securely receive images.

While the sample size is small, this pilot study provides promising results regarding the use of the Patient Connect app. Our findings demonstrate that the app can be effectively used across various types of wounds and healthcare settings. It has been utilized in hospital departments, such as the Division of Infectious Diseases at the Jewish General Hospital, as well as in ambulatory settings, including ostomy care and pressure ulcer prevention at Centenary Hospital, Scarborough Health Network, and Ontario Health at Home. No adverse outcomes or wound complications were recorded with the use of the Patient Connect application during the study period. No significant privacy or security issues arose as well as the app followed all regulatory protocols regarding data protection. However, a few participants, usually elderly patients, may have

highlighted the need to use assistance in taking pictures of wounds for difficult to reach or seen areas such as the sacrum or back. Lighting conditions also had an effect on the quality of the images, which indicated the need for further instruction or caregiver assistance in cases where optimal image capture was crucial.

Future studies are needed to rigorously evaluate the time savings associated with the use of the app, such as reductions in days lost due to unplanned hospital admissions or the average number of missed workdays. Additionally, research should investigate whether incorporating the app as part of a remote wound care strategy can deliver care that is comparable to or even superior to standard in-person appointments by measuring median days to heal and wound complication rates. Beyond clinical outcomes, the app's potential to reduce patient costs related to travel, time off work, and other logistical burdens associated with frequent healthcare visits highlights its value in remote care settings. As the current study had a three-month follow-up period, which may not fully capture the healing trajectory or wound recurrence for some wound types, an extended follow-up duration is recommended in future studies. Such insights will be critical in validating the app's role in enhancing accessibility, efficiency, and cost-effectiveness in wound care. Additionally, we are currently exploring the potential use cases of our technology for post-surgical sites, aiming to evaluate the effectiveness and feasibility of patient-centered wound images to detect infection. Understanding the potential use cases of generative AI for patient support may also be a worthwhile avenue for further exploration, for example, summarizing the AI/CV analysis of the images captured by patients and providing information on the next steps (e.g., clinician follow-up or continued self-management). AI and CV technology may offer patients and caregivers meaningful tools that empower them to understand better their condition, treatment options, and progress addressing gaps that chronic wounds face due to falling outside of a medical specialty. Further, this study explained and discussed the development of the Patient Connect app for feasible remote wound monitoring. Swift Medical further introduced advanced AI-enhanced features such as AutoDepth and SmartTissue™ to deal with any challenges surrounding the monitoring of complex wounds. For example, AutoDepth identifies wound edges, calculates dimensions, and pinpoints the deepest area of the wound in real-time. SmartTissue™ is capable of quantifying tissue types, namely, epithelial, granulation, slough, and eschar—irrespective of the skin tone (Gupta et al., 2024). These innovations enhance precision, introduce automation, and facilitate clinical decision-making. Future studies should examine the effect of the innovations on patient engagement, complex wound assessment, and treatment outcomes.

Limitations

This study was limited to a targeted patient group of 28 individuals across two hospitals, which may restrict the generalizability of our findings. Additionally, while images were collected from a variety of wound types, further research is needed to evaluate the applicability of the technology for complex versus simple wounds and location of wounds. For example, situations may exist where

caregiver support would be necessary like for wounds in inaccessible locations. However, differences in patient and caregiver technical proficiency with smartphones and applications were not standardized or controlled for as potential confounding factors. Further, understanding the relationship between the technological capability and the app's use, engagement level, and clinical outcome would provide valuable insight. Future studies could help inform the creation of training programs to increase adoption and usability in various patient and caregiver populations. Additionally, the study only included patients using iOS devices, potentially excluding the experience from a broader population who use Android or other platforms. Future research should evaluate the feasibility and usability, as well as the clinical advantages, of an Android-compatible version. Moreover, cross-platform studies comparing user experiences and engagement between iOS and Android users might give insight into possible differences in adoption, functionality, and effectiveness for remote wound monitoring.

Due to the nature of this as a feasibility study, the absence of a control group limits the ability to infer causality from the Patient Connect app to wound healing outcomes. However, feasibility studies are still important as they inform and guide the design of future large-scale trials. The findings from this study, where an observed median wound closure rate was 80%, offer preliminary insights into potential benefits. Such data could facilitate a sample size estimation in an RCT to be run in the future. Sample size calculation suggests that 81 per group (162 total) would be required to have a power of 80% to detect a statistically significant difference between wound healing outcomes in the intervention and standard care without it done with a level of significance of 5% ($\alpha = 0.05$), assuming a healing rate of 60% with standard care without intervention. These findings should be further investigated to understand their validity, as well as some other broader clinical and economic implications.

Conclusion

AI-powered medical tools exhibit tremendous potential in their ability to promote treatment optimization, patient satisfaction, treatment adherence, and overall health outcomes. Our pilot study found numerous clinical benefits using the novel patient-centered, CV-powered mobile application for chronic wound assessment. Similarly, the regular image capture by patients enabled physicians to conduct real-time wound assessments, thereby increasing patient adherence to management plans, as evidenced by an 80% wound closure rate within the participating sample. Considering the potential for technologies like the Patient Connect app to positively impact patient behavior and involvement within their own healthcare treatment journeys by collecting data that benefits their own self-awareness and clinical decision-making, future research should be conducted to understand the clinical, operational, and financial outcomes impacted by patient self-monitoring of wounds and chronic wounds. Factors that would help the widespread adoption of this innovation include more evidence-

based research from larger patient populations to demonstrate the application's effectiveness and benefits in helping deliver remote care, continued user-interface improvements, further maturation of the AI/CV wound assessment technology, patient education on the use of applications and general improvements in specific populations (e.g. the elderly) familiarity with technology, and access to high-speed internet, especially for rural populations.

Disclosure

Patient Connect is a software product of Swift Medical Inc. RDJF, SW & HM are current employees of Swift Medical, and JRG is a former employee of Swift Medical Inc.. All other authors have no disclosures to declare.

Figures and Tables

Variable	N = 28
Average Age	66.4 ± 18.5
Gender	Female = 14 (52%) Male = 13 (48%)
Type of lesion	Diabetic ulcer = 14 (52%) Venous ulcer = 7 (26%) Pressure ulcer = 4 (15%) Postsurgical = 2 (7%)
Median initial wound size (cm²)	3.71 (range 0.48 to 27.91)
Median follow up time (months)	3 (range 1 to 9)
Median number of images submitted	13 (range 4 to 45)
Average time between images (days)	8 (range 3 to 14)
Percentage of wound closure achieved	80% (range 15 to 100%)

Table 1: Patient characteristics. Data is presented as mean ± sd, median and range, or proportions




Figure 1. A simulated wound is used to illustrate the difference between the patient-facing mobile application and the clinician mobile applications. The patient user experience is simplified and provides educational content to support image capture and wound care best practices. The patient image history shows only images and access to information the patient submitted in the documents section, which includes basic screening questions for signs of infection and a free text (third image from the right). The clinician app has standardized documentation for wound assessment, treatment and progress to be documented. The AI/CV outputs are displayed to support clinical decision making.



Figure 2. A case series of a post-operative wound. First image on the left was captured by the clinician. Then the patient was taught to capture images and a second image the same day was documented. The two images on the right half show follow up monitoring submitted by the patient as the wound closed.




Figure 3. A case series of a hard-to-heal wound on the sacrum imaged by a caregiver during the patient journey. Images have adequate lightening, focus, colour correction, and AI/CV-based measurement is shown to the clinician monitoring the wound remotely.

References

1. Bowers S, Franco E. Chronic Wounds: Evaluation and Management. *Am Fam Physician*. 2020;101(3):159-166.
2. Gottrup F. A specialized wound-healing center concept: importance of a multidisciplinary department structure and surgical treatment facilities in the treatment of chronic wounds. *Am J Surg*. 2004;187(5):S38-S43. doi:10.1016/S0002-9610(03)00303-9
3. Mustoe TA, O'Shaughnessy K, Kloeters O. Chronic Wound Pathogenesis and Current

Treatment Strategies: A Unifying Hypothesis: *Plast Reconstr Surg.* 2006;117(SUPPLEMENT):35S-41S. doi:10.1097/01.prs.0000225431.63010.1b

4. Järbrink K, Ni G, Sönnerngren H, et al. Prevalence and incidence of chronic wounds and related complications: a protocol for a systematic review. *Syst Rev.* 2016;5(1):152. doi:10.1186/s13643-016-0329-y

5. Guest JF, Fuller GW, Vowden P, Vowden KR. Cohort study evaluating pressure ulcer management in clinical practice in the UK following initial presentation in the community: costs and outcomes. *BMJ Open.* 2018;8(7):e021769. doi:10.1136/bmjopen-2018-021769

6. Encarnação R, Manuel T, Palheira H, Neves-Amado J, Alves P. Artificial Intelligence in Wound Care Education: Protocol for a Scoping Review. *Nurs Rep.* 2024;14(1):627-640. doi:10.3390/nursrep14010048

7. Dave M, Patel N. Artificial intelligence in healthcare and education. *Br Dent J.* 2023;234(10):761-764. doi:10.1038/s41415-023-5845-2

8. Lindroth H, Nalaie K, Raghu R, et al. Applied Artificial Intelligence in Healthcare: A Review of Computer Vision Technology Application in Hospital Settings. *J Imaging.* 2024;10(4):81. doi:10.3390/jimaging10040081

9. Ethier O, Chan HO, Abdolahnejad M, et al. Using Computer Vision and Artificial Intelligence to Track the Healing of Severe Burns. *J Burn Care Res.* 2024;45(3):700-708. doi:10.1093/jbcr/irad197

10. Le DTP, Pham TD. Unveiling the role of artificial intelligence for wound assessment and wound healing prediction. *Explor Med.* Published online August 31, 2023:589-611. doi:10.37349/emed.2023.00163

11. Mohammed HT, Bartlett RL, Babb D, Fraser RDJ, Mannion D. A time motion study of manual versus artificial intelligence methods for wound assessment. Singh K, ed. *PLOS ONE.* 2022;17(7):e0271742. doi:10.1371/journal.pone.0271742

12. Mohammed H, Mannion D, Cassata A, Fraser R. Trends in Pressure Injury Prevalence Rates and Average Days to Healing Associated With Adoption of a Comprehensive Wound Care Program and Technology in Skilled Nursing Facilities in the United States. *Wounds Compend Clin Res Pract.* 2024;36(1):23-33. doi:10.25270/wnds/23089

13. Wang SC, Anderson JAE, Evans R, et al. Point-of-care wound visioning technology: Reproducibility and accuracy of a wound measurement app. Facchiano A, ed. *PLOS ONE.* 2017;12(8):e0183139. doi:10.1371/journal.pone.0183139

14. Wang SC, Anderson JA, Jones DV, Evans R. Patient perception of wound photography. *Int Wound J.* 2016;13(3):326-330. doi:10.1111/iwj.12293

15. Au Y, Beland B, Anderson JAE, Sasseville D, Wang SC. Time-Saving Comparison of Wound Measurement Between the Ruler Method and the Swift Skin and Wound App. *J Cutan Med Surg.* 2019;23(2):226-228. doi:10.1177/1203475418800942

16. Teresi JA, Yu X, Stewart AL, Hays RD. Guidelines for Designing and Evaluating Feasibility Pilot Studies. *Med Care.* 2022;60(1):95-103. doi:10.1097/MLR.0000000000001664

17. Liyana Arachige M, Tse WC, Zhang R, Ma H, Singhal S, Phan T. Estimating the cost of visiting hospital outpatient. *BMJ Neurol Open.* 2024;6(1):e000576. doi:10.1136/bmjno-2023-000576

18. Trading Economics. (2024). Canada average hourly earnings. Retrieved [02, 14, 2025], from <https://tradingeconomics.com/canada/average-hourly-earnings>

19. Labovitz DL, Shafner L, Reyes Gil M, Virmani D, Hanina A. Using Artificial Intelligence to Reduce the Risk of Nonadherence in Patients on Anticoagulation Therapy. *Stroke*. 2017;48(5):1416-1419. doi:10.1161/STROKEAHA.116.016281
20. Kong LY, Ramirez-GarciaLuna JL, Fraser RDJ, Wang SC. A 57-Year-Old Man with Type 1 Diabetes Mellitus and a Chronic Foot Ulcer Successfully Managed with a Remote Patient-Facing Wound Care Smartphone Application. *Am J Case Rep*. 2021;22. doi:10.12659/AJCR.933879
21. Gunter RL, Chouinard S, Fernandes-Taylor S, et al. Current Use of Telemedicine for Post-Discharge Surgical Care: A Systematic Review. *J Am Coll Surg*. 2016;222(5):915-927. doi:10.1016/j.jamcollsurg.2016.01.062




Supplementary Files

Responses to peer reviewers with in line references to updates made to the manuscript.

URL: <http://asset.jmir.pub/assets/11b26f3a7b127e70e3bcdbcddb6f919a.pdf>

Figures


A simulated wound is used to illustrate the difference between the patient-facing mobile application and the clinician mobile applications. The patient user experience is simplified and provides educational content to support image capture and wound care best practices. The patient image history shows only images and access to information the patient submitted in the documents section, which includes basic screening questions for signs of infection and a free text (third image from the right). The clinician app has standardized documentation for wound assessment, treatment and progress to be documented. The AI/CV outputs are displayed to support clinical decision making.



A case series of a post-operative wound. First image on the left was captured by the clinician. Then the patient was taught to capture images and a second image the same day was documented. The two images on the right half show follow up monitoring submitted by the patient as the wound closed.



A case series of a hard-to-heal wound on the sacrum imaged by a caregiver during the patient journey. Images have adequate lightening, focus, colour correction, and AI/CV-based measurement is shown to the clinician monitoring the wound remotely.



Multimedia Appendixes

Patient Connect instructions for patient or care giver support in adoption of AI-powered wound self-monitoring solution.
URL: <http://asset.jmir.pub/assets/b1a31f7f5297b4f8ff506e1bd18edd3.pdf>



CONSORT (or other) checklists

CONSORT Checklist for the feasibility study.

URL: <http://asset.jmir.pub/assets/ae43219a394b2146931addcf02aa293b.pdf>