

Effects of Digital Sleep Interventions on Sleep Among College Students and Young Adults: A Systematic Review and Meta-Analysis

Yi-An Lu, Hui-Chen Lin, Pei-Shan Tsai

Submitted to: Journal of Medical Internet Research
on: December 05, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 59

 Figures 60

 Figure 1..... 61

 Figure 2..... 62

 Multimedia Appendixes 63

 Multimedia Appendix 1..... 64

 Multimedia Appendix 2..... 64

 Multimedia Appendix 3..... 64

 Multimedia Appendix 4..... 64

 Multimedia Appendix 5..... 64

 Multimedia Appendix 6..... 64

Effects of Digital Sleep Interventions on Sleep Among College Students and Young Adults: A Systematic Review and Meta-Analysis

Yi-An Lu¹ MSN; Hui-Chen Lin^{1,2} PhD; Pei-Shan Tsai^{1,3,4} PhD

¹ School of Nursing, College of Nursing Taipei Medical University Taipei TW

² Research Center in Nursing Clinical Practice Wan Fang Hospital Taipei Medical University Taipei TW

³ Department of Nursing and Center for Nursing and Healthcare Research in Clinical Practice Application Wan Fang Hospital, Taipei Medical University Taipei TW

⁴ Research Center of Sleep Medicine Taipei Medical University Hospital Taipei TW

Corresponding Author:

Pei-Shan Tsai PhD

School of Nursing, College of Nursing
Taipei Medical University
250, Wuxing Street, Xinyi District
Taipei
TW

Abstract

Background: College students and young adults (aged 18-25 years) frequently experience poor sleep quality, with insomnia being particularly prevalent among this population. Given the widespread use of digital devices in the modern world, electronic device-based sleep interventions present a promising solution for improving sleep outcomes. However, their effects in this population remain underexplored.

Objective: To synthesize current evidence on the effectiveness of electronic device-based sleep interventions in enhancing sleep outcomes among college students and young adults.

Methods: A comprehensive search of PubMed, CINAHL, Cochrane Library, Embase, and Web of Science was conducted to identify randomized controlled trials (RCTs) on digital sleep interventions. Pooled estimates of postintervention and follow-up effects were calculated using Hedges' g and 95% confidence intervals (CIs) under a random-effects model. Heterogeneity was assessed with I^2 statistics, and subgroup, moderator, and meta-regression analyses were performed to explore sources of heterogeneity. Evidence quality was evaluated using the Grading of Recommendations, Assessment, Development, and Evaluations framework, and data analysis was performed using Comprehensive Meta-Analysis software, version 3.7.

Results: This study included 13 studies involving 5,251 participants (mean age 23.58 years; 69.55% women). Digital sleep interventions significantly improved subjective sleep quality (Hedges' g = -1.249; 95% CI: -1.834, -0.664), sleep efficiency (Hedges' g = 0.617; 95% CI: 0.182, 1.052), insomnia severity (Hedges' g = -4.082; 95% CI: -5.141, -3.022), dysfunctional beliefs and attitudes about sleep (DBAS; Hedges' g = -1.544; 95% CI: -3.329, -0.986), sleep hygiene (Hedges' g = -0.188; 95% CI: -0.343, -0.032), and sleep knowledge (Hedges' g = -0.268; 95% CI: 0.088, 0.447). The follow-up effects were significant for subjective sleep quality (Hedges' g = -0.532; 95% CI: -0.956, -0.107) and insomnia severity (Hedges' g = -2.649; 95% CI: -3.888, -1.409). The evidence quality ranged from moderate to low certainty across measured outcomes.

Conclusions: Digital sleep interventions are effective in improving subjective sleep quality and reducing insomnia severity, with moderate effects on dysfunctional beliefs and attitudes about sleep, sleep hygiene, and sleep knowledge. These interventions offer a viable approach to managing sleep problems in college students and young adults. Clinical Trial: The protocol for this study is registered with PROSPERO (registration number: CRD 42024595126).

(JMIR Preprints 05/12/2024:69657)

DOI: <https://doi.org/10.2196/preprints.69657>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/>, I will be able to make my accepted manuscript PDF available to anyone.

No. Please do not make my accepted manuscript PDF available to anyone.



Original Manuscript

Effects of Digital Sleep Interventions on Sleep Among College Students and Young Adults: A Systematic Review and Meta-Analysis

Yi-An Lu¹, MSN; Hui-Chen Lin^{1,2}, PhD; Pei-Shan Tsai^{1,3,4}, PhD

¹ School of Nursing, College of Nursing, Taipei Medical University, Taipei, Taiwan

² Research Center in Nursing Clinical Practice, Wan Fang Hospital Taipei Medical University, Taipei, Taiwan

³ Department of Nursing and Center for Nursing and Healthcare Research in Clinical Practice Application, Wan Fang Hospital, Taipei Medical University, Taipei, Taiwan

⁴ Research Center of Sleep Medicine, Taipei Medical University Hospital, Taipei, Taiwan

Corresponding Author:

Pei-Shan Tsai, PhD

School of Nursing, College of Nursing

Taipei Medical University

250, Wuxing Street, Xinyi District

Taipei City, 11031

Taiwan

Phone: +886-2-27361661#6321

Fax: +886-2-2377-2842

Email: ptsai@tmu.edu.tw

Abstract

Background: College students and young adults (≥ 18 -25 years) frequently experience poor sleep quality, with insomnia being particularly prevalent among this population. Given the widespread use of digital devices in the modern world, electronic device-based sleep interventions present a promising solution for improving sleep outcomes. However, their effects in this population remain underexplored.

Objective: To synthesize current evidence on the effectiveness of electronic device-based sleep interventions in enhancing sleep outcomes among college students and young adults.

Methods: Five electronic databases (PubMed, CINAHL, Cochrane Library, Embase, and Web of Science) were searched to identify randomized controlled trials on digital sleep intervention. Sleep interventions including cognitive behavioral therapy for insomnia, mindfulness, and sleep education programs delivered via online platforms or mobile applications, were evaluated for their effects on sleep quality, sleep parameters, and insomnia severity. Pooled estimates of postintervention and follow-up effects were calculated using Hedges' g and 95% confidence intervals (CIs) under a random-effects model. Heterogeneity was assessed with I^2 statistics, and moderator and meta-regression analyses were performed to explore sources of heterogeneity. Evidence quality was evaluated using the Grading of Recommendations, Assessment, Development, and Evaluations.

Results: This study included 13 studies involving 5,251 participants. Digital sleep interventions significantly improved sleep quality (Hedges' $g = -1.25$, 95% CI: -1.83, -0.66, $I^2 = 97\%$), sleep efficiency (Hedges' $g = 0.62$, 95% CI: 0.18, 1.05, $I^2 = 60\%$), insomnia severity (Hedges' $g = -4.08$, 95% CI: -5.14, -3.02, $I^2 = 99\%$), dysfunctional beliefs and attitudes about sleep (DBAS; Hedges' $g = -1.54$, 95% CI: -3.33, -0.99,

$I^2 = 85\%$), sleep hygiene (Hedges' $g = -0.19$, 95% CI: -0.34, -0.03, $I^2 = 0\%$), and sleep knowledge (Hedges' $g = -0.27$, 95% CI: 0.09, 0.45, $I^2 = 0\%$). The follow-up effects were significant for sleep quality (Hedges' $g = -0.53$, 95% CI: -0.96, -0.11, $I^2 = 78\%$) and insomnia severity (Hedges' $g = -2.65$, 95% CI: -3.89, -1.41, $I^2 = 99\%$). Moderator analyses revealed several significant sources of heterogeneity in the meta-analysis examining the effects of digital sleep interventions on sleep outcomes. Variability in sleep quality was influenced by the sleep assessment tool, intervention type and duration, therapist guidance, delivery mode, history of insomnia, and the use of intention-to-treat analysis (all $P < .05$). Heterogeneity in insomnia severity was primarily attributed to differences in the sleep assessment tool ($P < .001$), while the effect size on sleep efficiency varied based on intervention duration ($P = .02$). The evidence quality ranged from moderate to very low certainty across measured outcomes.

Conclusion: Digital sleep interventions are effective in improving sleep quality and reducing insomnia severity, with moderate effects on DBAS, sleep hygiene, and sleep knowledge. These interventions offer a viable approach to managing sleep problems in college students and young adults.

Trial Registration: This review was registered in PROSPERO (CRD 42024595126).

KEYWORDS

Emerging Adults; Insomnia; Technology-Assisted Sleep Interventions; Undergraduate Students

Introduction

College students and young adults aged between 18 and 25 years frequently experience poor sleep quality and insufficient sleep duration [1,2] accompanied by a high prevalence of insomnia. Studies have reported that 18.5% of college students have insomnia, whereas 22.6% of young adults experience difficulty falling asleep; these rates are higher than those observed in the general population [3,4]. Adequate sleep is essential for maintaining overall health, cognitive functioning, and overall well-being in college students and young adults [5,6]. Although the recommended sleep duration for individuals aged 18 to 25 years is 7 to 9 hours per night [7], a study indicated that 30% to 50% of college students and young adults fail to meet this recommendation [8].

Short sleep duration is associated with numerous adverse outcomes in college students and young adults. These include poor dietary habits, obesity [9], and an increased risk of mental health problems (e.g., depression, suicidal behavior, and substance abuse) [10-13]. Furthermore, insufficient sleep adversely affects academic performance and may lead to a need for prolonged years of study [14]. Poor sleep in college students and young adults can also contribute to long-term sleep disturbances that may persist into adulthood [15].

Meta-analyses conducted by Friedrich and Schlarb (2018) and Saruhanjan et al. (2021) have indicated that psychological interventions exert a substantial moderate-to-large effect on sleep-related outcomes in college students [16,17]. In recent years, rapid technological advancements and the widespread adoption of electronic media have drawn attention to the relationship between electronic media use and sleep. A study indicated that college students and young adults spend an average of 316 minutes per week using electronic media, such as smartphones, tablets, emails, and

video platforms [18]. This demographic also engages with the Internet more frequently than do other age groups, rendering it a central aspect of their daily routines [19]. Internet-based interventions have become a practical and promising solution to sleep problems, offering a form of therapy that individuals can access at their preferred time and location [20]. Although face-to-face interventions are highly effective [21], they are often costly, time-intensive, and impractical in many college or other settings [22,23]. By contrast, electronic device-based interventions offer numerous advantages, including convenience, affordability, privacy, ease of distribution, and consistent content delivery [24]. These interventions have demonstrated comparable effectiveness to face-to-face methods in addressing sleep difficulties among college students and young adults [17].

Several studies have systematically reviewed the effectiveness of digital psychological interventions for college students [25-27]. However, these reviews did not specifically evaluate sleep-related outcomes. One review included only two studies on sleep, neither of which focused on sleep as a primary outcome [26], thereby limiting the specificity of the analysis regarding sleep-related interventions. Furthermore, the effects of electronic device-based sleep interventions on sleep outcomes in college students remains underexplored. Four systematic reviews have examined the effectiveness of digital cognitive behavioral therapy for insomnia (CBT-i) [28-31]. Among these reviews, two [30,31] have included college students. Tsai et al. (2022) reviewed four studies, one of which included adolescents and three of which focused on college students. However, certain forms of digital CBT-i, such as interactive therapy and CBT-i delivered using mobile applications, were not analyzed, which considerably limits the applicability of their findings. Additionally, Zhang et al. (2023) analyzed digital sleep interventions in individuals aged 18 years and older with

insomnia or poor sleep quality. Although the review included diverse populations, only two studies specifically focused on college students. The other two reviews [28,29] focused on a broad range of participants aged 18 years and older, including adults and patients with sleep disorders. In addition, these reviews exclusively relied on subjective sleep outcome measures, not including objective data in their analyses [28,29]. Although these studies have provided strong evidence indicating improvements in sleep quality and insomnia symptoms, the heterogeneity in the characteristics of their participants may have affected the internal validity of their results, thereby limiting the generalizability of their findings.

Research indicates that psychological interventions can substantially improve sleep quality and mental health in college students. However, the impact of digital interventions on sleep among college students and young adults remains underexplored. Given the high prevalence of sleep difficulties in this population and their widespread use of digital devices and mobile applications, investigating the effectiveness of digital sleep interventions is both relevant and crucial. The current systematic review and meta-analysis focused on college students and young adults aged 18 to 25 years [32]; this is a critical developmental stage often marked by poor sleep health [33]. Importantly, this review addresses several limitations in the current literature by focusing solely on sleep-related outcomes in college students and young adults. Specifically, we examined both immediate and longer-term effects of digital sleep interventions on subjective and objective sleep measures. By including a wide variety of digital modalities, this review offers clearer insight into how technology-assisted sleep interventions can help improve sleep in this population.

This systematic review and meta-analysis thus aimed to synthesize and quantitatively evaluate current evidence regarding the effectiveness of electronic

device-based sleep interventions in enhancing sleep quality, reducing insomnia severity, and improving sleep outcomes among individuals aged 18 to 25 years, in comparison with control conditions.



Methods

Study Design

This systematic review and meta-analysis of randomized controlled trials (RCTs) investigated the effects of digital sleep interventions on college students and young adults aged 18 to 25. The included interventions were delivered via online platforms or mobile applications. The review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [34], which are detailed in Multimedia Appendix 1. The review protocol was registered with PROSPERO (CRD 42024595126).

Data Sources and Search Strategy

Two reviewers (Y.A.L. and H.C.L.) worked collaboratively to conduct a comprehensive search across five electronic databases, namely PubMed, CINAHL, Cochrane Library, Embase, and Web of Science, to identify relevant studies published from each database's inception to October 2024. The search strategy involved the use of the following keywords: (college OR university AND students OR young adults) AND (digital OR sleep AND intervention OR internet OR web-based OR online OR email OR mobile OR smartphone OR virtual reality) AND (sleep OR sleep quality OR insomnia) AND (randomized controlled trial). Details regarding the search strategies are provided in Multimedia Appendix 2-Table 1. In addition, the reference lists of related articles were manually screened for further eligible studies. No restrictions regarding the language of publication were applied.

Inclusion and Exclusion Criteria

In this systematic review, we included RCTs that met the following inclusion criteria: (1) involved participants who were college/university students or young adults (aged 18-25, as defined by the Society for Adolescent Health and Medicine);

(2) investigated the effects of sleep interventions, delivered through electronic technologies such as the Internet, computers, tablets, mobile devices, telehealth, or virtual reality; (3) included control groups comprising waiting-list control, placebo, no intervention, or standard/usual care groups or individuals receiving interventions delivered face-to-face through in-person coaching; (4) assessed sleep outcomes of interest by using either subjective measurements (e.g., sleep questionnaires or sleep diaries) or objective measurements (e.g., actigraphy); (5) reported both preintervention and postintervention outcome measures. No limitations regarding the language of publication were applied.

Studies were excluded if they: (1) involved participants prescribed medication for sleep; or (2) participants had clinically diagnosed severe sleep disorder (e.g., obstructive sleep apnea, parasomnias, narcolepsy), based on diagnostic criteria from the Diagnostic and Statistical Manual of Mental Disorders-5 (DSM-5), and requiring more intensive treatment; or (3) participants had psychiatric disorders (e.g., major depressive disorder, bipolar disorder, or schizophrenia), diagnosed by a licensed clinician.

Primary and Secondary Outcomes

The primary outcomes of this review were sleep quality, sleep parameters and insomnia severity. The secondary outcomes included dysfunctional beliefs and attitudes about sleep (DBAS), sleep hygiene, sleep knowledge, and pre-sleep arousal in both cognitive and somatic domains.

Study Selection and Data Extraction

Duplicate articles were identified, removed, and screened using EndNote 20 software. Two independent reviewers (Y.A.L and H.C.L) screened the titles and

abstracts of the remaining studies. Full-text reviews were conducted for studies meeting the inclusion criteria. Any disagreements between the two reviewers were resolved through discussion with a third reviewer (P.S.T).

Data were extracted using Excel with a standardized form by two independent reviewers (Y.A.L and H.C.L). Information on the following parameters was collected: study characteristics (e.g., first author, publication year, country, number of participants, mean age, number of female participants, and history of insomnia), intervention type (e.g., duration and delivery mode), postintervention dropout rates, use of intention-to-treat analysis, comparison groups, and outcome measurements (e.g., sleep quality and sleep parameters). Study authors were contacted if any data were missing or incomplete. Available data were used when no response was received from corresponding authors.

Assessment of Methodological Quality

Two researchers (Y.A.L and H.C.L) independently evaluated the quality of the included studies by using the Cochrane Risk of Bias (RoB 2.0) tool for RCTs [35]. This tool is used to evaluate bias across five domains: (1) bias arising from the randomization process, (2) bias due to deviations from the intended intervention, (3) bias due to missing outcome data, (4) bias in the measurement of the outcome, and (5) bias in the selection of reported results. The overall risk of bias was determined by synthesizing findings across these domains, and studies were categorized as having a low risk of bias, some concerns, or a high risk of bias. Discrepancies between the researchers were resolved through discussion or, if necessary, consultation with a third reviewer (P.S.T).

Data Synthesis and Analysis

We conducted a meta-analysis of the pooled quantitative data by using Comprehensive Meta-Analysis software, version 3.7. Effect sizes were calculated on the basis of reported means and standard deviations for each outcome. We analyzed preintervention to postintervention improvements and between-group differences in sleep-related outcomes. In addition, effect sizes for available follow-up measurements were calculated by comparing the intervention and control groups. A minimum of two studies per outcome was required to pool data for quantitative analyses.

A random-effects model was applied to account for anticipated heterogeneity across studies. The primary effect size, Hedges' g was calculated for continuous outcomes and reported with 95% confidence intervals (CIs) [36]. Effect sizes were interpreted as small (0.2), moderate (0.5), or large (0.8) [37]. Heterogeneity was assessed using the I^2 statistic, with values of 0%, 25%, 50%, and 75% indicating no, low, moderate, and high heterogeneity, respectively [38]. Substantial heterogeneity (i.e., $I^2 \geq 50\%$) was explored through moderator analyses and meta-regression. For outcomes with high heterogeneity and significant effects, we performed moderator analyses (for categorical variables) and meta-regression (for continuous variables) to identify potential sources of heterogeneity. Moderator analyses were limited to subgroups represented in at least two studies. A P value of $< .05$ was considered significant for both the moderator and the meta-regression analyses.

To ensure the robustness of our findings, sensitivity analyses were conducted by excluding studies with a high risk of bias and evaluating their effects on the overall results. High-risk studies were defined as those with severe systematic biases, such as unclear reporting of the randomization process, deviations from intended interventions, missing outcome data, problems with outcome measurements, or selective reporting of results [35]. Publication bias was assessed using Egger's test

and a funnel plot [39]. For outcomes reported in at least 10 studies, the trim-and-fill method was applied to estimate the potential impact of missing studies [40].

Assessment of the Certainty of Evidence

The quality of evidence was evaluated using the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE) framework. The quality of evidence was categorized as high, moderate, low, or very low on the basis of a comprehensive assessment of several factors, including risk of bias, inconsistency, indirectness, imprecision, and other relevant considerations [41].

Results

Study Selection

A total of 4,699 records were identified through database searches, and 1,572 duplicates were removed using EndNote 20. Two reviewers (Y.A.L and H.C.L) independently dual-screened the titles and abstracts of the remaining studies. After the titles and abstracts were screened, 3,093 records were deemed irrelevant and excluded. This process resulted in 34 studies being included for full-text review. Of these, 22 did not meet the inclusion criteria and were excluded from the systematic review. In addition, one study was identified through a manual search of reference lists in related articles. Finally, 13 studies were included in the meta-analysis [42-54]. The study selection process is summarized in Figure. 1.

Study Characteristics

The included studies involved a total of 5,251 college students and young adults, with a mean age of 23.6 years and 69.6% of participants being women. Most studies were conducted in the United States ($n = 6$), followed by in the United Kingdom ($n = 3$), Asia ($n = 3$), and the Middle East ($n = 1$). The sample sizes ranged from 40 to 3,755 participants. The average dropout rate was 17.7% in the intervention group and 12.6% in the control group. Nearly 75% of the participants ($n = 3,924$) reported sleep problems at baseline. The diagnostic criteria for assessing sleep problems included the Insomnia Severity Index (ISI) in two studies [52,53], the Sleep Condition Indicator (SCI) in one study [44], and the Sleep Quality Score in one study [49].

Most of the studies employed a two-arm trial design, whereas two studies adopted a three-arm trial design [48,49]. Jones et al. (2020) compared “text message-based sleep intervention” with “placebo” and “no intervention.” Additionally, Kim et al. (2024) compared “virtual reality-based meditation” with “concentration meditation” and “no intervention”.

The effectiveness of digital sleep interventions was evaluated in terms of various outcomes: 15 studies assessed subjective sleep quality, 5 examined objective sleep parameters, 4 evaluated insomnia severity, 2 focused on dysfunctional beliefs and attitudes regarding sleep, 4 assessed sleep hygiene, 3 evaluated sleep knowledge, and 2 analyzed pre-sleep arousal. The instruments used to assess subjective sleep quality included the Pittsburgh Sleep Quality Index (PSQI), the Korean Modified Leeds Evaluation Questionnaire, the Patient Reported Outcomes Measurement Information System (PROMIS), and sleep diaries. Objective sleep parameters were assessed using actigraphy. Insomnia severity was evaluated using the ISI or SCI. DBAS was measured using the Dysfunctional Beliefs and Attitudes about Sleep-16. Sleep hygiene and knowledge were assessed using the Sleep Hygiene Index or the Sleep Hygiene Practice Scale. Pre-sleep arousal was measured using the Pre-Sleep Arousal Scale (PSAS). The details of the included studies are presented in Table 1.

Quality Assessments

Six studies had a low overall risk of bias, three had a high overall risk of bias, and four had some concerns. Among the high-risk studies, two [42,46] did not report their randomization methods. In addition, one of these studies [46] had a high rate of missing data, compromising the integrity of the results because of its excessive data loss without explanation. Another study [47] was classified as high risk because it explicitly stated that blinding of participants and assessors was not feasible.

Four studies were categorized as having some concerns because of unclear information regarding specific domains. These included uncertainties about the randomization process [44], blinding of participants [50,54], and outcome measurement procedures [49]. All studies demonstrated a low risk of bias concerning selective reporting, and none provided sufficient information to evaluate other

potential sources of bias. The details of the risk of bias assessment are illustrated in Figure. 2






Table 1. Summary of characteristics of included studies.

No	Author(s), year, country	No. of participants (I/C)	Age (Mean $\pm$ SD)	Women (%) [n]	History of insomnia	Dropout [PI (%)]	Use of ITT	Comparison	Outcome measurements
1	Baber et al., (2017), USA [42]	43/35	I: 20.67 $\pm$ 2.52 C: 20.33 $\pm$ 1.66	I: 29 (37.2) C: 18 (23.1)	No	I: 0 C: 0	Yes	Passive (waiting list)	1. Actigraphy ^a 2. SHI 3. Sleep quality
2	Denis et al., (2020), UK [43]	99/100	I: 19.73 $\pm$ 2.94 C: 20.22 $\pm$ 5.69	I: 99 (100) C: 100 (100)	No	I: 32.3 C: 22.0	No	Active (puzzles)	1. DBAS-16 2. MCQ 3. PSAS 4. PSQI 5. SCI
3	Freeman et al., (2017), UK [44]	1,891/1,864	I: 24.83 $\pm$ 7.70 C: 24.60 $\pm$ 7.60	I: 1,361 (72.0) C: 1,315 (71.0)	Yes	I: 59.1 C: 38.7	No	Passive (usual practice)	1. ISI 2. SCI 3. Sleep diaries
4	Fucito et al., (2017), USA [45]	21/21	I: 24.83 $\pm$ 7.7 C: 24.60 $\pm$ 7.6	I: 10 (48.0) C: 10 (48.0)	No	I: 9.5 C: 9.5	Yes	Active (healthy behaviors)	1. Actigraphy ^a 2. PSQI 3. PROMIS-SRI-SF 4. Sleep diaries
5	Hershner et al., (2018), USA [46]	254/295	I: 21.90 $\pm$ 4.01 C: 22.00 $\pm$ 4.35	I: 140 (55.8) C: 176 (60.5)	No	I: 42.9 C: 30.2	No	Passive (no intervention)	1. ESS 2. MEQ 3. PSQI 4. SHI
6	Huberty et al., (2019), USA [47]	56/53	I: 20.41 $\pm$ 2.31 C: 21.85 $\pm$ 6.3	I: 36 (41.0) C: 43 (49.0)	No	I: 26.8 C: 11.3	No	Passive (waiting list)	1. PROMIS
7	Jones et al., (2020), USA [48]	53/C ₁ :52 /C ₂ :51	I: 18.16 $\pm$ 0.37 C ₁ : 18.17 $\pm$ 0.38 C ₂ : 18.03 $\pm$ 0.17	I: 38 (79.2) C ₁ : 29 (69.1) C ₂ : 36 (80.0)	No	I: 9.4 C ₁ : 19.2 C ₂ : 11.8	No	C ₁ : Passive (placebo) C ₂ : Passive (no intervention)	1. PSQI 2. SHI
8	Kim et al., (2024), Korea [49]	20/C ₁ :20 /C ₂ :20	I: 21.60 $\pm$ 1.47 C ₁ : 21.75 $\pm$ 1.48 C ₂ : 22.35 $\pm$ 1.35	I: 15 (75.0) C ₁ : 13 (65.0) C ₂ : 13 (65.0)	Yes	I: 0 C ₁ : 0 C ₂ : 0	Yes	C ₁ : Active (concentration meditation) C ₂ : Passive (no intervention)	1. KMLSEQ-LSEQ 2. Actigraphy ^a
No	Author(s), year, country	No. of participants	Age (Mean $\pm$ SD)	Women (%) [n]	History of	Dropout [PI (%)]	Use of	Comparison	Outcome measurements

		(I/C)			insomnia		ITT			
9	Liu et al., (2024), Singapore [50]	30/29	I: 21.6 ± 4.26 C: 22.4 ± 1.31	I: 12 (40.0) C: 9 (31.0)	No	I: 6.7 C: 10.0	No	Active (sleep system)	QA	1. Actigraphy ^a 2. PSQI 3. Sleep diaries
10	Morris et al., (2016), UK [51]	48/47	I: 20.69 ± 2.61 C: 20.27 ± 1.56	I: 29 (60.4) C: 33 (70.2)	No	I: 25.0 C: 6.4	Yes	Passive (waiting list)		1. PSQI
11	Okajima et al., (2022), Japan [52]	24/24	I and C: 19.56 ± 1.56	I: 16 (66.7) C: 16 (66.7)	Yes	I: 12.5 C: 16.7	Yes	Passive (self- monitoring)		1. DBAS-16 2. FIRST 3. ISI 4. PSAS 5. SHPS 6. Sleep diaries
12	Short et al., (2020), USA [53]	32/29	I and C: 19.43 ± 2.04	I and C: 51 (84.0)	Yes	I: 6.3 C: 10.3	No	Active (physical health education)		1. ISI 2. SRBQ
13	Yıkılmaz et al., (2023), Turkey [54]	20/20	I: 20.70 ± 1.21 C: 21.05 ± 0.94	I: 3 (15.0) C: 2 (10.0)	No	I: 0 C: 0	Yes	Passive (no intervention)	(no)	1. PSQI

Table 1. Summary of characteristics of included studies. (*continued*).

C = control group, DBAS-16 = Dysfunctional Beliefs and Attitudes about Sleep-16, ESS = Epworth Sleepiness Scale, FIRST = Ford Insomnia Response to Stress Test, I = intervention group, ISI = Insomnia Severity Index, ITT = intention-to-treat analysis, KMLSEQ = Korean Modified Leeds Evaluation Questionnaire, PI = postintervention, PROMIS = Patient Reported Outcomes Measurement Information System, PROMIS-SRI-SF = Patient Reported Outcomes Measurement Information System Sleep-Related Impairment Short-Form, PSAS = Pre-Sleep Arousal Scale, PSQI = Pittsburgh Sleep Quality Index, QA = question answering, SCI = Sleep Condition Indicator, SD = standard deviation, SHI = Sleep Hygiene Index, SHPS = Sleep Hygiene Practice Scale, SRBQ = Sleep-Related Behaviors Questionnaire, UK = the United Kingdom, USA = the United States of America, ^aObjective measures.

Table 2. Summary of digital intervention features of included studies.

No	Author(s), year, country	Type	Duration	Mode of delivery
1	Baber et al., (2017), USA [42]	Sleep hygiene	1 week (22 min and 33 s)	Online (through PowerPoint)
2	Denis et al., (2020), UK [43]	Virtual therapist-led CBT-i (cognitive techniques, relaxation techniques, sleep hygiene, sleep restriction, stimulus control)	6 weeks (6 sessions/week, 20–25 min/session)	Online (through a video platform)
3	Freeman et al., (2017), UK [44]	CBT-i (cognitive techniques, psychoeducation, relaxation techniques, sleep hygiene, sleep restriction, stimulus control)	10 weeks (6 sessions, average of 20 min/session)	Online (through a website)
4	Fucito et al., (2017), USA [45]	CBT-i (cognitive techniques, psychoeducation, relaxation techniques, sleep hygiene, stimulus control)	4 weeks (intervention consisted of four modules; 1 module/week)	Online (web-based)
5	Hershner et al., (2018), USA [46]	Sleep education (a personality profile, two videos on sleep hygiene and its effects, and information on sleep behavior)	8 weeks (intervention consisted of sleep modules of 20 min)	Online (through a website)
6	Huberty et al., (2019), USA [47]	Mindfulness meditation (body scan, breath focus, and kindness)	8 weeks (at least 10 min/session)	Mobile phone (application)
7	Jones et al., (2020), USA [48]	Health belief model education (theory based)	6 weeks (2 texts [first 2 weeks]; 1 text [second 2 weeks]; 1 text every other day [final 2 weeks])	Online (through a text message)
8	Kim et al., (2024), Korea [49]	Meditation (4 video types: sea, night sky, walking through a forest in the fall, and a green forest)	5 days (30 min/day)	VR-based (through a video)

Table 2. Summary of digital intervention features of included studies. (*continued*).

No	Author(s), year, country	Type	Duration	Mode of delivery
9	Liu et al., (2024), Singapore [50]	Sleep QA system with health coaching	4 weeks (30 min/week)	Online (through text)
10	Morris et al., (2016), UK [51]	CBT-i (psychoeducation, relaxation techniques, sleep hygiene, sleep restriction, stimulus control)	6 weeks (intervention consisted of 7 modules; 20 min/week)	Online (through text and email)
11	Okajima et al., (2022), Japan [52]	CBT-i (cognitive techniques, relaxation techniques, sleep hygiene, sleep restriction, stimulus control)	8 weeks (30 min/session; 8 sessions/week)	Online (through an email with PDF files)
12	Short et al., (2020), USA [53]	Motivation, psychoeducation, behavioral tools, and behavior change modules	1 week (intervention consisted of four modules of 45 min)	Online (through text)
13	Yikilmaz et al., (2023), Turkey [54]	Physiotherapist-led, Internet-based BBAT intervention program	6 weeks (60 min/session; 3 sessions/week)	Videoconferencing

BBAT = basic body awareness therapy, CBT-i = cognitive behavioral therapy for insomnia, QA = question answering,

VR = virtual reality.





Figure 2. Risk of bias graph for included studies

Digital Sleep Interventions for College Students and Young Adults

The components of the digital sleep interventions varied across the included studies. Among the 13 studies, 5 (38.5%) employed CBT-i, 5 (38.5%) focused on sleep-related education or information, 2 (15.4%) incorporated meditation, and 1 (7.7%) used body awareness therapy. The intervention durations ranged from 5 days to 10 weeks, with four studies [43-45,53] also assessing follow-up effects after the intervention period. Approximately half of the studies ($n = 7$, 53.8%) were exclusively conducted in university settings. The detailed results are provided in Table 1, Table 2, and Multimedia Appendix 3-Table 1.

Comparators

The majority of the included studies ($n = 10$, 66.7%) included a passive control group. In these studies, participants in the control group received usual or standard care, were placed on a waiting list, or did not receive any intervention. By contrast, the active control groups participated in sleep programs, received information on sleep education and healthy behaviors, or engaged in concentration meditation. The detailed results are provided in Table 1.

Effectiveness of Digital Sleep Interventions

Sleep Quality

The effects of digital sleep interventions on sleep quality were analyzed for both postintervention and follow-up periods. The pooled results revealed a large and significant postintervention effect (Hedges' $g = -1.25$, 95% CI: -1.83, -0.66, $P < .001$, $I^2 = 97\%$) and a moderate but significant follow-up effect (Hedges' $g = -0.53$, 95% CI: -0.96, -0.11, $P = .014$, $I^2 = 78\%$). Substantial heterogeneity was observed between the studies. The detailed results are provided in Table 3.

Although sleep quality is a critical outcome for college students and young adults, the certainty of evidence for the postintervention effects was rated as “low” (Multimedia Appendix 4-Table 1). This rating was downgraded because of risks of bias, inconsistency, and imprecision but upgraded because of the large magnitude of the effect. For follow-up effects, the evidence was rated as having “moderate” certainty. These findings suggest that digital sleep interventions are likely to significantly improve sleep quality in both postintervention and follow-up assessments.

Sleep Parameters

The pooled effects on sleep parameters, including sleep efficiency (SE), total sleep time (TST), wake after sleep onset (WASO), and the number of awakenings (NWAK), were analyzed. Digital sleep interventions demonstrated a significant medium effect on SE (Hedges' $g = 0.62$, 95% CI: 0.18, 1.05, $P = .005$, $I^2 = 60\%$), with substantial heterogeneity among studies. However, nonsignificant effects were observed for NWAK ($P = .27$), TST ($P = .07$), and WASO ($P = .18$). The detailed results are provided in Table 3.



Although sleep parameters are crucial outcomes, the certainty of the evidence ranged from “very low” to “low” (Multimedia Appendix 4-Table 1). This low rating was because of increased risks of bias, inconsistency, and imprecision primarily resulting from variations in measurement methods for sleep parameters. Overall, the evidence remained highly uncertain regarding the postintervention effects of digital sleep interventions on sleep parameters.


Kim et al., 2024 ^b [49]	-4.11	0.65	0.42	-5.38, -2.83	< .001
Liu et al., 2024 [50]	-0.04	0.27	0.07	-0.56, 0.48	.88
Morris et al., 2016 [51]	-0.55	0.21	0.04	-0.96, -0.14	.01
Okajima et al., 2022 [52]	-7.04	0.72	0.60	-8.55, -5.52	< .001
Short et al., 2020 [53]	-0.20	0.27	0.07	-0.72, 0.32	.45
Yikilmaz et al., 2023 [54]	-1.23	0.34	0.12	-1.90, -0.57	< .001
Total	-1.25	0.30	0.09	-1.83, -0.66	< .001
Test for heterogeneity: Q value = 495.09, df = 14 (<i>P</i> < .001), <i>I</i> ² = 97%					
Test for overall effect: Z = -4.18 (<i>P</i> < .001)					

Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (*continued*).

Study (year)	Statistics for each study					Hedges' g and 95% confidence interval
	Hedges' g	Standard error	Variance	Lower, upper limit	P	
Long-term effect: Follow-up (n = 3)						
Denis et al., 2020 [43]	-0.09	0.19	0.04	-0.47, 0.29	.64	
Freeman et al., 2017 [44]	-0.66	0.05	0.003	-0.78, -0.56	< .001	
Fucito et al., 2017 [45]	-0.93	0.32	0.10	-1.55, 0.30	< .01	
Total	-0.53	0.22	0.05	-0.96, -0.11	.01	
Test for heterogeneity: Q value = 9.01, df = 2 (P = .01), I ² = 78 %						
Test for overall effect: Z = -2.46 (P = .01)						

Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (*continued*).

Sleep Parameters						Hedges' g and 95% confidence interval
Study (year)	Hedges' g	Standard error	Variance	Lower, upper limit	P	
Short-term effect: Postintervention						
A) Number of awakenings (n = 2)						
Baber et al., 2017 [42]	−0.45	0.23	0.05	−0.90, −0.001	.049	
Liu et al., 2024 [50]	−0.004	0.27	0.07	−0.53, −0.52	.99	
Total	−0.25	0.22	0.05	−0.68, 0.19	.27	
Test for heterogeneity: Q value = 1.61, df = 1 (P = .20), I² = 38%						
Test for overall effect: Z = −1.12 (P = .27)						
B) Sleep efficiency (n = 5)						
Baber et al., 2017 [42]	0.21	0.23	0.05	−0.23, 0.65	.35	
Fucito et al., 2017 [45]	1.21	0.33	0.11	0.57, 1.87	< .001	
Kim et al., 2024 ^a [49]	0.31	0.38	0.14	−0.43, 1.06	.41	
Kim et al., 2024 ^b [49]	1.24	0.41	0.17	0.43, 2.04	.003	
Liu et al., 2024 [50]	0.35	0.27	0.07	−0.18, 0.88	.19	
Total	0.62	0.22	0.05	0.18, 1.05	.01	
Test for heterogeneity: Q value = 10.11 df = 4 (P = 0.04), I² = 60%						
Test for overall effect: Z = 2.78 (P = 0.01)						



Insomnia Severity

The pooled effects of digital sleep interventions on insomnia severity, as assessed using the ISI or SCI questionnaires, were analyzed for both postintervention and follow-up periods. The results indicated significant reductions in insomnia severity at postintervention (Hedges' $g = -4.08$, 95% CI: -5.14, -3.02, $P < .001$, $I^2 = 99\%$) and follow-up assessments (Hedges' $g = -2.65$, 95% CI: -3.89, -1.41, $P < .001$, $I^2 = 99\%$). Heterogeneity was observed in insomnia severity outcomes. The detailed results are provided in Table 3.

Insomnia severity is a critical outcome, and the evidence supporting the postintervention and follow-up effects of digital sleep interventions was rated as having a “moderate” level of certainty (Multimedia Appendix 4-Table 1). This rating was downgraded because of inconsistency and imprecision but was then upgraded because of the large magnitude of the observed effects. These findings suggest that digital sleep interventions are likely to significantly reduce insomnia severity in both postintervention and follow-up assessments.

Dysfunctional Beliefs and Attitudes about Sleep

The pooled results of two studies reporting the postintervention effects of digital sleep interventions on DBAS indicated a large and significant effect (Hedges' $g = -2.11$, 95% CI: -3.33, -0.97, $P < .001$, $I^2 = 86\%$). The detailed results are provided in Table 3.

The certainty of evidence for this outcome was rated as “low,” reflecting the limited importance of DBAS as a measure (Multimedia Appendix 4-Table 1). This rating was downgraded due to inconsistency and imprecision in the findings. Although the evidence suggested that digital sleep interventions moderately affected DBAS in postintervention assessments, the clinical significance of this outcome remained uncertain.



Sleep Hygiene

Analysis of the pooled postintervention effects of digital sleep interventions on sleep hygiene revealed a small but significant improvement (Hedges' $g = -0.19$, 95% CI: $-0.34, -0.03$, $P = .02$, $I^2 = 0\%$). The detailed results are provided in Table 3.

The evidence for this outcome was rated as having a “moderate” level of certainty, although sleep hygiene was considered an outcome of limited importance (Multimedia Appendix 4-Table 1). The certainty rating was downgraded because of an increased risk of bias. These findings suggest that digital sleep interventions led to a slight improvement in sleep hygiene following the intervention.

Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (continued).

Insomnia Severity


Study (year)	Statistics for each study					Hedges' g and 95% confidence interval
	Hedges' g	Standard error	Variance	Lower, upper limit	P	
Short-term effect: Postintervention						
A) Insomnia (n = 5)						
Denis et al., 2020 ^c [43]	-10.40	0.63	0.40	-11.64, -9.16	<0.001	
Freeman et al., 2017 ^d [44]	-2.05	0.06	0.003	-2.16, -1.93	<0.001	
Freeman et al., 2017 ^c [44]	-2.95	0.07	0.004	-3.08, -2.82	<0.001	
Okajima et al., 2022 ^d [52]	-7.04	0.77	0.60	-8.55, -5.52	<0.001	
Short et al., 2020 ^d [53]	-0.20	0.27	0.07	-0.72, 0.32	0.45	
Total	-4.08	0.54	0.29	-5.14, -3.02	<0.001	
Test for heterogeneity: Q value = 371.45, df = 4 (P < 0.001), I ² = 99%						
Test for overall effect: Z = -7.55 (P < 0.001)						
Long-term effect: Follow-up						
B) Insomnia (n = 3)						
Denis et al., 2020 ^c [43]	-8.27	0.59	0.35	-9.42, -7.11	<0.001	
Freeman et al., 2017 ^d [44]	-0.66	0.05	0.003	-0.77, -0.56	<0.001	
Freeman et al., 2017 ^c [44]	-2.15	0.07	0.004	-2.27, -2.02	<0.001	
Short et al., 2020 ^d [53]	-0.46	0.30	0.09	-1.04, 0.12	0.12	
Total	-2.65	0.63	0.40	-3.89, -1.41	<0.001	
Test for heterogeneity: Q value = 457.18, df = 2 (P < 0.001), I ² = 99%						
Test for overall effect: Z = -4.19 (P < 0.001)						

^c Sleep Condition Indicator questionnaire, ^d Insomnia Severity Index questionnaire


Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (continued).

Dysfunctional Beliefs and Attitudes about Sleep

Study (year)	Statistics for each study					Hedges' g and 95% confidence interval
	Hedges' g	Standard error	Variance	Lower, upper limit	P	



Test for heterogeneity: Q value = 6.83, df = 1 (*P* = .01), *I*² = 85%
Test for overall effect: Z = -3.68 (*P* < .001)



Test for heterogeneity: Q value = 0.96, df = 3 (*P* = .81), *I*² = 0%
Test for overall effect: Z = -2.36 (*P* = .02)

Sleep Knowledge

The pooled results indicated a small but significant postintervention effect of digital sleep interventions on sleep knowledge (Hedges' $g = -0.27$, 95% CI: 0.09, 0.45, $P = .003$, $I^2 = 0\%$). The detailed results are provided in Table 3.

The evidence for this outcome was rated as having a “moderate” level of certainty, although sleep knowledge was considered an outcome of limited importance (Multimedia Appendix 4-Table 1). The certainty rating was downgraded because of an increased risk of bias. These findings suggest that digital sleep interventions led to a slight improvement in sleep knowledge following the intervention.

Pre Sleep Arousal

The effects of digital sleep interventions on pre sleep arousal, including cognitive and somatic domains, were analyzed. The pooled results indicated nonsignificant postintervention effects for both the cognitive domain (Hedges' $g = -2.02$, 95% CI: -5.86, 1.82, $P = .30$, $I^2 = 98\%$) and the somatic domain (Hedges' $g = -1.90$, 95% CI: -5.04, 1.23, $P = .23$, $I^2 = 98\%$). The detailed results are provided in Table 3.

The certainty of evidence for these outcomes was rated as “low,” reflecting the limited importance of pre sleep arousal (cognitive and somatic domains) as a measure (Multimedia Appendix 4-Table 1). The rating was downgraded because of inconsistency and imprecision in the findings. Overall, the evidence remained uncertain regarding the postintervention effects of digital sleep interventions on pre sleep arousal in both domains.

Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (*continued*).






Table 3. Effect sizes for outcomes of interest in studies evaluating digital sleep interventions (*continued*).





Moderator Analysis

Moderator analysis identified several significant sources of heterogeneity in the effects of digital sleep interventions on sleep quality. Key factors included the type of sleep assessment tool used (PSQI vs. other questionnaires, $P < .001$), intervention type (digital CBT-i vs. other therapies, $P < .001$), intervention duration (≤ 6 weeks vs. > 6 weeks, $P < .001$), therapist guidance (guided vs. unguided, $P < .001$), mode of delivery (email/text message vs. video materials, $P < .01$), history of insomnia (yes vs. no, $P < .001$), and the use of intention-to-treat analysis (yes vs. no, $P = .001$).

Additionally, heterogeneity in insomnia severity outcomes, both postintervention (ISI vs. SCI, $P = .001$) and at follow-up (ISI vs. SCI, $P < .001$), was primarily attributed to differences in the sleep assessment tool. Moreover, variation in sleep efficiency was linked to intervention duration (1 week vs. 4 weeks). The detailed results of the moderator (subgroup) analyses are provided in Multimedia Appendix 5-Table 1.

Meta-regression Analysis

Meta-regression analyses indicated that neither mean age nor the percentage of women significantly contributed to heterogeneity in the effects of digital sleep interventions on various sleep-related outcomes (all $P > .05$).

Specifically, the effect size on sleep quality was not significantly associated with mean age ($\beta = -0.18$, 95% CI: -0.46, 0.11, $P = .23$) or the percentage of women ($\beta = 0.002$, 95% CI: -0.03, 0.03, $P = .90$). Similarly, SE was not significantly influenced by mean age ($\beta = 0.06$, 95% CI: -0.55, 0.68, $P = .85$) or the percentage of women ($\beta = 0.003$, 95% CI: -0.04, 0.04, $P = .90$).

The effect size on WASO also showed no significant association with mean age ($\beta = -1.18$, 95% CI: -6.30, 3.94, $P = .65$) or the percentage of women ($\beta = -0.02$, 95%

CI: -0.05, 0.01, $P = .12$).

Similarly, insomnia severity, both postintervention and at follow-up, was not significantly associated with mean age ($\beta = 0.31$, 95% CI: -0.39, 1.01, $P = .39$; $\beta = 0.02$, 95% CI: -0.36, 0.40, $P = .90$, respectively). The percentage of women also showed no significant association with insomnia severity postintervention ($\beta = 0.13$, 95% CI: -0.02, 0.27, $P = .09$) or at follow-up ($\beta = 0.01$, 95% CI: -0.07, 0.09, $P = .85$).

Finally, SE was not significantly associated with mean age ($\beta = -0.03$, 95% CI: -0.15, 0.08, $P = .54$) or the percentage of women ($\beta = 0.01$, 95% CI: -0.02, 0.04, $P = .39$).

Sensitivity Analysis

Sensitivity analyses were conducted to assess the robustness of the findings of the meta-analysis. Exclusion of the data of three high-risk studies from the pooled data on sleep quality had a minimal impact on the results ($P < .001$), despite the substantial heterogeneity observed in postintervention sleep quality ($I^2 = 97\%$). Additionally, a sensitivity analysis in which the data of one high-risk study were excluded from the pooled data on SE for sleep parameters revealed negligible changes in the results ($P = .004$), further supporting the robustness of the findings. The detailed results are provided in Multimedia Appendix 5-Table 2.

Publication Bias

Publication bias for sleep quality was assessed using Egger's test and a funnel plot. Egger's test indicated no significant publication bias ($P = .24$). The funnel plot is presented in Multimedia Appendix 6.

Discussion

Overview

This systematic review and meta-analysis evaluated the effects of digital sleep interventions on sleep-related outcomes in college students and young adults (aged 18 to 25 years). The findings demonstrate that digital sleep interventions significantly improved sleep quality, SE as a sleep parameter, insomnia severity, DBAS, sleep hygiene, and sleep knowledge. These results indicate the value of evidence-based digital sleep interventions for addressing sleep problems in this population.

Sleep Quality and Insomnia Severity

Our results are consistent with those of previous studies demonstrating that digital sleep interventions [28,31] effectively alleviate sleep disturbances. Reviews have reported moderate-to-large effect sizes for improvements in sleep quality and reductions in insomnia severity. These results are particularly relevant given the high prevalence of poor sleep quality and insomnia among college students and young adults [55,56]. Digital interventions offer accessible and scalable solutions for managing sleep problems. The present review noted that the beneficial effects of digital sleep interventions on sleep quality were sustained for 3 to 6 weeks postintervention, whereas improvements in insomnia severity persisted for up to 1 to 6 months. These findings highlight the potential of digital sleep interventions as a sustainable and effective treatment option for this demographic.

Sleep Parameters

Digital sleep interventions significantly improved SE but demonstrated limited effects on other sleep parameters. These findings are in line with those reported by Maurer et al. (2024) who examined the impact of digital CBT-i on sleep parameters in

individuals with insomnia disorder [57]. The observed improvements in SE suggest that digital interventions can meaningfully enhance the proportion of time spent asleep while in bed. However, no significant effects were found for other objective measures, such as NWAK, TST, and WASO. This lack of consistency may be attributable to the variability in the methods used to measure sleep parameters across studies, which could have affected the outcomes [58]. Future research should prioritize standardizing sleep measures to improve consistency and reliability.

Dysfunctional Beliefs and Attitudes about Sleep

Digital interventions effectively reduced DBAS, consistent with findings reported by Thakral et al. (2020) who indicated the effectiveness of CBT-i in helping individuals develop healthier sleep-related beliefs and attitudes [59]. However, this result contrasts with that reported by Linardon et al. (2024), whose meta-analysis found that smartphone app-based interventions for insomnia and sleep disturbances did not effectively reduce DBAS [60]. A potential explanation for this discrepancy is that the intervention in Linardon et al. did not specifically address participants' individual sleep-related beliefs and attitudes, which may have reduced the persuasiveness and impact of the CBT-i components. Our findings highlight the critical role of CBT-i as a core element in digital sleep interventions, given its demonstrated effectiveness in improving DBAS outcomes [61-63].

Sleep Hygiene and Sleep Knowledge

Although the improvements in sleep hygiene and sleep knowledge were smaller than those for other outcomes, they remain significant. Consistent with findings reported by Bauducco et al. (2020) and Inhulsen et al. (2022), Chehri et al. (2023) demonstrated that sleep hygiene practices are closely linked to sleep quality [64-66].

Promoting better sleep hygiene and increasing awareness of healthy sleep habits are essential strategies for addressing sleep problems. Sleep-related education that promotes knowledge and encourages behavioral changes is particularly effective at addressing sleep problems [67]. Such initiatives are especially valuable for raising awareness and establishing healthier sleep routines among college students and young adults (aged 18-25 years).

Pre Sleep Arousal

The effect sizes for PSAS scores related to pre sleep arousal in both cognitive and somatic domains were nonsignificant, likely because the number of eligible RCTs was limited. Pre sleep arousal is a known predisposing factor for insomnia, with high levels in either domain commonly observed in individuals with sleep disorders, which considerably interferes with the ability to fall asleep [68-69]. Future interventions should incorporate additional components such as mindfulness or relaxation techniques to more effectively address this aspect of sleep disruption.

Moderator

Moderator (subgroup) analyses revealed that digital CBT-i had greater effects on improving sleep quality than did other digital sleep interventions, and therapist-guided interventions outperformed unguided ones. These findings suggest that therapist-guided digital CBT-i is a particularly effective approach to managing sleep problems [28]. Unlike the review by Hasan et al. (2022), which focused on general populations, we incorporated both subjective and objective sleep data and specifically targeted college students and young adults. This focused approach provided deeper insights into the effectiveness of therapist-guided digital CBT-i in improving sleep outcomes for this population.

Strengths and Limitations

To the best of our knowledge, this study is the first to comprehensively investigate the effects of various digital sleep interventions on sleep parameters among college students and young adults. By examining multiple sleep-related outcomes, we provided a holistic understanding of how digital interventions affect different aspects of sleep. However, several limitations of this study should be acknowledged. First, significant heterogeneity was observed across the studies due to variations in intervention regimens and outcome measures, which likely contributed to the inconsistencies in the results. Second, the inclusion of studies with a high risk of bias may have reduced the overall quality of evidence and affected the validity of the findings. However, sensitivity analyses in which high-risk studies were excluded revealed no significant changes in the outcomes, reinforcing the reliability of the results. Third, most studies relied heavily on self-reported measures, which are prone to biases, and objective assessments were underutilized. Finally, the lack of long-term follow-up data in many of the studies limits our ability to provide insights into the sustained impact of these interventions over time.

Relevance for Implementation

We obtained robust evidence indicating that digital sleep interventions are an effective approach to addressing a variety of sleep problems among college students and young adults. These interventions offer a convenient and accessible method for managing sleep disturbances, serving as a low-intensity, cost-effective alternative for individuals who cannot or prefer not to access traditional sleep treatments. By helping users develop effective sleep-related skills, digital interventions can significantly

improve sleep outcomes in this population. However, further research is required to enhance their effectiveness. Personalized digital tools that tailor interventions to individuals' specific sleep patterns and behaviors should be explored. In addition, future RCTs should compare the effectiveness of guided versus unguided digital CBT-i, as therapist involvement may possibly exert attention effects and improve treatment adherence. Future research could benefit from incorporating qualitative approaches to gain deeper insights into user experiences with digital sleep interventions. Additionally, combining both subjective and objective measures, such as actigraphy, would offer a more comprehensive evaluation of the interventions' effectiveness. Longer follow-up periods are also required to enable assessment of the durability of intervention effects, ensuring these tools optimize their potential to improve sleep outcomes for college students and young adults.

Conclusion

This review provides up-to-date evidence on the effectiveness of digital sleep interventions in improving various sleep-related outcomes among college students and young adults. According to the GRADE assessment, the certainty of evidence ranged from moderate to very low. Specifically, moderate certainty was found for sleep quality, insomnia severity, sleep hygiene, and sleep knowledge, while other outcomes had low or very low certainty.

We conclude that digital sleep interventions can improve overall perceived sleep quality and insomnia severity in both postintervention and follow-up assessments. They may also have a slight impact on DBAS, sleep hygiene, and sleep knowledge in postintervention assessments. However, the evidence remains highly uncertain regarding their effects on sleep parameters and somewhat uncertain concerning their impact on pre-sleep arousal in both cognitive and somatic domains. These findings may be influenced by high heterogeneity, variations in interventions and outcome measures, reliance on self-reported data, and the absence of long-term follow-up, which limits the ability to assess the sustained effects of digital sleep interventions. Future research should aim to address these limitations to improve the applicability and effectiveness of digital sleep interventions.

Acknowledgments

None.

Data Availability

All raw data required to produce the aforementioned findings are available upon request.

Authors' Contributions

Yi-An Lu contributed to writing-original draft, visualization, methodology, software, formal analysis, data curation, conceptualization. Hui-Chen Lin contributed to writing-review and editing, validation, software, data curation. Pei-Shan Tsai contributed to writing-review and editing, validation, supervision, conceptualization. All the authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors have no conflicts of interest to declare.

Multimedia Appendix 1

PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) 2020 checklist.

Multimedia Appendix 2

Literature search strategy.

Multimedia Appendix 3

Detailed description of digital sleep interventions based on the TiDIER checklist.

Multimedia Appendix 4

Grading of Recommendations Assessment, Development, and Evaluation (GRADE) summary.

Multimedia Appendix 5

Additional tables (Summary of moderator analysis and sensitivity analyses).

Multimedia Appendix 6

Funnel plot for publication bias.

References

1. Brown FC, Buboltz WC Jr, Soper B. Relationship of sleep hygiene awareness, sleep hygiene practices, and sleep quality in university students. *Behav Med*. 2022; 28(1): 33-8. [doi:10.1080/08964280209596396] [PMID: 12244643]
2. Sivertsen B, Vedaa Ø, Harvey AG, Glozier N, Pallesen S, Aarø LE, et al. Sleep patterns and insomnia in young adults: A national survey of Norwegian university students. *J Sleep Res*. 2019; 28(2): e12790. [doi: 10.1111/jsr.12790] [PMID: 30515935]
3. Jiang XL, Zheng XY, Yang J, Ye CP, Chen YY, Zhang ZG, et al. A systematic review of studies on the prevalence of insomnia in university students. *Public Health*. 2015; 129(12): 1579-84. [doi: 10.1016/j.puhe.2015.07.030] [PMID: 26298588]
4. Kocavska D, Lysen TS, Dotinga A, Koopman-Verhoeff ME, Luijk MP, Antypa N, et al. Sleep characteristics across the lifespan in 1.1 million people from The Netherlands, United Kingdom and United States: a systematic review and meta-analysis. *Nat Hum Behav*. 2021; 5(1): 113-22. [doi:10.1038/s41562-020-00965-x] [PMID: 33199855]
5. Dewald JF, Meijer AM, Oort FJ, Kerkhof GA, Bögels SM. The influence of sleep quality, sleep duration and sleepiness on school performance in children and adolescents: a meta-analytic review. *Sleep Med Rev*. 2010; 14(3): 179-89. [doi: 10.1016/j.smr.2009.10.004] [PMID: 20093054]
6. Jennum P, Bonke J, Clark AJ, Flyvbjerg A, Garde AH, Hermansen K, et al. Søvn og sundhed. København: vidensråd for Forebyggelse, 2015; 1-224. URL: <https://vidensraad.dk/rapport/soevn-og-sundhed#> [accessed 2024-11-04]
7. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep health*. 2015; 1(1): 40-3. [doi:10.1016/j.sleh.2014.12.010] [PMID: 29073412]
8. Lund HG, Reider BD, Whiting AB, Prichard JR. Sleep patterns and predictors of disturbed sleep in a large population of college students. *J Adolesc Health*. 2010; 46(2): 124-32. [doi: 10.1016/j.jadohealth.2009.06.016] [PMID: 20113918]
9. Riemann D, Benz F, Dressle RJ, Espie CA, Johann AF, Blanken TF, et al. Insomnia disorder: state of the science and challenges for the future. *J Sleep Res*. 2022;

- 31(4): e13604. [doi:10.1111/jsr.13604] [PMID: 35460140]
10. Glozier N, Martiniuk A, Patton G, Ivers R, Li Q, Hickie I, et al. Short sleep duration in prevalent and persistent psychological distress in young adults: the DRIVE study. *Sleep*. 2010; 33(9): 1139-45. [doi:10.1093/sleep/33.9.1139] [PMID: 20857859]
 11. Short MA, Booth SA, Omar O, Ostlundh L, Arora T. The relationship between sleep duration and mood in adolescents: a systematic review and meta-analysis. *Sleep Med Rev*. 2020; 52, 101311. [doi:10.1016/j.smrv.2020.10131112] [PMID: 32240932]
 12. Littlewood DL, Kyle SD, Carter LA, Peters S, Pratt D, Gooding P. Short sleep duration and poor sleep quality predict next-day suicidal ideation: an ecological momentary assessment study. *Psychol Med* 2019; 49(3): 403-11. [doi:10.1017/S0033291718001009] [PMID: 29697037]
 13. Russell K, Allan S, Beattie L, Bohan J, MacMahon K, Rasmussen S. Sleep problem, suicide and self-harm in university students: A systematic review. *Sleep Med Rev*. 2019; 44: 58-69. [doi:10.1016/j.smrv.2018.12.008] [PMID: 30721844]
 14. Paudel K, Adhikari TB, Khanal P, Bhatta R, Paudel R, Bhusal S, Basel P. Sleep quality and its correlates among undergraduate medical students in Nepal: A cross-sectional study. *PLOS Glob Public Health*. 2022; 2(2): e0000012. [doi:10.1371/journal.pgph.0000012] [PMID: 36962248]
 15. Dregan A, Armstrong D. Adolescence sleep disturbances as predictors of adulthood sleep disturbances-a cohort study. *J Adolesc Health*. 2010; 46(5): 482-7. [doi:10.1016/j.jadohealth.2009.11.197] [PMID: 20413085]
 16. Friedrich A, Schlärp AA. Let's talk about sleep: a systematic review of psychological interventions to improve sleep in college students. *J Sleep Res*. 2018; 27(1): 4-22. [doi:10.1111/jsr.12568] [PMID: 28618185]
 17. Saruhanjan K, Zarski AC, Bauer T, Baumeister H, Cuijpers P, Spiegelhalder K, et al. Psychological interventions to improve sleep in college students: A meta-analysis of randomized controlled trials. *J Sleep Res*. 2021; 30(1): e13097. [doi:10.1111/jsr.13097] [PMID: 32672865]
 18. Kulturministeriet. Internetbrug og sociale medier.
URL: <https://kum.dk/kulturomraader/medier/mediernes-udvikling/publikationer/internetbrug-og-sociale-medier-2021> [accessed 2024-11-04]
 19. Brautsch LA, Lund L, Andersen MM, Jennum PJ, Folker AP, Andersen S. Digital

- media use and sleep in late adolescence and young adulthood: A systematic review. *Sleep Med Rev.*2023; 68: 101742. [doi:10.1016/j.smrv.2022.101742] [PMID: 36638702]
20. Younes N, Chollet A, Menard E, Melchior M. E-mental health care among young adults and help-seeking behaviors: a transversal study in a community sample. *J Med Internet Res.*2015; 17(5): e123. [doi:10.2196/jmir.4254] [PMID: 25979680]
21. Carey KB, Scott-Sheldon LA, Carey MP, Demartini KS. Individual-level interventions to reduce college student drinking: a meta-analytic review. *Addict Behav.* 2007; 32(11): 2469-94. [doi:10.1016/j.addbeh.2007.05.004] [PMID: 17590277]
22. Carey KB, Scott-Sheldon LA, Elliott JC, Garey L, Carey MP. Face-to-face versus computer-delivered alcohol interventions for college drinkers: a meta-analytic review, 1998 to 2010. *Clin Psychol Rev.* 2012; 32(8): 690-703. [doi:10.1016/j.cpr.2012.08.001] [PMID: 23022767]
23. Nelson TF, Toomey TL, Lenk KM, Erickson DJ, Winters KC. Implementation of NIAAA College Drinking Task Force recommendations: how are colleges doing 6 years later? *Alcohol Clin Exp Res.* 2010; 34(10): 1687-93. [doi:10.1111/j.1530-0277.2010.01268.x] [PMID: 20626728]
24. Leeman RF, Perez E, Nogueira C, Demartini KS. Very-brief, web-based interventions for reducing alcohol use and related problems among college students: a review. *Front Psychiatry.* 2015; 6: 129. [doi:10.3389/fpsy.2015.00129] [PMID: 26441690]
25. Ferrari M, Allan S, Arnold C, Eleftheriadis D, Alvarez-Jimenez M, Gumley A, et al. Digital Interventions for Psychological Well-being in University Students: Systematic Review and Meta-analysis. *J Med Internet Res.* 2022; 24(9): e39686. [doi:10.2196/39686] [PMID: 36169988]
26. Harrer M, Adam SH, Baumeister H, Cuijpers P, Karyotaki E, Auerbach RP, et al. Internet interventions for mental health in university students: A systematic review and meta-analysis. *Int J Methods Psychiatr Res.* 2019; 28(2): e1759. [doi:10.1002/mpr.1759] [PMID: 30585363]
27. Lattie EG, Adkins EC, Winkquist N, Stiles-Shields C, Wafford QE, Graham AK. Digital Mental Health Interventions for Depression, Anxiety, and Enhancement of Psychological Well-Being Among College Students: Systematic Review. *J*

- Med Internet Res. 2019; 21(7): e12869. [doi:10.2196/12869] [PMID: 31333198]
28. Hasan F, Tu YK, Yang CM, James Gordon C, Wu D, Lee HC, et al. Comparative efficacy of digital cognitive behavioral therapy for insomnia: A systematic review and network meta-analysis. *Sleep Med Rev.* 2022; 61: 101567. [doi:10.1016/j.smrv.2021.101567] [PMID: 34902820]
29. Lin W, Li N, Yang L, Zhang Y. The efficacy of digital cognitive behavioral therapy for insomnia and depression: a systematic review and meta-analysis of randomized controlled trials. *Peer J.* 2023; 31(11): e16137. [doi:10.7717/peerj.16137] [PMID: 37927792]
30. Tsai HJ, Yang AC, Zhu JD, Hsu YY, Hsu TF, Tsai SJ. Effectiveness of Digital Cognitive Behavioral Therapy for Insomnia in Young People: Preliminary Findings from Systematic Review and Meta-Analysis. *Pers Med.* 2022; 12(3): 481. [doi:10.3390/jpm12030481] [PMID: 35330480]
31. Zhang H, Yang Y, Hao X, Qin Y, Li K. Effects of digital sleep interventions on sleep and psychological health during the COVID-19 pandemic: A systematic review and meta-analysis. *Sleep Med.* 2023; 110: 190-200. [doi:10.1016/j.sleep.2023.07.036] [PMID: 37625326]
32. Society for Adolescent Health and Medicine. Young Adult Health and Well-Being: A Position Statement of the Society for Adolescent Health and Medicine. *J Adolesc Health.* 2017; 60(6): 758-9. [doi:10.1016/j.jadohealth.2017.03.021] [PMID: 28532650]
33. Taylor DJ, Bramoweth AD, Grieser EA, Tatum JI, Roane BM. Epidemiology of insomnia in college students: Relationship with mental health, quality of life, and substance use difficulties. *Behav Ther.* 2013; 44(3): 339-48. [doi:10.1016/j.beth.2012.12.001] [PMID: 23768662]
34. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021; 372: n71.[doi:10.1136/bmj.n71] [PMID: 33782057]
35. McGuinness LA, Higgins JP. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods.* 2021; 12(1): 55-61. [doi:10.1002/jrsm.1411] [PMID: 32336025]
36. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods.* 2010; 1(2): 97-111. [doi:10.1002/jrsm.12] [PMID: 26061376]

37. Cohen J. Statistical Power Analysis for the Behavioral Sciences. Routledge, New York. 2013. [doi:10.4324/9780203771587]
38. Ahn E, Kang H. Introduction to systematic review and meta-analysis. Korean J Anesthesiol. 2018; 71(2): 103-12. [doi:10.4097/kjae.2018.71.2.103] [PMID:29619782]
39. Lin LF, Chu HT. Quantifying publication bias in meta-analysis. Biometrics. 2018; 74(3): 785-94. [doi:10.1111/biom.12817] [PMID: 29141096]
40. Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics. 2000; 56(2): 455-63. [doi:10.1111/j.0006-341X.2000.00455.x] [PMID: 10877304]
41. Prasad M. Introduction to the GRADE tool for rating certainty in evidence and recommendations. Clinical Epidemiology and Global Health. 2024; 25: 101484. [doi:10.1016/j.cegh.2023.101484]
42. Barber LK, Cucalon MS. Modifying the Sleep Treatment Education Program for Students to include technology use (STEPS-TECH): Intervention effects on objective and subjective sleep outcomes. Stress Health. 2017; 33(5): 684-90. [doi:10.1002/smi.2746] [PMID: 28156049]
43. Denis D, Eley TC, Rijdsdijk F, Zavos HMS, Keers R, Espie CA, et al. Is digital cognitive behavioural therapy for insomnia effective in treating sub-threshold insomnia: a pilot RCT. Sleep Med. 2020; 66: 174-83. [doi:10.1016/j.sleep.2019.10.007] [PMID: 31901759]
44. Freeman D, Sheaves B, Goodwin GM, Yu LM, Nickless A, Harrison PJ, et al. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. Lancet Psychiatry. 2017; 4(10): 749-58. [doi:10.1016/S2215-0366(17)30328-0] [PMID: 28888927]
45. Fucito LM, DeMartini KS, Hanrahan TH, Yaggi HK, Heffern C, Redeker NS. Using Sleep Interventions to Engage and Treat Heavy-Drinking College Students: A Randomized Pilot Study. Alcohol Clin Exp Res. 2017; 41(4): 798-809. [doi:10.1111/acer.13342] [PMID: 28118486]
46. Hershner S, O'Brien LM. The impact of a randomized sleep education intervention for college students. J Clin Sleep Med. 2018; 14(3): 337-47. [doi:10.5664/jcsm.6974] [PMID: 29510791]
47. Huberty J, Green J, Glissmann C, Larkey L, Puzia M, Lee C. Efficacy of the mindfulness meditation mobile app "calm" to reduce stress among college

- students: Randomized controlled trial. *JMIR Mhealth Uhealth*. 2019; 7(6): e14273. [doi:10.2196/14273] [PMID: 31237569]
48. Jones KE, Evans R, Forbes L, Schoenberger YM, Heaton K, Snyder S. Research on freshman and sleeping habits: A text message-based sleep intervention. *J Am Coll Health*. 2020; 68(8): 864-71. [doi:10.1080/07448481.2019.1626860] [PMID: 31237569]
49. Kim KY, Hur MH, Kim WJ. Effects of Virtual Reality (VR)-Based Meditation on Sleep Quality, Stress, and Autonomic Nervous System Balance in Nursing Students. *Healthcare (Basel)*. 2024; 12(16): 1581. [doi:10.3390/healthcare12161581] [PMID: 39201140]
50. Liu J, Ito S, Ngo TM, Lawate A, Ong QC, Fox TE, et al. A pilot randomised controlled trial exploring the feasibility and efficacy of a human-AI sleep coaching model for improving sleep among university students. *Digital Health*. 2024; 10: 20552076241241244. [doi:10.1177/20552076241241244] [PMID: 38638406]
51. Morris J, Firkins A, Millings A, Mohr C, Redford P, Rowe A. Internet-delivered cognitive behavior therapy for anxiety and insomnia in a higher education context. *Anxiety Stress Coping*. 2016; 29(4): 415-31. [doi:10.1080/10615806.2015.1058924] [PMID: 26079158]
52. Okajima I, Tanizawa N, Harata M, Suh S, Yang CM, Li SX, Trockel MT. Can an E-mail-delivered CBT for insomnia validated in the west be effective in the east? A randomized controlled trial. *Int J Environ Res Public Health*. 2022; 19(1): 186. [doi:10.3390/ijerph19010186] [PMID: 35010445]
53. Short N A, Schmidt N B. Developing and Testing a Novel, Computerized Insomnia and Anxiety Intervention to Reduce Safety Aids Among an at-Risk Student Sample: A Randomized Controlled Trial. *Behav Ther*. 2020; 51(1): 149-61. [doi:10.1016/j.beth.2019.05.012] [PMID: 32005332]
54. Yıkılmaz SK, Güloğlu S, Özek Ö, Demirtaş M, Aras GN. Effects of basic body awareness therapy on pain, sleep, quality of life during COVID-19 pandemic-a randomized controlled study. *Annals Clin Analyt Med*. 2023; 14(1): 65-9. [doi:10.4328/acam.21435]
55. Gardani M, Bradford DRR, Russell K, Allan S, Beattie L, Ellis JG, et al. A systematic review and meta-analysis of poor sleep, insomnia symptoms and stress in undergraduate students. *Sleep Medi Rev*. 2022; 61: 101565.

- [doi:10.1016/j.smr.2021.101565] [PMID: 34922108]
56. Schlarb AA, Friedrich A, Claßen M. Sleep problems in university students-An intervention. *Neuropsychiatr Dis Treat*. 2017; 13: 1989-2001. [doi:10.2147/NDT.S142067] [PMID: 28794633]
57. Maurer LF, Bauermann P, Karner L, Müller C, Lorenz N, Giesermann A. Investigating the efficacy of digital cognitive behavioural therapy in comparison to a sleep-monitoring application via integrated diary and actigraphy: A randomized-controlled trial. *J Sleep Res*. 2024; 34(1):e14255. [doi:10.1111/jsr.14255] [PMID: 38895830]
58. Haghayegh S, Khoshnevis S, Smolensky MH, Diller KR, Castriotta RJ. Accuracy of wristband Fitbit models in assessing sleep: Systematic review and meta-analysis. *J Med Internet Res*. 2019; 21(11): e16273. [doi:10.2196/16273] [PMID: 31778122]
59. Thakral M, Von Korff M, McCurry SM, Morin CM, Vitiello MV. Changes in dysfunctional beliefs about sleep after cognitive behavioral therapy for insomnia: A systematic literature review and meta-analysis. *Sleep Med Rev*. 2020; 49: 101230. [doi:10.1016/j.smr.2019.101230] [PMID: 31816582]
60. Linardon J, Anderson C, McClure Z, Liu C, Messer M. The effectiveness of smartphone app-based interventions for insomnia and sleep disturbances: A meta-analysis of randomized controlled trials. *Sleep Med*. 2024; 122: 237-44. [doi:10.1016/j.sleep.2024.08.025] [PMID: 39213858]
61. Okajima I, Akitomi J, Kajiyama I, Ishii M, Murakami H, Yamaguchi M. Effects of a tailored brief behavioral therapy application on insomnia severity and social disabilities among workers with insomnia in Japan: a randomized clinical trial. *JAMA Netw Open*. 2020; 3(4): e202775. [doi:10.1001/jamanetworkopen.2020.2775] [PMID: 32286659]
62. Werner-Seidler A, Li SH, Spanos S, Johnston L, O'Dea B, Torok M, et al. The effects of a sleep-focused smartphone application on insomnia and depressive symptoms: a randomised controlled trial and mediation analysis. *J Child Psychol Psychiatry*. 2023; 64(9): 1324-35. [doi:10.1111/jcpp.13795] [PMID: 36991537]
63. Zhang C, Liu Y, Guo X, Liu Y, Shen Y, Ma J. Digital Cognitive Behavioral Therapy for Insomnia Using a Smartphone Application in China: A Pilot Randomized Clinical Trial. *JAMA Netw Open*. 2023; 6(3): e234866.

- [doi:10.1001/jamanetworkopen.2023.4866] [PMID: 36972049]
64. Bauducco SV, Flink IK, Boersma K, Linton SJ. Preventing sleep deficit in adolescents: Long-term effects of a quasi-experimental school-based intervention study. *J Sleep Res.* 2020; 29(1): e12940. [doi:10.1111/jsr.12940] [PMID: 31691408]
65. Inhulsen MMR, Busch V, van Stralen MM. Effect Evaluation of a School-Based Intervention Promoting Sleep in Adolescents: A Cluster-Randomized Controlled Trial. *J Sch Health.* 2022; 92(6): 550-560. [doi:10.1186/ISRCTN36701918] [PMID: 5315076]
66. Chehri A, Shetabi M, Khazaie H, Zakiei A. Sleep hygiene and sleep quality in Iranian adolescents during the COVID-19 pandemic. *BMC Psychol.* 2023; 11(1): 125. [doi:10.1186/s40359-023-01165-8] [PMID: 37076887]
67. Chung KF, Chan MS, Lam YY, Lai CSY, Yeung WF. School-Based Sleep Education Programs for Short Sleep Duration in Adolescents: A Systematic Review and Meta-Analysis. *J Sch Health.* 2017; 87(6): 4018. [doi:10.1111/josh.12509] [PMID: 28463450]
68. Basishvili T, Oniani N, Sakhelashvili I, Eliazishvili M, Khizanashvili M, Arabidze M, et al. Insomnia, pre-sleep arousal, psychosocial factors and changes in sleep pattern during the second wave lockdown of the covid-19 pandemic in Georgia. *Brain Sci.* 2022; 12(1): 17. [doi:10.3390/brainsci12010017] [PMID: 35053761]
69. Maskevich S, Cassanet A, Allen NB, Trinder J, Bei B. Sleep and stress in adolescents: the roles of pre-sleep arousal and coping during school and vacation. *Sleep Med.* 2020; 66:130-8. [doi: 10.1016/j.sleep.2019.10.006] [PMID: 31877504]

Abbreviations

CBT-i: Cognitive Behavioral Therapy for Insomnia

CI: Confidence Intervals

DBAS: Dysfunctional Beliefs and Attitudes about Sleep

DSM-5: Diagnostic and Statistical Manual of Mental Disorders-5

GRADE: Grading of Recommendations, Assessment, Development, and Evaluations

ISI: Insomnia Severity Index

NWAK: Number of Awakenings

PSAS: Pre Sleep Arousal Scale

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses

PROMIS: Patient Reported Outcomes Measurement Information System

PSQI: Pittsburgh Sleep Quality Index

RCTs: Randomized Controlled Trials

RoB: Risk of Bias

SCI: Sleep Condition Indicator

SE: Sleep Efficiency


TST: Total Sleep Time

WASO: Wake After Sleep Onset


Supplementary Files

Figures

PRISMA flow diagram for study selection.



Risk of bias graph for included studies.



Multimedia Appendixes

PRISMA Checklist.

URL: <http://asset.jmir.pub/assets/37eb93e8c40b362bc09d5abf8950a641.docx>

Literature search strategy.

URL: <http://asset.jmir.pub/assets/2c6293257477a57da56f0f1d98905705.docx>

TiDIER checklist.

URL: <http://asset.jmir.pub/assets/ba78b563800708261f097e70be2db695.docx>

Summary of GRADE assessment for certainty of evidence.

URL: <http://asset.jmir.pub/assets/b1ddcdf3ee68041252fbe5a964b53b85.docx>

Additional tables (Summary of moderator analysis and sensitivity analyses).

URL: <http://asset.jmir.pub/assets/1c0d3dff6582a782a43c2822bd1b0330.docx>

Funnel plot for publication bias.

URL: <http://asset.jmir.pub/assets/af5978a08a8bb5351132672ac813d0ce.docx>