

Stigma Attitudes Towards HIV/AIDS From 2011 Through 2023 in Japan: Retrospective Study in Japan

Yi Piao, Nao Taguchi, Kunihiro Hirahara, Yosuke Takaku, John Austin, KuanYeh Lee, Yui Shiozawa, Yun-Fei Cheng, Yoji Inoue

Submitted to: Journal of Medical Internet Research
on: December 05, 2024

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 52

Figures 53

Figure 1..... 54

Figure 2..... 55

Figure 3..... 56

Figure 4..... 57

Figure 5..... 58

Stigma Attitudes Towards HIV/AIDS From 2011 Through 2023 in Japan: Retrospective Study in Japan

Yi Piao¹ PhD; Nao Taguchi¹ PhD; Kunihiro Hirahara¹; Yosuke Takaku²; John Austin³ PhD; KuanYeh Lee¹ MD; Yui Shiozawa⁴; Yun-Fei Cheng⁴; Yoji Inoue⁵

¹ Gilead Sciences K.K. Tokyo JP

² Japanese Network of People Living with HIV/AIDS Tokyo JP

³ Gilead Sciences (United States) Foster City US

⁴ Deloitte Tohmatsu Consulting LLC Tokyo JP

⁵ Accelight Inc. Tokyo JP

Corresponding Author:

Yi Piao PhD

Gilead Sciences K.K.

Gran Tokyo South Tower 16F, 1-9-2 Marunouchi, Chiyoda-ku

Tokyo

JP

Abstract

Background: Stigma associated with HIV continues to be a major barrier to prevention, testing, treatment, and care despite advances in medical treatments and increased awareness. HIV stigma can influence health behaviors, including reduced likelihood of taking screening tests and decreased adherence to scheduled medical visits. Surveys of the general public in Japan also demonstrated substantial gaps in knowledge of HIV. The content of tweets from the social networking service X (formerly Twitter) has been studied to identify stigmas in mental illness, obesity, and other disorders but has not yet been used to study HIV stigma in Japan.

Objective: To examine the variations and evolution of stigma related to HIV over an extended period using tweets from X, and to investigate the stigma toward people with HIV associated with various demographic segments.

Methods: Japanese tweets related to HIV/AIDS posted on X were retrospectively collected; the Phase 1 feasibility study collected tweets from 2011, 2014, and 2017, and the Phase 2 analysis included tweets from each third year from 2011 through 2023. Individual tweets were labeled with the messages they conveyed (stigma and corresponding antistigma types included: labels, marks, responsibility, peril, insults, and fear; tweets without stigma or antistigma messages were considered general education or neutral) along with demographic characteristics and locations; results collected in Phase 1 were used to develop a machine learning model that was applied in Phase 2. The labeled data from Phase 2 were used to answer the study's research questions concerning yearly changes in HIV stigma and proportions of stigma across population segments.

Results: A total of 2,016,826 tweets related to HIV/AIDS were identified over the study period; 1,648,556 (81.7%) were from individual accounts, and the remainder were from organizational accounts. In total, 574,687 (28.5%) tweets indicated stigma attitudes, while 1,119,852 (55.5%), 207,320 (10.3%), and 114,967 (5.7%) showed neutral, antistigma, or general education attitudes, respectively. The time series showed the proportion of tweets with stigma (approximately 26% in 2011) increased notably during the COVID-19 pandemic and did not return to the starting level in 2023. Tweets including peril, fear, or insult comprised over 80% of tweets with stigma. The greatest number of tweets were made by people in their 20s, whereas people in their 20s and 60s had the greatest proportions of tweets with stigma.

Conclusions: This study contributes to the understanding of the types and attributes of HIV stigma expressed in Japan and shows the usefulness of social media for studying stigma. The extent and type of HIV stigma in Japanese tweets changed from before to after the COVID-19 pandemic. Insights from this study can be used to develop future activities and educational programs to combat HIV-related stigma. Clinical Trial: Not applicable.

(JMIR Preprints 05/12/2024:69696)

DOI: <https://doi.org/10.2196/preprints.69696>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [http://www.jmir.org/](#)

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Stigma Attitudes Towards HIV/AIDS From 2011 Through 2023 in Japan: Retrospective Study in Japan

Yi Piao¹; Nao Taguchi¹; Keisuke Harada¹; Kunihiro Hirahara¹; Yosuke Takaku²; John Austin³; KuanYeh Lee¹; Yui Shiozawa⁴; Yun-Fei Cheng⁴; Yoji Inoue⁵

¹Gilead Sciences K.K., Tokyo, Japan; ²Japanese Network of People Living with HIV/AIDS, Tokyo, Japan;

³Gilead Sciences, Inc., Foster City, CA, USA; ⁴Deloitte Tohmatsu Consulting LLC, Tokyo, Japan;

⁵Accelight Inc., Tokyo, Japan

Abstract (words: 488; limit: 450)

Background: Stigma associated with HIV/AIDS continues to be a major barrier to prevention, testing, treatment, and care despite advances in medical treatments and increased awareness. HIV stigma can negatively influence health behaviors. Surveys of the general public in Japan also demonstrated substantial gaps in knowledge of HIV/AIDS. Tweets from the social networking service X (formerly Twitter) have been studied to identify stigmas in mental illness, obesity, and other disorders but have not yet been used to study HIV stigma in Japan.

Objective: To examine the variations in stigma related to HIV over an extended period using tweets from X and to investigate the stigma toward people with HIV associated with various demographic segments.

Methods: Japanese tweets from X related to HIV/AIDS were retrospectively collected; the Phase 1 feasibility study collected tweets from 2011, 2014, and 2017, and the Phase 2 analysis included tweets from each third year from 2011 through 2023. Individual tweets were labeled with the messages they conveyed (stigma and corresponding antistigma types included: labels, marks, responsibility, peril, insults, and fear; tweets without stigma or antistigma messages were considered general education or neutral) along with demographic characteristics and locations; Phase 1 results were used to develop a machine learning model that was applied in Phase 2. The labeled data from Phase 2 were used to answer the study's research questions concerning yearly changes in HIV stigma and proportions of stigma across population segments.

Results: A total of 2,016,826 tweets related to HIV/AIDS were identified over the study period; 1,648,556 (81.7%) were from individual accounts, with the remainder from organizational accounts. In total, 574,687 (28.5%) tweets indicated stigma attitudes, while 1,119,852 (55.5%), 207,320 (10.3%), and 114,967 (5.7%) showed neutral, antistigma, or general education attitudes, respectively. Tweets including peril, fear, or insult comprised over 80% of tweets with stigma. The

greatest number of tweets were made by people in their 20s, whereas people in their 20s and 60s had the greatest proportions of tweets with stigma (35% of tweets in both age groups). Peril and fear made up 60% of stigma tweets from people in their 20s. The proportion of tweets with stigma (approximately 20% in 2017) increased notably during the COVID-19 pandemic (approximately one-third in 2017), remaining at the same level in 2023. Tweets from health care practitioners had 1.68 times the odds of having antistigma messages vs those from others.

Conclusions: This study contributes to the understanding of HIV stigma expressed in Japan and shows the usefulness of social media for studying stigma. The extent and type of HIV stigma changed from before to after the COVID-19 pandemic. Insights from this study can be used to develop future activities and educational programs to combat HIV-related stigma.

Keywords: HIV; machine learning; stigma; social media; Twitter; X

Introduction

With the availability of effective antiretroviral treatment, mortality among people with HIV (PWH) has decreased considerably; in Japan, the 10-year survival rate for PWH no longer differs significantly from that for the general population [1]. Further, modern antiretroviral therapy is able to suppress viremia sufficiently to prevent sexual transmission of HIV [2]. Yet, public perception may not necessarily reflect such progress. In the 2018 “Public Opinion Survey on HIV Infection and AIDS” of Japanese people randomly selected by the Cabinet Office, approximately 33% of respondents identified AIDS as a disease with no known cause, indicating that the public’s understanding of HIV/AIDS is still considerably insufficient [3]. Moreover, internet search data indicated a decrease in searches relating to HIV test centers in Japan over the course of the COVID-19 pandemic [4]. Despite advances in medical treatments and improved availability of information, the stigma associated with HIV infection continues to be a major barrier to prevention, testing, treatment, and care [5-8]. Perceived stigma may influence PWH to withdraw from social relationships to avoid potential discrimination [9], leading to social isolation and reduced opportunities for social support. In health care settings, stigma decreases medication adherence and reduces trust in health care practitioners (HCPs) among those suffering from HIV stigma [10]. The effects of HIV stigma are not limited to health care behaviors or social interaction but also contribute to an overall poor quality of life [11-13]. HIV stigma can lead to disease progression by delaying care, as shown by the United Nations Development Programme’s correlation between stigma and disease, and many of the people most susceptible to HIV already face stigma, prejudice, and discrimination in their daily lives. This puts them on the margins of society, where poverty and fear make it difficult for them to access health care and HIV services [14]. A study by Yu et al also reported that the prevalence of anticipated stigma was significantly higher among PWH from East Asia (China, Japan, South Korea, and Taiwan) than those from outside Asia [15].

Stigma refers to negative attitudes, beliefs, and behaviors toward a group (for example, PWH) [16]. Stigma can manifest as various *messages*, eliciting *message responses*, and ultimately causing *message effects*, as defined by Smith's stigma-communication model, which can be used to analyze tweets from X (formerly known as Twitter) [17]. A *message* refers to the intent of a tweet, such as a label intended to reinforce societal prejudices. A *message response* is a reaction or interaction generated by another user in relation to the initial message, such as accessing relevant social attitudes or stereotypes. *Message effects* relate to the impact and consequences of these messages and responses on individuals affected by HIV stigma, particularly the development of stigmatizing attitudes [18].

The usefulness of studying the content of X tweets has been reported in recent years [19], and content from X has been employed in studies examining the stigma associated with mental illness, obesity, and other disorders [18, 20, 21]. Ireland et al also showed an association between more frequent "action" tweets and lower HIV incidence in the United States [22]. Japan has the second-highest number of X users across all countries, with approximately 70 million users [23]. However, there have been no studies using X to evaluate attitudes toward HIV in Japan. The sheer volume of posts on X makes it impractical for researchers to personally review a representative portion for analysis. Natural language processing models are increasingly used for health informatics purposes, as they offer the possibility of analyzing volumes of data beyond the scale of practical human labor, including social media output [24]. The adoption of the Bidirectional Encoder Representations from Transformers (BERT) approach allows for the pretraining of models before they are applied to broader tasks [25].

The purpose of this study was to clarify the reality of HIV stigma in Japan by analyzing tweets on the social networking service X.

Methods

Study Phases

The study was conducted in 2 phases. A feasibility study phase was conducted first to explore and contrast the different methodologies of analysis to better understand which approach best answers the research question associated with the study objectives. Then, the outcome of the feasibility study was used as reference to perform an expanded study on a larger dataset obtained from mining tweets collected over a longer period.

Research Objectives

The primary research objective was to examine the variations and transformations in stigma related to HIV over an extended period, and the secondary objective was to analyze the unique stigma associated with demographic segments, including age, sex, and geographic area groups; HCPs or non-HCPs; and individual or organizational accounts.

Ethical Considerations

This study did not directly involve human participants or include any interventions but instead used publicly available tweets. Nonetheless, the investigator submitted this protocol to the independent research ethics committee NPO MINS (Tokyo, Japan) and ensured compliance with the research ethics principles outlined in the Declaration of Helsinki (7th revision, 2013). Before implementation, the investigator received and documented approval from NPO MINS for any modifications made to the protocol. Personal identifying information was limited to information the X users chose to make public.

Inclusion and Exclusion Criteria

The study referenced previous HIV/AIDS infodemiology papers [26-28] to create the initial list of search keywords for obtaining target tweets using the X application programming interface (API) and to develop the methodology for selecting keywords. *HIV*, *AIDS*, and their Japanese synonyms were selected as the keywords. The list was intentionally kept simple to reduce the “noise” that can be generated by using X as a data source.

Certain stop words were selected to remove unrelated data, such as the omission of records with “猫” [cat] to remove unwanted feline immunodeficiency tweets; these stop words are listed in **Table 1**. Unrelated tweets were excluded, as were bot and spam tweets with no identifiable human input.

Table 1. Unrelated topics and corresponding keywords used to exclude tweets from analysis of stigma from 2011 through 2023 in Japan.

Topic	Keywords
Band-aids	● band-aid
Cat	● 猫 (cat) ● 猫 (cat in Hiragana) ● 猫 (cat’s crying sound) ● 猫 (pet owner) ● 猫 (foster parent) ● 猫 (wild cat in Hiragana) ● 猫 (wild cat in Katakana) ● 猫 (wild cat in Kanji)
Chernobyl-aids	● チェルノブイリ (Chernobyl)
Fish/Guppy-aids	● 魚 (fish)

	● 魚 (Guppy) ● 尾 (unit of counting fish)
Hearing	● 助听器 (Hearing-aid machine) ● hearing

Data Collection

Tweet data and metadata, such as profile descriptions and follower counts, were obtained using the X API. Tweet data included the tweeter, the tweet itself, the time of the tweet, and the prefecture (**Table 2**). Demographic segmentation information was extracted from the profile to the extent possible, including age bracket, sex, prefecture, profile text, and profession (HCP or non-HCP). Tweet data and profile data were linked in a master dataset. Proportions of the data unsuitable for the analyses could be removed.

Table 2. Data collected about tweets for categorization in analysis of stigma from 2011 through 2023 in Japan.

Collected Field	Description
tweet_id	A unique combination of integers used to identify all tweets.
tweet_text	The UTF-8 ^a text of the tweet.
is_retweet	A Boolean (true or false) used to identify whether a tweet is original or retweeted. Retweeted tweets may be one of the following: quotation, retweet, or comment.
tweet_author	A unique combination of integers used to identify the author of the tweet. Forms a 1-to-1 relation with the author_id field.
conversation_id	The tweet_id of the original tweet of the conversation (which includes direct replies and replies of replies). Used to reconstruct a conversation or dialogue.

tweet_geo	The geolocation information of the tweet in text format. Available if the author of a tweet included a geolocation tag and NaN ^b otherwise.
tweet_created_time	The UTC ^c of tweet in ISO8601 format.
referenced_tweets_id_list	A list including each tweet_id the associated tweet references. Only available if is_retweet is True, and NaN otherwise.
entities_expanded_url_list	A list of URL ^d entities that have been parsed out of the tweet_text by X.
public_metrics_retweet_count	The integer retweet count of the associated tweet. A retweet occurs when a user shares another user's tweet. Retweets do not include quotation tweets (retweets with a new comment).
public_metrics_reply_count	The integer reply count of the associated tweet. A reply occurs when a user comments under another user's tweet.
public_metrics_like_count	The integer "like" count of the associated tweet. A like occurs when a user clicks the heart button of another user's tweet.
public_metrics_quote_count	The integer quoted count of the associated tweet. A quotation occurs when a user tweets some original text upon referencing/citing another user's tweet.
public_metrics_bookmark_count	The integer bookmarked count of the associated tweet. A bookmark occurs when a user saves another user's tweet for later access.
public_metrics_impression_count	The integer impression count of the associated tweet. An impression occurs when a tweet becomes visible anywhere on a viewer's screen.
author_metrics_followers_count	The integer accumulated follower count of the author of the associated tweet.
author_metrics_following_count	The integer accumulated following count of the author of the associated tweet.
author_metrics_tweet_count	The integer accumulated tweet count of the author of the associated tweet.
author_metrics_listed_count	The integer accumulated list count of the author of the associated tweet. A list indicates a categorization or grouping of X users into specific subject matters.
author_metrics_like_count	The integer accumulated "like" count of the author of the associated tweet.
author_description	The UTF-8 profile text of an author. Also known as a bio.
author_id	A unique combination of integers used to identify the author of the tweet. Forms a 1-to-1 relation with the tweet_author field.
author_name	The UTF-8 text name of the author as defined in the

	profile.
author_username	The UTF-8 text screen name, handle, or alias. Unique to each author but subject to change.
author_created_time	The UTC of user account creation in ISO8601 format.
author_location	The UTF-8 text location of the author. NaN if the author does not provide a text value.

^aUTF: Unicode transformation format.

^bNaN: not a number.

^cUTC: coordinated universal time.

^dURL: uniform resource locator.

Japanese tweet data were available from April 2008 onwards, when X was launched in Japan. However, over the first 3 years, there was little use of the platform in Japan. For this study, 2011 was set as a new initial year, as the use of X in Japan is reported to have substantially increased after the 2011 earthquake off the Pacific Coast of Tohoku [29, 30].

Annotation

Message Category

Labels were applied to each tweet so the analysis result could be grouped and observed based on stigma message and socioeconomic/geographic segments. Tweets were classified into 1 of the message categories: stigma (labels, marks, responsibility, peril, insults, and fear), corresponding antistigma (messages attempting to combat or rectify labels, marks, responsibility, peril, insults, and fear), general education, or neutral, as shown in the illustrative examples in **Table 3**. Based on the categories outlined by Brown et al for investigation using Twitter [18], an additional stigmatic theme called *fear* was identified, as some tweets expressed fear toward HIV/AIDS without necessarily inciting social peril. General education was separated from antistigma as an independent category;

this was used for tweets that may have provided useful information for PWH and/or the general population while not necessarily directly addressing or combating stigma. The initial categories were examined through the labeling processes, and their definitions (the codebook) were updated/ elaborated as needed.

Table 3. Stigma subclassification definitions used from analysis of stigma in tweets from 2011 through 2023 in Japan.

Types of stigma	Definition	Example tweets
Label	- Labels are othering terms used to refer to a stigmatized group - Labels may cause people to consider the stigmatized persons as a distinct group, highlight group differences, or promote stereotypes by highlighting particular characteristics associated with the PSHCD^a	- I don't want to be with homosexuals because I don't want to get AIDS - Black people have quite a lot of HIV infections
Marks	- Marks refer to potentially stigmatizing images, describing ways to identify members of a stigmatized group - To be most effective, marks may be visible physical features, perceived as disgusting, recognized quickly, and remembered	- I was chatting with an infectious disease doctor and the topic of AIDS came up. Apparently, there are HIV specialists who can tell if someone has HIV just by a glance. They say things like, "I mean, even the way they open the door seems suggestive." I wonder if there are cases where a person mimics the behavior or habits of an HIV-negative person and unknowingly contracts HIV - Kaposi's sarcoma all over the face and body. If you see it like that, it's AIDS
Responsibility	Responsibility messages attribute blame	AIDS, huh... This

	or fault to an individual or group because of their experiences of the target PSHCD Such messages may ascribe personal responsibility for the origin, persistence, or severity of the target PSHCD	one is all about personal responsibility. If you want to prevent infection, you have to take precautions yourself. And you really shouldn't go to brothels in the first place. The risk is naturally higher for those who frequent such places compared to the average person. You just have to understand that and make your choices accordingly HIV, because homosexuals do whatever the hell they want to do to each other
Peril	- Peril is content that describes the physical or social threat to a community's effective functioning - To be most effective, the peril may be painful, deadly, and socially taboo	- I heard that more and more foreigners with AIDS virus are coming to Japan and prostituting themselves in Japan. Be careful, sex workers! - A 28-year-old woman infected with HIV is having sex with more than 300 people to get revenge on a man, and she can't be arrested, so she's on the loose, Welcome to the AIDS world
Insults	- Insulting stigma includes messages that are devaluing, derogatory, or abusive toward those living with potentially stigmatized health conditions and disorders - The life of an individual or group may be devalued by content that suggests someone's status, character, credibility, or value may be lessened because of their	Record number of AIDS cases, huh. But the rise in new infections is even more worrying. Increases among people in their 40s and 60s... Gross old folks causing problems

	experiences of the target PSHCD ▪ Derogatory language refers to communications that employ terms associated with the target PSHCD in a disparaging or disrespectful manner ▪ This includes referencing terms associated with the target PSHCD outside of a health context. Abusive communication is content that involves online harassment by direct or indirect reference to the target PSHCD	▪ AIDS lectures, what a pain in the ass. Sex, gays, etc., really annoying
Fear	▪ Fear is the demonstration of baseless personal anxiety toward HIV/AIDS in general or phobia of the people involved ▪ It also involves distrust and discomfort over a situation that is, in fact, within one's control or already contained	▪ I'm scared of AIDS and hemorrhoids, so I don't want to do the real fuck ▪ I'm afraid I'll get HIV if I shake their hand

Based on Brown et al [18].

^aPSHCD: potentially stigmatized health condition or disorder.

A portion of tweets was manually labeled by a group of coders to serve as the teaching data for supervised learning in both the primary and secondary analyses. The methodology was established in accordance with previous research [31] and included sampling and distribution of up to 10 people for labeling. To reduce bias, a small sample of tweets (approximately 1500 tweets per labeler) distributed to the labelers received overlapping treatment, thereby affording them a chance to be labeled by at least 2 people. This promoted consistency in labeling across different labelers. To establish further rigor, the coding team lead was designated in advance and tasked to maintain an audit trail of the manual coding processes. A guiding codebook was developed through interaction with the team lead to ensure agreement. Any contradictory interpretations were resolved through reflective practice and systematic consultations with the entire team. The team could decide to review auxiliary information, such as linked websites and tweets immediately before and after the extracted tweets, in a pre-agreed manner. The discrepancy resolution process was repeated until

80% of the distributed data were labeled consistently across the annotators. After an initial round of labeling based on randomly sampled posts, categories with low frequency (e.g., antistigma) were chosen for additional sampling. Generative AI (ChatGPT) with message definitions included in the prompt was used to screen the posts for annotation. However, all posts were ultimately manually annotated for machine learning training purposes.

Socioeconomic/Geographic Segments

Data segmentation was also performed using a tagging task to identify the age, sex, and other information regarding the user (**Table 4**), using the user profile text as input, to address the secondary objective of examining HIV stigma associated with various demographic segments. The age segment was divided into brackets of 10 years. The segment of geographic area was used instead of prefecture due to a scarcity of tweets with prefecture label; the geographic area segment was divided into metropolis, metropolis peripheral city, medium-sized city, or others. The sex and HCP/non-HCP segments were categorized using binary labels (sex other than male or female could not be included due to lack of representation). Furthermore, the representation and media segments were given particular attention. Subgroup analysis was used to delve into the segmentation of stigmatizing messages. A more detailed examination was performed, if tweet volume permitted, to explore the interplay between different segment combinations or investigate the temporal dynamics of these segments.

Table 4. Segments of categorization from analysis of stigma in tweets from 2011 through 2023 in Japan.

Category	Segment	Label
Binary label	Sex of tweeter	Male/female ^a

	Profession of tweeter	Health care practitioner/not health care practitioner	
	Representation of tweeter	Individual/organization (media/general)	
Multilabel	Age of tweeter	10–19	40–49
		20–29	50–59
		30–39	60+
	Prefecture of tweeter ^b	Tokyo	Kyoto
		Any of the other 47 prefectures of Japan	

^aSex other than male or female was planned for inclusion but could not be included due to lack of representation. ^bA segmentation category of “area” was introduced to gain further insight into stigma levels between urban vs non-urban environments, as not enough tweets with prefecture labels were available, and the results were not necessarily reliable at the prefecture level.

Modeling

Message Category

To fulfill the primary objective of classifying HIV stigma type/message on tweets, the research included “stigma modeling” using a BERT-based large language model (LLM) [32]. Transfer learning from Hugging Face’s publicly available LLM was used to fine-tune this classification task [33]. A stigma model and an antistigma model were developed (each considering the categories of fear, insult, label, etc.). The performance of the models was evaluated using the area under the curve (AUC) based on the receiver operating characteristic curve. An AUC of 1 indicates a perfect model, whereas an AUC of 0.5 signifies a model that is no better than random chance.

Socioeconomic/Geographic Segments

The methodology employed for the segmentation analysis involved constructing models tasked with categorizing the data into specific segments. Models (machine learning or rule based) were built to label each of the segments in the segmentation step for the secondary analyses (**Table 5**). In addition to the BERT model employed for assigning message categories, there was a need for models that are quick to train and lightweight for the purpose of profiling demographics. A random forest model, known for its efficiency and robustness when examining unbalanced data [34], was employed with MeCab tokenization and Tf-idf vectorization to assign demographic segments, including sex, health care profession, and representation. To enhance accuracy, label-identifying keywords and characters were used as separate predictors. For instance, Chinese characters corresponding to “father,” “male,” “uncle,” etc., in English, were chosen for the sex model. We decided to develop simple rule-based models to categorize location and age brackets. A pattern of age or birthday/birth year writing was first identified by examining the profiles, and those that matched the pattern were extracted. For example, numerical values preceding characters corresponding to “year old” and “generation” were extracted. For locations, all possible iterations of a prefecture name and its major city in Chinese characters, Japanese hiragana, katakana, and Roman alphabets were extracted.

Table 5. Models used to classify tweets in analysis of stigma from 2011 through 2023 in Japan.

Task	Model	Input	Processing
Stigma message	Japanese BERT ^a	“tweet_text”	• BERT Japanese tokenization
Sex	Random forest	“author_name” “author_description”	• MeCab tokenization • Tf-idf^b vectorization

			• Add male flag if either input contains sex-identifying keywords: ➤ 男 (Male) ➤ 父 (Father) ➤ 叔叔 (Uncle) ➤ 伯伯 (Uncle) • Add female flag if either input contains sex-identifying keywords: ➤ 女 (Female) ➤ 母 (Mother) ➤ 女人 (Woman)
Health care practitioner	Random forest	"author_name" "author_description"	• MeCab tokenization • Tf-idf vectorization • Add medical flag if either input contains keywords: ➤ 医 (Medical) ➤ 护理 (Nursing care) ➤ 福利 (Welfare) ➤ 医院 (Hospital)
Representation	Random forest	"author_name" "author_description" "author_metrics_followers_count"	• MeCab tokenization • Tf-idf vectorization • Add media flag if author_name or author_description contains keywords: ➤ 新闻 (News) ➤ News ➤ 报纸 (Newspaper) • Add group flag if author_name or

			author_description contains keywords: ➤ ☐ (Department) ➤ ☐ (Association)
--	--	--	--

^aBERT: Bidirectional Encoder Representations from Transformers.

^bTf-idf: term frequency–inverse document frequency.

The performance of all the trained models was evaluated using the AUC based on the receiver operating characteristic curve. For multi-class models, the one-vs-rest scheme was employed. The results were cross-validated to ensure consistent performance across the models. The statistics of the results from rule-based models were evaluated against publicly available information, such as Japanese demography across prefectures, to identify potential bias and to fine-tune the rules.

All models were based on the profiles and only the tweets with the defined HIV-related keywords [35]. The post history of the author accounts, pictures, and other interactions on social media, such as likes, follows, and followers, were not used for this study.

Data Analysis

The next step in the research process involved conducting quantitative analyses. The frequency of message choice was calculated for each message-choice category (labels, marks, responsibility, peril, insult, and fear for stigma and corresponding categories for antistigma). The yearly frequency of each message-choice category was aggregated between 2011 and 2023.

Time-series charts were used to plot the stigma and antistigma counts and proportions of total tweets, providing a clear visualization of the yearly change in public stigma. Subgroup analyses were conducted to investigate the potential intersectionality of stigma levels among the assessed segments. The proportion of stigmatizing and antistigmatizing tweets was calculated for each pair of


compared segments (e.g., male vs female). Fisher's exact test with Holm's method was performed to assess any differences across the segmented subgroups. Multivariate logistic regression was used to examine the characteristics associated with stigma.

Results

Characteristics of Users Posting Tweets

A total of 2,016,826 tweets related to HIV/AIDS were identified (**Figure 1**). Among these, gender could be identified for 1,614,983 (80.1%), age group could be identified for 79,694 (4.0%), and prefecture could be identified for 481,383 (23.9%).

Figure 1. Quantification of tweets identified in analysis of stigma from 2011 through 2023 in Japan.



^aURL: uniform resource locator.

Characteristics of X users included in the study are shown in **Table 6**. Of 2,016,826 tweets, 1,648,556 (81.7%) were from individuals; the remainder came from organizations. Among tweets from individuals, 1,610,060 (97.7%) were from non-HCPs, more tweets were from people in their 20s compared to any other decade (27,331 [34.3%] of the 79,694 whose age group could be

identified), and among those tweets from individuals whose location could be identified (481,383), the majority came from people in a metropolis (296,241 [61.5%]).

Table 6. Characteristics of tweets and X users in analysis of stigma from 2011 through 2023 in Japan.

Attributes		Total: subcategory (proportion of main category)	Neutral: number of tweets (proportion of row)	Stigma: number of tweets (proportion of row)	Antistigma: number of tweets (proportion of row)	General education : number of tweets (proportion of row)
Representation		2,016,826 (100.0%)	1,119,852 (55.5%)	574,687 (28.5%)	207,320 (10.3%)	114,967 (5.7%)
	Individual	1,648,556 (81.7%)	898,186 (54.5%)	513,961 (31.2%)	177,421 (10.8%)	58,988 (3.6%)
	Organization: general	344,801 (17.1%)	204,765 (59.4%)	56,258 (16.3%)	28,291 (8.2%)	55,487 (16.1%)
	Organization: media	23,469 (1.2%)	16,901 (72.0%)	4,468 (19.0%)	1,608 (6.9%)	492 (2.1%)
Sex		1,614,983 (100.0%)	877,197 (54.3%)	506,335 (31.4%)	173,796 (10.8%)	57,655 (3.6%)
	Male	758,667 (47.0%)	401,123 (52.9%)	254,263 (33.5%)	75,960 (10.0%)	27,321 (3.6%)
	Female	856,316 (53.0%)	476,074 (55.6%)	252,072 (29.4%)	97,836 (11.4%)	30,334 (3.5%)
HCP^a/non-HCP		1,648,556 (100.0%)	898,186 (54.5%)	513,961 (31.2%)	177,421 (10.8%)	58,988 (3.6%)
	HCP	38,496 (2.3%)	21,790 (56.6%)	8,723 (22.7%)	6,421 (16.7%)	1,562 (4.1%)
	Non-HCP	1,610,060 (97.7%)	876,396 (54.4%)	505,238 (31.4%)	171,000 (10.6%)	57,426 (3.6%)
Age group		79,694	40,172	22,986	10,053	6,483

		(100.0%)	(50.4%)	(28.8%)	(12.6%)	(8.1%)
	10s	17,254 (21.7%)	7,699 (44.6%)	3,774 (21.9%)	1,338 (7.8%)	4,443 (25.8%)
	20s	27,331 (34.3%)	13,707 (50.2%)	9,650 (35.3%)	3,249 (11.9%)	725 (2.7%)
	30s	14,031 (17.6%)	7,259 (51.7%)	4,052 (28.9%)	2,179 (15.5%)	541 (3.9%)
	40s	10,551 (13.2%)	5,789 (54.9%)	2,654 (25.2%)	1,806 (17.1%)	302 (2.9%)
	50s	8,911 (11.2%)	4,908 (55.1%)	2,298 (25.8%)	1,309 (14.7%)	396 (4.4%)
	60s	1,616 (2.0%)	810 (50.1%)	558 (34.5%)	172 (10.6%)	76 (4.7%)
Area		481,383 (100.0%)	299,682 (62.3%)	115,652 (24.0%)	53,709 (11.2%)	12,340 (2.6%)
	Metropolis	296,241 (61.5%)	182,234 (61.5%)	72,006 (24.3%)	34,775 (11.7%)	7,226 (2.4%)
	Metropolis peripheral city	59,159 (12.3%)	35,173 (59.5%)	15,685 (26.5%)	6,858 (11.6%)	1,443 (2.4%)
	Medium-sized city	86,779 (18.0%)	57,866 (66.7%)	18,778 (21.6%)	8,010 (9.2%)	2,125 (2.4%)
	Others	39,204 (8.1%)	24,409 (62.3%)	9,183 (23.4%)	4,066 (10.4%)	1,546 (3.9%)


^aHCP: health care practitioner.

Characteristics of Tweets

In terms of attitude, 1,119,852 (55.5%) of all tweets were neutral vs 574,687 (28.5%), 207,320 (10.3%), and 114,967 (5.7%) with stigma, antistigma, or general education attitudes, respectively (**Table 6**). The performance of the machine learning models was found as an AUC of 0.72 for the stigma model and 0.77 for the antistigma model (**Figure 2**), indicating moderate performance. The stigma model had a slightly lower performance due to the lack of training data for the mark category (less than 1% incidence rate), though it was within the acceptable range. In the antistigma


model, the mark category was excluded due to its extremely low incident rate. The additional model constructed to classify tweets that were considered neither stigma nor antistigma into general education, irrelevant, or neutral categories had an AUC of 0.92. The proportion of tweets with stigma, approximately 20%, increased significantly during the COVID-19 pandemic; even toward the end of 2023, the proportion of tweets with stigma substantially exceeded the pre-COVID-19 pandemic level (**Figure 3**). Until 2020, absolute numbers rose, the stigma ratio decreased, and the general education ratio increased. After the onset of the COVID-19 pandemic in 2020, except for only a slight increase in general education, all classifications saw a large surge in numbers. By 2023, numbers decreased from 2020, but the stigma ratio remained higher than in 2011. When comparing data before and after the COVID-19 pandemic, stigma rates increased for peril and decreased for insult. In 2020, the numbers of tweets including peril, fear, and responsibility surged, corresponding with the COVID-19 pandemic. Changes in mentions of other stigma were relatively constant after peaking in 2014, although label and insult decreased in absolute numbers relative to their 2014 peak. There were more tweets with stigma messages than antistigma messages at each time point, and the relative frequency of antistigma tweets, compared to stigma tweets, remained relatively consistent before and after the pandemic. Post 2020, the total tweet count surged; it may be the case that antistigma tweets increased in response to the increase in stigma messages during the COVID-19 pandemic. On the other hand, within the breakdown of antistigma types, there was no notable trend before and after the pandemic, and proportions were relatively constant compared to the stigma breakdown ratio.

Figure 2. Receiver operating characteristic curves showing evaluation of (top panel) stigma model and (bottom panel) antistigma model used in analysis of tweets from 2011 through 2023 in Japan.



The performance of the machine learning models was found to have an AUC of 0.72 for the stigma model and 0.77 for the antistigma model, indicating moderate performance. AUC: area under the curve.

Figure 3. Message classification trends for analysis of tweets from 2011 through 2023 in Japan: (top panel) number and percentage of tweets with stigma, neutral, general education, or antistigma labels; (middle panel) number and percentage of tweets with different types of stigma; and (bottom panel) number and percentage of tweets with type of antistigma label. Toward the end of 2023, the proportion of tweets with stigma substantially exceeded the pre-COVID-19 pandemic level.



In terms of stigma subcategory, peril (47.3%) was the most prevalent, followed by fear (27.6%) and insult (12.6%), but the proportion of stigma categories varied depending on the attributes of the individuals posting the tweets (**Table 7**). The proportion of tweets, including fear, was greater among HCPs than among non-HCPs (31.8% vs 27.6%). Among X users in their 20s, 9.1%, 0.1%,

17.4%, 21.7%, 38.6%, and 13.1% of tweets with stigma included label, mark, responsibility, peril, fear, and insult, respectively, vs 6.1%, 0.2%, 4.7%, 47.3%, 36.7%, and 5.0% for X users in their 60s.

Table 7. Stigma and attribute classification from analysis of stigma in tweets from 2011 through 2023 in Japan.

Attribute		Stigma total: subcategory (proportion of main category)	Label: # of tweets (% of row)	Mark: # of tweets (% of row)	Responsibility: # of tweets (% of row)	Peril: # of tweets (% of row)	Fear: # of tweets (% of row)	Insult: # of tweets (% of row)
Representation		574,687 (100.0%)	31,124 (5.4%)	733 (0.1%)	40,696 (7.1%)	271,614 (47.3%)	158,328 (27.6%)	72,192 (12.6%)
	Individual	513,961 (89.4%)	26,070 (5.1%)	652 (0.1%)	38,111 (7.4%)	238,103 (46.3%)	142,274 (27.7%)	68,751 (13.4%)
	Organization: general	56,258 (9.8%)	4,656 (8.3%)	50 (0.1%)	2,410 (4.3%)	31,093 (55.3%)	15,148 (26.9%)	2,901 (5.2%)
	Organization: media	4,468 (0.8%)	398 (8.9%)	31 (0.7%)	175 (3.9%)	2,418 (54.1%)	906 (20.3%)	540 (12.1%)
Gender		506,335 (100.0%)	25,986 (5.1%)	651 (0.1%)	37,983 (7.5%)	236,012 (46.6%)	139,777 (27.6%)	65,926 (13.0%)
	Female	254,263 (50.2%)	12,793 (5.0%)	346 (0.1%)	18,212 (7.2%)	117,528 (46.2%)	65,952 (25.9%)	39,432 (15.5%)
	Male	252,072 (49.8%)	13,193 (5.2%)	305 (0.1%)	19,771 (7.8%)	118,484 (47.0%)	73,825 (29.3%)	26,494 (10.5%)
HCP/non-HCP^a		513,961 (100.0%)	26,070 (5.1%)	652 (0.1%)	38,111 (7.4%)	238,103 (46.3%)	142,274 (27.7%)	68,751 (13.4%)
	HCP	8,723 (1.7%)	419 (4.8%)	17 (0.2%)	507 (5.8%)	4,236 (48.6%)	2,772 (31.8%)	772 (8.9%)
	Non-	505,238	25,651	635	37,604	233,867	139,502	67,979

	HCP	(98.3%)	(5.1%)	(0.1%)	(7.4%)	(46.3%)	(27.6%)	(13.5%)
Gener- ation		22,986 (100.0%)	1,777 (7.7%)	26 (0.1%)	2,869 (12.5%)	7,196 (31.3%)	8,220 (35.8%)	2,898 (12.6%)
	10s	3,774 (16.4%)	220 (5.8%)	3 (0.1%)	294 (7.8%)	1,450 (38.4%)	1,172 (31.1%)	635 (16.8%)
	20s	9,650 (42.0%)	881 (9.1%)	7 (0.1%)	1,678 (17.4%)	2,095 (21.7%)	3,724 (38.6%)	1,265 (13.1%)
	30s	4,052 (17.6%)	327 (8.1%)	8 (0.2%)	482 (11.9%)	1,178 (29.1%)	1,555 (38.4%)	502 (12.4%)
	40s	2,654 (11.5%)	191 (7.2%)	2 (0.1%)	213 (8.0%)	1,119 (42.2%)	877 (33.0%)	252 (9.5%)
	50s	2,298 (10.0%)	124 (5.4%)	5 (0.2%)	176 (7.7%)	1,090 (47.4%)	687 (29.9%)	216 (9.4%)
	60s	558 (2.4%)	34 (6.1%)	1 (0.2%)	26 (4.7%)	264 (47.3%)	205 (36.7%)	28 (5.0%)
Area		115,652 (100.0%)	5,713 (4.9%)	193 (0.2%)	7,408 (6.4%)	57,813 (50.0%)	32,811 (28.4%)	11,714 (10.1%)
	Metrop- olis	72,006 (62.3%)	3,518 (4.9%)	134 (0.2%)	4,639 (6.4%)	35,540 (49.4%)	20,832 (28.9%)	7,343 (10.2%)
	Metrop- olis periphe- ral city	15,685 (13.6%)	756 (4.8%)	27 (0.2%)	990 (6.3%)	7,973 (50.8%)	4,365 (27.8%)	1,574 (10.0%)
	Mediu- m-sized city	18,778 (16.2%)	908 (4.8%)	26 (0.1%)	1,155 (6.2%)	9,753 (51.9%)	5,151 (27.4%)	1,785 (9.5%)
	Others	9,183 (7.9%)	531 (5.8%)	6 (0.1%)	624 (6.8%)	4,547 (49.5%)	2,463 (26.8%)	1,012 (11.0%)


^aHCP: health care practitioner.

Characteristics Associated With Stigma

Models constructed to classify sex, individuals vs organizations, and HCPs vs non-HCPs had AUCs of 0.78, 0.89, and 0.94, respectively (models for age brackets and location were rule based, and no AUCs applied). **Figure 4** shows characteristics associated with a stigma message. Tweets from organizations were less likely than tweets from individuals to be associated with stigma (odds ratios [OR]s and 95% confidence intervals [CIs] of 0.43 [0.43–0.43] and 0.52 [0.50–0.54] for general organizations and media organizations, respectively, vs individuals). Tweets from people in their 20s, 30s, or 60s were all more likely to bear a stigma message vs tweets from people in their 40s (ORs [95% CIs] of 1.62 [1.54–1.71], 1.21 [1.14–1.28], and 1.57 [1.40–1.76], respectively). Tweets from organizations were more likely to be associated with the label stigma than were tweets from individuals (ORs [95% CIs] of 1.69 [1.63–1.74] and 1.83 [1.65–2.03] for general organizations and media organizations, respectively, vs individuals). Tweets from people in their 20s and 30s were more likely than tweets from people in their 40s to bear responsibility stigma (ORs [95% CIs] of 2.41 [2.07–2.81] and 1.55 [1.30–1.84], respectively). Tweets from organizations were more likely to bear the stigma of peril (ORs [95% CIs] of 1.43 [1.41–1.46] for general organizations vs individuals and 1.37 [1.29–1.45] for media organizations vs individuals). Tweets from organizations were less likely to be associated with fear stigma vs those from individuals (ORs [95% CIs] of 0.96 [0.94–0.98] and 0.66 [0.62–0.72] for general organizations and media organizations vs individuals, respectively). Tweets from people in their 20s, 30s, and 60s were more likely to be associated with fear stigma than those from people in their 40s (ORs [95% CI] of 1.27 [1.16–1.40], 1.26 [1.14–1.40], and 1.18 [0.97–1.43],

respectively). Tweets from people in a metropolis or the peripheral cities of a metropolis were more likely to bear fear stigma than tweets from people in other locations (ORs [95% CIs] of 1.12 [1.08–1.17] and 1.06 [1.01–1.12], respectively). Tweets from people in their teens, 20s, and 30s were more likely to bear insult stigma than tweets from people in their 40s (ORs [95% CIs] of 1.93 [1.65–2.26], 1.44 [1.25–1.67], and 1.35 [1.15–1.59], respectively).


Figure 4. Characteristics associated with stigma messages in analysis of tweets from 2011 through 2023 in Japan: (top center) overall; (top of left column) label; (middle of left column) mark; (bottom of left column) responsibility; (top of right column) peril; (middle of right column) fear; and (bottom of right column) insult.



CI: confidence interval; HCP: health care practitioner; OR: odds ratio.

Tweets from general organizations were more likely than tweets from individuals to have a general education perspective (OR [95% CI] of 5.17 [5.11–5.23]), whereas tweets from media organizations were less likely than tweets from individuals to have a general education perspective (0.58 [0.53–0.63]; **Figure 5**). Tweets from HCPs were more likely than tweets from non-HCPs to have an antistigma perspective (OR [95% CI] of 1.68 [1.64–1.73]), whereas tweets from people in a metropolis were more likely than tweets from people in other areas to have an antistigma perspective 1.14 [1.11–1.17].

Figure 5. Characteristics associated with (top) general education and (bottom) antistigma in analysis of tweets from 2011 through 2023 in Japan.



CI: confidence interval; HCP: health care practitioner; OR: odds ratio.

Discussion

Principal Results

In this retrospective study of tweets from Japan mentioning HIV/AIDS from 2011 through 2023 processed using machine learning, the greatest proportion of tweets were neutral; however, there were more tweets with stigma messages than antistigma messages. The tweets were overwhelmingly from individuals in metropolitan areas, which is consistent with expectations for social media. Disappointingly, the number of tweets with HIV-related stigma rose over the course of the study along with the increasing number of X users in Japan. The details identified in this study of population segments most likely to express stigma and the type of stigma expressed can help inform more effective educational activities in the future.

HIV-related tweets were predominantly posted by people in their 20s, indicating that young people may be more concerned about sexually transmitted infections such as HIV than are older people; however, it should be noted that X is a social networking site that may attract young people. Stigma increased during the COVID-19 pandemic (2020 vs 2011), including the percentage of tweets with the peril label. Conspiracy theories such as HIV being contained in the COVID-19 vaccine were spread on social networking sites during this period [36, 37], and these tweets could be a possible reason for the increase in HIV-related tweets.

In tweets from both people in their 20s and those in their 60s, approximately 35% contained stigma statements, the highest proportions among age groups. Stigma statements were more common among people in peripheral cities vs other locations; this may suggest directions for educational activities. For example, young people's limited knowledge of sexual health and lack

of understanding of HIV/AIDS may lead to stigma, such as peril and fear [38]. The high number of tweets may indicate this demographic could benefit from more educational activities. It should be noted that the people in their 20s whose tweets were recorded in 2011 were not the same people in their 20s whose tweets were recorded in 2023, and these different groups of people could have different relationships both to HIV and social media.

The label category made up a proportionately greater part of antistigma tweets than stigma tweets. This could represent a desire to avoid othering sexual minority groups. Some tweets condemned discrimination against people infected with COVID-19, citing past discrimination against PWH and lesbian/gay/bisexual/transgender/queer people, which could be a contributing factor.

Although the percentage and number of tweets from HCPs with stigma statements were small, they may have more impact than those from non-HCPs and should be considered important. It may be worthwhile to develop ways to amplify the influence of HCPs who transmit antistigma statements, for example. Organizations that promote awareness (including professional and academic institutions dedicated to infectious disease) could work together and/or in parallel with HCPs who wish to disseminate antistigma messages addressing all types of stigma. In addition, misinformation has spread, leading to HIV being conflated with COVID-19, including the false rumor that HIV was contained in COVID-19 vaccines [39]. Therefore, it is necessary to disseminate research information with full consideration of the literacy of the recipients of the information.

Comparison With Prior Work

The percentage of stigma messages in this study (28%) was similar to that seen in English-speaking countries [18]. Brown et al reported that label (34%) and insult (31%) were particularly high among stigma messages in English-language tweets [18], whereas peril (47%) and fear (28%) were high in the present study, suggesting the nature of HIV stigma may differ between the populations. Differences in stigma perceptions across nations have been found in previous research; in a global survey, concern about disclosing one's HIV status was notably higher in the Asia-Pacific region vs North America [40]. In addition, since the incidence rate is lower in Japan than in other countries [41], HIV is perceived as a "foreign" disease [42], and the overall awareness of and familiarity with the disease is low; therefore, members of the general public may not be aware of any PWH in their lives, and this unfamiliarity can readily lead to fear and misunderstanding. Based on the results of the "Public Opinion Survey on HIV Infection and AIDS" [3], it is highly possible that although medical advances have made drugs more effective for treating the disease, the concept of "undetectable equals untransmittable" (U=U) is not yet sufficiently widespread among the general public. More detailed studies on the reasons for differences in stigma across countries are needed. Previous reports showed that understanding U=U improves the acceptance of PWH [43], so disseminating this concept and other accurate information via influencers may be a solution to reduce stigma among younger generations. Conducting regular awareness campaigns such as short movies illustrating the U=U concept on social networking sites may help, as such approaches were effective in reducing COVID-19-related stigma [44].

This study has several limitations. User bias is a potential limitation; although Japan has

many X users [23], those users do not necessarily accurately represent the entirety of society. The user attributes recorded are based on self-declaration by the users and cannot be verified; both intentional misrepresentation and outdated information (e.g., ages in profiles) may be present, and there is potential for bias between users who choose to self-disclose information and users who do not. Further, the selection of tweets in Japanese may not necessarily represent attitudes of Japan, as tweets in any language may be posted from different places. There are cases where certain users tweet very frequently (bots, advertisements, etc.), which may influence the statistical results, although we attempted to eliminate consideration of such accounts. An additional limitation includes potential bias in the sample of tweets. HIV-related tweets were limited to those published before 2024 (and not deleted at the time of data retrieval) with specific keywords, and potentially HIV-related tweets that did not mention specific keywords were not included in the study. In particular, antistigma may be tweeted in the form of replies, in which case, there is likely no mention of keywords, which may lead to the underestimation of antistigma tweets. Lastly, there is the possibility of classification bias. Tweet classification was carried out through discussions within the team, but there were tweets with unclear intentions or tweets that were difficult to interpret, and decisions ultimately depended on the interpretation of individuals who annotated a certain label. Furthermore, the trained models and rule-based models are imperfect, with the AUC values reported (0.72 for the stigma model and 0.77 for the antistigma model) indicating moderate performance; there may be some misclassification, which may not be random and therefore biased toward a particular category. Future work can refine the training dataset by oversampling underrepresented categories or using transfer learning from models pretrained on larger datasets.

Conclusion

This study contributes to a better understanding of the types and attributes of HIV stigma in Japan and illustrates the usefulness of social media for describing stigma. The results indicate that HIV stigma is still prevalent, particularly among young adults, who were not alive during the early days of the AIDS panic. This information can be useful in the future development of more effective educational activities. To respond more accurately to the Japanese public's educational needs in the future, it may be important to clarify how better to reach these X users (i.e., which influencers they follow and support, what topics they are interested in) and how to approach them to consider more efficient educational activities in the future.

Acknowledgments

This study was funded by Gilead Sciences K.K. Funding for this analysis and for the Rapid Service Fee was provided by Gilead Sciences K.K. The sponsors participated in the planning, execution, and interpretation of the research. The authors acknowledge the contributions of Yusuke Hirabuki, Yuki Hori, and Taichi Kido for the manual labeling of tweets to develop the model for this study.

Medical writing support was provided by Rob Coover, MPH, of Red Nucleus, and funded by Gilead Sciences K.K.

Conflicts of Interest

YP, NT, Keisuke Harada, Kunihiro Hirahara, and KL are employees of Gilead Sciences K.K. and shareholders of Gilead Sciences, Inc.

JA is an employee of Gilead Sciences, Inc.

YS and **YFC** are employees of Deloitte Tohmatsu Consulting LLC.

YI and **YT** report no conflicts of interest.

Abbreviations

API, application programming interface

AUC, area under the curve

BERT, Bidirectional Encoder Representation from Transformers

CI, confidence interval

HCP, health care practitioner

LLM, large language model

OR, odds ratio

PWH, people with HIV

Data Sharing

Gilead Sciences shares anonymized individual patient data upon request or as required by law or regulation with qualified external researchers based on submitted curriculum vitae and reflecting non-conflict of interest. The request proposal must also include a statistician. Approval of such requests is at Gilead Science's discretion and is dependent on the nature of the request, the merit of the research proposed, the availability of the data, and the intended use of the data. Data requests should be sent to datarequest@gilead.com.

References

1. Tsuda H, Koga M, Nojima M, Senkoji T, Kubota M, Kikuchi T, et al. Changes in survival and causes of death among people living with HIV: Three decades of surveys from Tokyo, one of the Asian metropolitan cities. *J Infect Chemother*. 2021;27(7):949-956. [doi: 10.1016/j.jiac.2021.02.003]
2. Broyles LN, Luo R, Boeras D, Vojnov L. The risk of sexual transmission of HIV in individuals with low-level HIV viraemia: a systematic review. *Lancet*. 2023;402(10400):464-471. [doi: 10.1016/S0140-6736(23)00877-2]
3. Government Public Relations Office Cabinet Office Government of Japan. Public opinion survey on HIV infection and AIDS. 2018. URL: <https://survey.gov-online.go.jp/hutai/h29/h29-hivg.pdf> [accessed
4. Kanamori R, Umemura F, Uemura K, Miyagami T, Valenti S, Fukui N, et al. Web-based search volume for HIV tests and HIV-testing preferences during the COVID-19 pandemic in Japan: infodemiology study. *JMIR Form Res*. 2024;8:e52306. [doi: 10.2196/52306]
5. Golub SA, Gamarel KE. The impact of anticipated HIV stigma on delays in HIV testing behaviors: findings from a community-based sample of men who have sex with men and transgender women in New York City. *AIDS Patient Care STDS*. 2013;27(11):621-627. [doi: 10.1089/apc.2013.0245]
6. Young SD, Bendavid E. The relationship between HIV testing, stigma, and health service usage. *AIDS Care*. 2010;22(3):373-380. [doi: 10.1080/09540120903193666]
7. Babel RA, Wang P, Alessi EJ, Raymond HF, Wei C. Stigma, HIV risk, and access to HIV prevention and treatment services among men who have sex with men (MSM) in the United

States: a scoping review. *AIDS Behav.* 2021;25(11):3574-3604. [doi: 10.1007/s10461-021-03262-4]

8. Norcini-Pala A, Stringer KL, Kempf MC, Konkle-Parker D, Wilson TE, Tien PC, et al. Longitudinal associations between intersectional stigmas, antiretroviral therapy adherence, and viral load among women living with HIV using multidimensional latent transition item response analysis. *Soc Sci Med.* 2024;366:117643. [doi: 10.1016/j.socscimed.2024.117643]

9. Ninnoni JP, Agyemang SO, Bennin L, Agyare E, Gyimah L, Senya K, et al. Coping with loneliness and stigma associated with HIV in a resource-limited setting, making a case for mental health interventions; a sequential mixed methods study. *BMC Psychiatry.* 2023;23(1):163. [doi: 10.1186/s12888-023-04643-w]

10. Turan B, Budhwani H, Fazeli PL, Browning WR, Raper JL, Mugavero MJ, et al. How does stigma affect people living with HIV? The mediating roles of internalized and anticipated HIV stigma in the effects of perceived community stigma on health and psychosocial outcomes. *AIDS Behav.* 2017;21(1):283-291. [doi: 10.1007/s10461-016-1451-5]

11. Andersson GZ, Reinius M, Eriksson LE, Svedhem V, Esfahani FM, Deuba K, et al. Stigma reduction interventions in people living with HIV to improve health-related quality of life. *Lancet HIV.* 2020;7(2):e129-e140. [doi: 10.1016/S2352-3018(19)30343-1]

12. Rueda S, Mitra S, Chen S, Gogolishvili D, Globerman J, Chambers L, et al. Examining the associations between HIV-related stigma and health outcomes in people living with HIV/AIDS: a series of meta-analyses. *BMJ Open.* 2016;6(7):e011453. [doi: 10.1136/bmjopen-2016-011453]

13. Hoang VTH, Pham HT, Nguyen LTP, Tran NA, Le-Thi VQ. The relationship between HIV-related stigma and quality of life among HIV infected outpatients: A cross-sectional study in

- Vietnam. *J Public Health Res.* 2024;13(1):22799036241238667. [doi: 10.1177/22799036241238667]
14. UNAIDS. HIV and stigma and discrimination. 2021. URL: https://www.unaids.org/sites/default/files/media_asset/07-hiv-human-rights-factsheet-stigma-discrimination_en.pdf [accessed November 1, 2024]
15. Yu F, Hsiao YH, Park S, Kambara K, Allan B, Brough G, et al. The influence of anticipated HIV stigma on health-related behaviors, self-rated health, and treatment preferences among people living with HIV in East Asia. *AIDS Behav.* 2023;27(4):1287-1303. [doi: 10.1007/s10461-022-03865-5]
16. Centers for Disease Control. Stigma and HIV. 2024. URL: <https://www.cdc.gov/hiv/health-equity/index.html> [accessed November 4,]
17. Smith RA, Bishop RE. Insights into stigma management communication theory: considering stigmatization as interpersonal influence. *J Appl Commun Res.* 2019;47(5):571-590. [doi: 10.1080/00909882.2019.1675894]
18. Brown R, Sillence E, Coventry L, Branley-Bell D, Murphy-Morgan C, Durrant AC. Health stigma on Twitter: investigating the prevalence and type of stigma communication in tweets about different conditions and disorders. *Frontiers.* 2023. [doi: 10.3389/fcomm.2023.1264373]
19. Toriumi F. Big data on Twitter. *Organization Science.* 2015;48(4):47-59.
20. Lazarus JV, Kakalou C, Palayew A, Karamanidou C, Maramis C, Natsiavas P, et al. A Twitter discourse analysis of negative feelings and stigma related to NAFLD, NASH and obesity. *Liver Int.* 2021;41(10):2295-2307. [doi: 10.1111/liv.14969]
21. Stupinski AM, Alshaabi T, Arnold MV, Adams JL, Minot JR, Price M, et al. Quantifying

changes in the language used around mental health on Twitter over 10 years: observational study. *JMIR Ment Health*. 2022;9(3):e33685. [doi: 10.2196/33685]

22. Ireland ME, Chen Q, Schwartz HA, Ungar LH, Albarracin D. Action tweets linked to reduced county-level HIV prevalence in the United States: online messages and structural determinants. *AIDS Behav*. 2016;20(6):1256-1264. [doi: 10.1007/s10461-015-1252-2]

23. Statista. Leading countries based on number of X (formerly Twitter) users as of April 2024(in millions). 2024. URL: <https://www.statista.com/statistics/242606/number-of-active-twitter-users-in-selected-countries/> [accessed September 11, 2024]

24. Hao T, Huang Z, Liang L, Weng H, Tang B. Health Natural Language Processing: Methodology Development and Applications. *JMIR Med Inform*. 2021;9(10):e23898. [doi: 10.2196/23898]

25. Elbattah M, Arnaud E, Gignon M, G D, editors. The Role of Text Analytics in Healthcare: A Review of Recent Developments and Applications. Proceedings of the 14th International Joint Conference on Biomedical Engineering Systems and Technologies (BIOSTEC 2021); 2021.

26. Adrover C, Bodnar T, Huang Z, Telenti A, Salathe M. Identifying adverse effects of HIV drug treatment and associated sentiments using Twitter. *JMIR Public Health Surveill*. 2015;1(2):e7. [doi: 10.2196/publichealth.4488]

27. Gabarron E, Serrano JA, Wynn R, Lau AY. Tweet content related to sexually transmitted diseases: no joking matter. *J Med Internet Res*. 2014;16(10):e228. [doi: 10.2196/jmir.3259]

28. Li Z, Qiao S, Jiang Y, Li X. Building a social media-based HIV risk behavior index to inform the prediction of HIV new diagnosis: a feasibility study. *AIDS*. 2021;35(Suppl 1):S91-S99. [doi: 10.1097/QAD.0000000000002787]

29. Lu X, Brelsford C. Network structure and community evolution on Twitter: human behavior change in response to the 2011 Japanese earthquake and tsunami. *Sci Rep*. 2014;4:6773. [doi: 10.1038/srep06773]
30. Tsubokura M, Onoue Y, Torii HA, Suda S, Mori K, Nishikawa Y, et al. Twitter use in scientific communication revealed by visualization of information spreading by influencers within half a year after the Fukushima Daiichi nuclear power plant accident. *PLoS One*. 2018;13(9):e0203594. [doi: 10.1371/journal.pone.0203594]
31. Dsouza VS, Rajkhowa P, Mallya BR, Raksha DS, Mrinalini V, Cauvery K, et al. A sentiment and content analysis of tweets on monkeypox stigma among the LGBTQ+ community: a cue to risk communication plan. *Dialogues Health*. 2023;2:100095. [doi: 10.1016/j.dialog.2022.100095]
32. Devlin J. BERT: Pre-training of deep bidirectional transformers for language understanding. *arXiv*. 2018;1810.04805.
33. Ptaszynski E, Masui F. Automatic cyberbullying detection: emerging research and opportunities. Hershey, PA IGI Globla; 2018.
34. Tholke P, Mantilla-Ramos YJ, Abdelhedi H, Maschke C, Dehgan A, Harel Y, et al. Class imbalance should not throw you off balance: Choosing the right classifiers and performance metrics for brain decoding with imbalanced data. *Neuroimage*. 2023;277:120253. [doi: 10.1016/j.neuroimage.2023.120253]
35. Onikoyi B, Nnamoko N, Korkontzelos I. Gender prediction with descriptive textual data using a Machine Learning approach. *Nat Lang Proc*. 2023;4(100018):100018.
36. Hakim MS. SARS-CoV-2, Covid-19, and the debunking of conspiracy theories. *Rev Med Virol*. 2021;31(6):e2222. [doi: 10.1002/rmv.2222]

37. Zhang JM, Wang Y, Mouton M, Zhang J, Shi M. Public Discourse, User Reactions, and Conspiracy Theories on the X Platform About HIV Vaccines: Data Mining and Content Analysis. *J Med Internet Res*. 2024;26:e53375. [doi: 10.2196/53375]
38. Watanabe K TH, Totsu Y. Sources of Sexual Knowledge for High School Students in Tokyo, Japan. *Creative Education*. 2018;9(2394-2404).
39. Associated Press. COVID-19 vaccines don't contain HIV. 2023. URL: <https://apnews.com/article/fact-check-coronavirus-vaccine-hiv-185375755407> [accessed
40. Nachega JB, Morroni C, Zuniga JM, Sherer R, Beyrer C, Solomon S, et al. HIV-related stigma, isolation, discrimination, and serostatus disclosure: a global survey of 2035 HIV-infected adults. *J Int Assoc Physicians AIDS Care (Chic)*. 2012;11(3):172-178. [doi: 10.1177/1545109712436723]
41. Otani M, Shiino T, Hachiya A, Gatanaga H, Watanabe D, Minami R, et al. Association of demographics, HCV co-infection, HIV-1 subtypes and genetic clustering with late HIV diagnosis: a retrospective analysis from the Japanese Drug Resistance HIV-1 Surveillance Network. *J Int AIDS Soc*. 2023;26(5):e26086. [doi: 10.1002/jia2.26086]
42. YS K. Japan Addresses the Global HIV/AIDS Crisis: The Roles of Media and Civil Society in Shaping Perceptions and Aid. *Asian Perspective*. 2015;39:483-511.
43. Togari T, Abe S, Inoue Y. [HIV-related public stigma and knowledge regarding the campaign slogan "undetectable=untransmittable" among Japanese people]. *Nihon Koshu Eisei Zasshi*. 2022;69(2):146-157. [doi: 10.11236/jph.21-005]
44. Valeri L, Amsalem D, Jankowski S, Susser E, Dixon L. Effectiveness of a Video-Based Intervention on Reducing Perceptions of Fear, Loneliness, and Public Stigma Related to COVID-


19: A Randomized Controlled Trial. *Int J Public Health*. 2021;66:1604164. [doi: 10.3389/ijph.2021.1604164]




Supplementary Files

Figures


Quantification of tweets identified. aURL: uniform resource locator.




Receiver operating characteristic curves showing evaluation of (top panel) stigma model and (bottom panel) antistigma model. AUC: area under the curve.



Message classification trends for 2011 through 2023: (top panel) number and percentage of tweets with stigma, neutral, general education, or antistigma labels; (middle panel) number and percentage of tweets with different types of stigma; and (bottom panel) number and percentage of tweets with type of antistigma label.



Characteristics associated with stigma message: (top center) overall; (top of left column) label; (middle of left column) mark; (bottom of left column) responsibility; (top of right column) peril; (middle of right column) fear; and (bottom of right column) insult.



Characteristics associated with (top) general education and (bottom) antistigma.

