

Enhancing Breast Ultrasound Diagnosis with ChatGPT-4: A Large Language Model for BI-RADS Classification and Malignancy Prediction

Su MiaoJiao, Wu SongSong, Liang Xia, Zeng Xian Tao, Hong Zhi Liang, Cheng Sheng

Submitted to: JMIR Medical Informatics
on: January 06, 2025

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 18

..... 18

..... 18

Figures 19

Figure 1..... 20

Figure 2..... 21

Figure 3..... 22

Figure 4..... 23

Enhancing Breast Ultrasound Diagnosis with ChatGPT-4: A Large Language Model for BI-RADS Classification and Malignancy Prediction

Su Miaojiao^{1*} MME; wu songsong^{2*} PhD; Liang Xia^{1*} MME; Zeng Xian Tao¹ MME; Hong Zhi Liang¹ PhD; Cheng Sheng¹ MME

¹ Department of Ultrasound, Fujian Provincial Hospital Fujian Provincial Hospital Fuzhou CN

² Fujian Provincial Hospital Fuzhou CN

*these authors contributed equally

Corresponding Author:

wu songsong PhD

Fujian Provincial Hospital

No. 134, Dong Street, Gulou District, Fuzhou City, Fujian Province, China

Fuzhou

CN

Abstract

Objective: To explore the feasibility of using publicly available mainstream large language models (LLMs) to evaluate the consistency and diagnostic accuracy of standardized ultrasound imaging reports, using pathology as the reference standard. Materials and

Methods: This retrospective study collected ultrasound imaging data of breast nodules with pathological diagnoses obtained at our hospital from June 2019 to June 2024. ChatGPT-4 was employed to evaluate the BI-RADS classification and benign or malignant nature of the nodules. The diagnostic performance of the LLM and the human-machine dialogue method (image-to-text-LLM) was assessed, including accuracy, sensitivity, specificity, and the area under the receiver operating characteristic curve (AUC).

Results: A total of 671 patients (mean age 45.82 ± 9.20 years [SD]; age range 26–75 years) with 671 breast nodule ultrasound images (385 benign, 286 malignant) were included. ChatGPT-4 achieved an overall accuracy of 96.87% in identifying BI-RADS classifications, surpassing the performance of two junior radiologists. For image interpretation, ChatGPT-4 achieved an AUC of 0.82 (95% CI: 0.79–0.85), an accuracy of 80.63% (541 of 671 cases), a sensitivity of 90.56% (259 of 286 cases), and a specificity of 73.51% (283 of 385 cases). Its diagnostic performance was comparable to that of two senior radiologists and superior to two junior radiologists. When utilizing the image-to-text-LLM, diagnostic performance, including AUC, accuracy, sensitivity, and specificity, improved for all four radiologists.

Conclusion: The application of LLMs (particularly the image-to-text-LLM) demonstrates potential value in diagnosing breast ultrasound imaging data.

(JMIR Preprints 06/01/2025:70924)

DOI: <https://doi.org/10.2196/preprints.70924>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in a peer-reviewed journal, I will be able to make the full manuscript available to all users.

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in <https://preprints.jmir.org/preprint/70924>



Original Manuscript

Enhancing Breast Ultrasound Diagnosis with ChatGPT-4: A Large Language Model for BI-RADS Classification and Malignancy Prediction

Abstract

Background Breast ultrasound is essential for evaluating breast nodules, with BI-RADS providing standardized classification. However, interobserver variability among radiologists can affect diagnostic accuracy. Large language models (LLMs) like ChatGPT-4 have shown potential in medical imaging interpretation. This study explores its feasibility in improving BI-RADS classification consistency and malignancy prediction compared to radiologists.

Objective: This study aims to evaluate the feasibility of using LLMs, particularly ChatGPT-4, to assess the consistency and diagnostic accuracy of standardized breast ultrasound imaging reports, using pathology as the reference standard.

Materials and Methods: This retrospective study analyzed breast nodule ultrasound data from 671 female patients (mean age 45.82 ± 9.20 years; range 26–75 years) who underwent biopsy or surgical excision at our hospital between June 2019 and June 2024. ChatGPT-4 was used to interpret BI-RADS classifications and predict benign versus malignant nodules. The study compared the model's performance to that of two senior radiologists (≥ 15 years of experience) and two junior radiologists (< 5 years of experience) using key diagnostic metrics, including accuracy, sensitivity, specificity, area under the receiver operating characteristic curve (AUC), P-values, and odds ratios (OR) with 95% confidence intervals (CI). Two diagnostic models were evaluated: (1) Image interpretation model, where ChatGPT-4 classified nodules based on BI-RADS features; (2) Image-to-text-LLM model, where radiologists provided textual descriptions, and ChatGPT-4 determined malignancy probability based on keywords. Radiologists were blinded to pathological outcomes, and BI-RADS classifications were finalized through consensus.

Results: ChatGPT-4 achieved an overall BI-RADS classification accuracy of 96.87%, outperforming junior radiologists (91.95% and 90.01%, $P < 0.01$). For malignancy prediction, ChatGPT-4 achieved an AUC of 0.82 (95% CI: 0.79–0.85), an accuracy of 80.63% (541/671 cases), a sensitivity of 90.56% (259/286 cases), and a specificity of 73.51% (283/385 cases). The image interpretation model demonstrated performance comparable to senior radiologists, while the image-to-text-LLM model further improved diagnostic accuracy for all radiologists, increasing their sensitivity and specificity significantly ($P < 0.001$). Statistical analyses, including McNemar's test and DeLong's test, confirmed that ChatGPT-4 outperformed junior radiologists ($P < 0.01$) and showed non-inferiority compared to senior radiologists ($P > 0.05$). Pathological diagnoses served as the reference standard, ensuring robust evaluation reliability.

Conclusion: Integrating ChatGPT-4 into an image-to-text-LLM workflow improves BI-RADS classification accuracy and supports radiologists in breast ultrasound diagnostics. These results demonstrate its potential as a decision-support tool to enhance diagnostic consistency and reduce variability.

Keywords: Large Language Models, BI-RADS, Breast Ultrasound, ChatGPT-4, Image-to-Text-LLM

Introduction

In recent years, large language models (LLMs) have advanced rapidly, demonstrating significant potential in healthcare. Their diverse applications have garnered considerable attention[1-4], driving revolutionary progress across various fields, including medicine[5]. LLMs are built upon large-scale datasets and continuously refined through iterative training, enabling them to handle a wide range of

tasks, including medical diagnosis. LLMs provide diagnostic recommendations by processing key inputs, extracting relevant information, and performing associative reasoning[6]. In medical imaging analysis, the integration of LLMs with image-text conversion models enables the creation of a continuous interactive system that seamlessly integrates with clinical practice, delivering more precise and personalized healthcare services[7]. This approach enhances medical technology, broadens clinical applications, and partially alleviates the challenges posed by limited healthcare resources.

GPT-4, developed by OpenAI, is one of the most widely recognized LLMs. Although its applications in medical imaging analysis have shown promise, the specific role and diagnostic accuracy of LLMs in this domain remain insufficiently explored. Initial research on LLMs primarily focused on evaluating their language generation capabilities. Most discussions surrounding LLMs, particularly ChatGPT, have revolved around comments, replies, or question-and-answer formats[8-11]. The findings have been promising[12]. However, many studies have primarily focused on response evaluation rather than quantitative assessments of diagnostic accuracy and have lacked reference standards such as pathology or laboratory examinations[13]. Additionally, the ability of LLMs to generate structured diagnostic reports, such as BI-RADS classifications, has not been systematically evaluated, raising concerns about their reliability in clinical decision-making. Furthermore, LLMs may inadvertently perpetuate existing biases and disparities.

Therefore, to address this gap, it is essential to demonstrate the application of LLMs in medical imaging. Breast cancer serves as an optimal case study for this purpose due to its high incidence, critical need for early detection, and reliance on imaging for diagnosis. Breast cancer is one of the most life-threatening diseases, second only to lung cancer[14,15]. The likelihood of breast cancer survival is largely determined by the stage at which it is detected, which significantly impacts chemotherapy outcomes[16-18]. Early diagnosis and effective treatment of breast cancer can significantly reduce mortality rates[19]. Unlike lung cancer, which primarily relies on computed tomography (CT) imaging, or neurological disorders, which depend on magnetic resonance imaging (MRI) and functional imaging, breast cancer diagnosis heavily incorporates ultrasound due to its accessibility, cost-effectiveness, and suitability for dense breast tissue—particularly in younger patients and in regions with high breast cancer incidence, such as China[20]. Therefore, in China, ultrasound has become the preferred imaging modality for breast lesion screening and preoperative evaluation[21]. Breast ultrasound not only facilitates routine detection of breast cancer but also aids in resolving inconclusive mammographic findings[22-24]. The Breast Imaging Reporting and Data System (BI-RADS), proposed by the American College of Radiology (ACR), is a standardized system designed to describe breast imaging findings[25]. It assists radiologists and breast surgeons in standardizing the assessment of malignancy probability in breast lesions and in guiding subsequent diagnostic and treatment plans. Physicians classify breast lesions based on specific characteristics, including shape, orientation, margins, echogenic pattern, posterior features, and the presence of calcifications and associated findings. However, unlike structured reporting systems in CT-based lung cancer screening or MRI-based neurological assessments, BI-RADS assessment in breast ultrasound remains highly dependent on operator experience. This reliance leads to interobserver variability and diagnostic inconsistency, as radiologists with different levels of expertise may provide varying interpretations. Additionally, ultrasound evaluation requires extensive training and certification, making it resource-intensive and subject to human variability.

Given these challenges, breast ultrasound serves as an ideal testbed for evaluating LLM-assisted imaging interpretation. First, the structured nature of BI-RADS makes it well-suited for AI-driven standardization, potentially reducing subjective variability. Second, the widespread clinical use of ultrasound in breast cancer detection allows for large-scale validation of LLM-generated reports against pathology results. Unlike lung CT or brain MRI, where AI-based segmentation and feature extraction already play a significant role, breast ultrasound interpretation lacks a robust AI-driven standardization framework, highlighting the potential value of LLM integration. LLMs offer a

promising solution by providing standardized interpretations, reducing interobserver variability, and assisting in structured reporting. Their ability to generate BI-RADS reports based on ultrasound findings suggests a potential role in improving diagnostic consistency and streamlining clinical workflows. Despite these advantages, the reliability and diagnostic accuracy of LLM-generated BI-RADS assessments remain unclear.

To bridge this research gap, this study aims to systematically evaluate the diagnostic performance, consistency, and clinical utility of mainstream LLMs in breast nodule assessment. By comparing LLM-generated BI-RADS categorizations with pathology-confirmed diagnoses, we seek to determine whether LLMs can enhance reproducibility, improve standardization, and assist radiologists in breast cancer diagnosis.

Materials and Methods

This study is a retrospective study, and informed consent was waived due to its retrospective design. No patient privacy information was provided to the LLMs.

Study Design

To compare the diagnostic performance of the LLMs and radiologists with different levels of experience in interpreting breast ultrasound images.

First, among the widely used LLMs, ChatGPT and Gemini are prominent. However, since Gemini does not support image analysis, we selected OpenAI's ChatGPT-4 (San Francisco, CA, USA) for this study[26], utilizing the BI-RADS standard for diagnosing breast nodules. For clarity, while LLMs refer to a broad category of AI models, this study specifically focuses on ChatGPT-4. We analyzed the accuracy of ChatGPT-4 in classifying breast nodules using the BI-RADS system and assessed its diagnostic performance in distinguishing between benign and malignant breast nodules. In subsequent analyses, the performance of two different models was evaluated:

(a) **Image-to-text-LLM model:** Radiologists first interpreted the ultrasound images and generated textual descriptions using a standardized BI-RADS reporting template. This template included predefined descriptors for nodule shape, margin, echogenicity, and other key ultrasound features. ChatGPT-4 then analyzed the text and determined whether the nodule was benign or malignant based on the extracted keywords (Figure 3). To minimize variability in descriptions, all radiologists were trained in the standardized reporting format, and their reports were randomly reviewed by a senior radiologist to ensure consistency.

(b) **Image interpretation model:** The ultrasound images were provided directly, and the model performed benign or malignant diagnosis based on BI-RADS classification and image analysis (Figure 4).

All radiologists were blinded to the pathological results. The radiologists were divided into two senior and two junior groups. The senior radiologists each had over 15 years of experience in breast ultrasound, while the junior radiologists had less than 5 years of experience.

The BI-RADS classifications were determined by consensus between two senior radiologists. If their conclusions differed, a third senior radiologists (W.S.S., with 21 years of experience in breast ultrasound) was consulted to reach a final decision through discussion.

Patient Data Collection

Retrospective collection of ultrasound images of breast nodules diagnosed through biopsy or surgical excision at Fujian Provincial Hospital, Fuzhou University Affiliated Hospital, between June 2019 and June 2024 (385 benign and 286 malignant cases). Patients without ultrasound examinations, those with low-quality images, or those with inconclusive pathological results were excluded (Figure 1). Each image corresponded to an individual breast nodule. For nodules suspected to be benign on biopsy, follow-up of at least six months was required. All lesions were ultimately classified as benign or malignant based on histopathological results.

Evaluation index

The diagnostic performance of ChatGPT-4 in two different models (Image-to-text-LLM model and image interpretation model) was compared with that of radiologists with varying levels of

experience. The analysis included the accuracy of BI-RADS classifications by junior radiologists and the calculation of key performance metrics, including sensitivity, specificity, accuracy, and the area under the receiver operating characteristic curve (AUC) with 95% confidence intervals (Figure 2).

Statistical Analysis

All statistical analyses were performed using SPSS software (IBM, version 20.0) and MedCalc (MedCalc Software, version 12.7). Continuous data were expressed as mean $\pm$ standard deviation (SD), and categorical data were presented as frequencies and percentages. The analysis included the calculation of key performance metrics, including accuracy, sensitivity, specificity, and the area under the receiver operating characteristic curve (AUC) with 95% confidence intervals. Finally, McNemar's test and DeLong's test were used to compare the diagnostic performance of ChatGPT-4 under different deployment strategies. A p -value of < 0.05 was considered statistically significant.

Results

Patient Characteristics:

A total of 671 female patients were included in the study (mean age 45.82 ± 9.20 years [SD]; age range 26–75 years). A total of 671 ultrasound images of breast nodules were collected, including 385 benign cases (57.38%) and 286 malignant cases (42.62%), with 428 nodules on the left side and 243 on the right side (Table 1). Pathological diagnoses within three months after ultrasound examination were based on surgical resection or core needle biopsy. Patients with benign pathology from core needle biopsy were followed up for an average of approximately 11.2 months. The inclusion and exclusion criteria for patients and images are detailed in Figure 1.

The representation of ChatGPT-4 in classification

As shown in Table 2, the BI-RADS classifications for the 671 ultrasound images include categories BI-RADS 3, 4 (A-C), and 5. The overall accuracy of the ChatGPT-4 in identifying BI-RADS classifications was 96.87%, surpassing junior radiologist 1 (91.95%) and junior radiologist 2 (90.01%). Notably, the model demonstrated significant differences in classification performance compared to both junior radiologists for BI-RADS 4A and to junior radiologist 2 for BI-RADS 3 and 4C.

The performance of image interpretation model in predicting benign and malignant breast nodules:

As shown in Table 3, Senior radiologist 1 demonstrated excellent performance in diagnosing benign and malignant breast lesions via ultrasound, with an AUC of 0.85 (95% CI: 0.81–0.88), an accuracy of 83.16% (558/671 cases), a sensitivity of 91.61% (262/286 cases), and a specificity of 77.40% (298/385 cases). ChatGPT-4 exhibited comparable performance to the image interpretation model of both senior radiologists in terms of accuracy, sensitivity, and specificity. Notably, it outperformed senior radiologist 2 in AUC, with the difference being statistically significant ($P < 0.01$). The image interpretation model showed diagnostic performance comparable to that of the two senior radiologists, particularly achieving a sensitivity of 90.56% (259/286 cases). Moreover, it consistently outperformed the predictive models of the junior radiologists in AUC, accuracy, and sensitivity ($P < 0.01$), while showing a statistically significant difference in specificity compared to junior radiologist 1 ($P < 0.01$).

The performance of ChatGPT-4 in predicting benign and malignant breast nodules based on image-to-text-LLM model

As shown in Table 4, the image interpretation model of ChatGPT-4 yielded an AUC of 0.82 (95% CI: 0.79–0.85), an accuracy of 80.63% (541/671 cases), a sensitivity of 90.56% (259/286 cases), and a specificity of 73.51% (283/385 cases). Under the Image-to-text-LLM model, ChatGPT-4 demonstrated performance comparable to the two senior radiologists while surpassing the models of junior radiologists in AUC, accuracy, sensitivity, and specificity ($P < 0.001$). The four radiologists with varying levels of experience showed improvements in AUC, accuracy, sensitivity, and

specificity after image-to-text–LLM model compared to image-to-text–LLM model.

Table 1: Demographic and Clinical Characteristics of Patients

Characteristic	Value
Age (y)*	45.82±9.20□26-75□
Location	
Left	428□63.79□
Right	243□36.21□
Nodule Pathology	
Benign	385□57.38□
Malignant	286□42.62□

Note: Unless otherwise specified, the data represents the number of survey results, with percentages in parentheses. *Data are presented as mean ± standard deviation, with the range in parentheses.

Table 2: The effectiveness of ChatGPT-4 in classification

Classification	N	ChatGPT-4	Junior Physicians 1	P	Junior Physicians2	P
BI-RADS 3	93	90	86	0.29	82	0.04
BI-RADS 4A	374	361	345	0.02	341	<0.01
BI-RADS 4B	98	95	91	0.22	90	0.13
BI-RADS 4C	56	55	50	0.06	47	<0.01
BI-RADS 5	50	49	45	0.13	44	0.06

Note: The *p*-value represents the comparison with ChatGPT-4

Table 3: The performance of image interpretation model in predicting benign and malignant breast nodules

Category	AUC*	P	Accuracy (%)	P	Sensitivity (%)	P	Specificity (%)	P
ChatGPT-4	0.82□0.79□0.85□		80.63□541/671□		90.56(259/286)		73.51(283/385)	
Senior Radiologist 1	0.81□0.77,0.84□	0.10	79.88□536/671□	0.88	86.01(246/286)	0.18	75.32(290/385)	0.50
Senior Radiologist 2	0.79□0.76,0.83□	<0.01	78.09□524/671□	0.24	86.71(248/286)	0.23	71.69(276/385)	0.63
Junior Radiologist 1	0.66□0.62,0.70□	<0.01	65.57□440/671□	<0.01	63.33(181/286)	<0.01	67.27(259/385)	<0.01
Junior Radiologist 2	0.67□0.62,0.71□	<0.01	66.47□446/671□	<0.01	66.78(191/286)	<0.01	66.23(255/385)	0.13

Note.—Except were indicated, numbers in parentheses are numbers of nodules. AUC = area under a receiver operating characteristic curve.
* Data in parentheses for the AUC metric are 95% CIs.

Table 4: The performance of ChatGPT-4 in predicting benign and malignant breast nodules based on image-to-text–LLM model

Diagnostic Performance	ChatGPT-4	Senior Radiologist 1	Senior Radiologist 2	Junior Radiologist 1	Junior Radiologist 2
AUC*	0.82□0.79□0.85□	0.85□0.81,0.88□	0.82□0.78,0.85□	0.64□0.62,0.70□	0.63□0.59,0.67□
P		0.10	0.77	<0.01	<0.01

Accuracy (%)	80.63[541/671]	83.16[558/671]	81.37[546/671]	72.43[486/671]	69.00[463/671]
		0.13	0.72	<0.01	<0.01
Specificity (%)	90.56(259/286)	91.61(262/286)	86.71(248/286)	83.92(240/286)	74.13(212/286)
		0.87	0.29	0.02	<0.01
Sensitivity (%)	73.51(283/385)	77.40(298/385)	76.36(294/385)	64.42(248/385)	65.71(253/385)
		0.10	0.18	<0.01	0.01

Note.—Except were indicated, numbers in parentheses are numbers of nodules. AUC = area under a receiver operating characteristic curve, LLM = large language model.
* Data in parentheses for the AUC metric are 95% CIs.

Figure 1: Flow chart




Figure 2: Study Design




Figure 3: The screenshot demonstrates ChatGPT-4 (OpenAI; <https://chat.openai.com/>) analyzing a single breast nodule in image-to-text-LLM model. This response was recorded as a malignant diagnosis.





Figure 4: The screenshot demonstrates ChatGPT-4 (OpenAI; <https://chat.openai.com/>) analyzing a single breast nodule in image interpretation model. This response was recorded as a malignant diagnosis.

Do you know the ACR BI-RADS of breast ultrasound

Yes, I'm familiar with the ACR BI-RADS for breast ultrasound. The **ACR BI-RADS** (Breast Imaging Reporting and Data System) was developed by the American College of Radiology to standardize breast imaging reporting, improve diagnostic accuracy, and ensure clear communication among healthcare providers. This system is used across various breast imaging modalities, including ultrasound, mammography, and MRI, and provides a framework for assessing the likelihood of malignancy and guiding patient management. Here's a summary of the BI-RADS categories:

1. **BI-RADS 0:** Incomplete - Additional imaging evaluation is needed.
2. **BI-RADS 1:** Negative - No significant findings.
3. **BI-RADS 2:** Benign findings - Definitely benign.
4. **BI-RADS 3:** Probably benign - Follow-up is recommended, as the finding has a low likelihood of malignancy.
5. **BI-RADS 4:** Suspicious abnormality - Biopsy should be considered; this is subdivided into:
 - **4A:** Low suspicion for malignancy
 - **4B:** Moderate suspicion for malignancy
 - **4C:** High suspicion for malignancy
6. **BI-RADS 5:** Highly suggestive of malignancy - Biopsy or appropriate intervention is strongly recommended.



This is an image of a breast ultrasound, whether you think it's benign or malignant

In examining this ultrasound image with color Doppler flow, several features suggest a high suspicion for malignancy:

1. **Increased Internal Vascularity:** The nodule exhibits significant internal blood flow, with both red and blue signals, indicating active blood supply within the lesion. Increased vascularity is commonly associated with malignant tumors, as they often induce angiogenesis to support rapid growth.
2. **Echotexture and Margins:** Although the margins are not completely clear in this view, the internal structure appears heterogeneous, which is often a concerning feature in breast imaging. Malignant nodules tend to have irregular or heterogeneous echotextures, as opposed to the homogeneous texture typically seen in benign lesions.

Based on these observations, this lesion would likely be classified as **BI-RADS category 4C or 5**, indicating a high suspicion of malignancy. A biopsy is strongly recommended to confirm the diagnosis.

Discussion

In medical diagnostics, logical reasoning often forms the foundation of treatment, requiring higher-order cognitive skills such as application, analysis, and evaluation. Bhayana et al.[27] demonstrated that ChatGPT nearly passed a radiology board-style exam despite lacking radiology-specific pretraining or direct access to imaging data. Despite the promising potential of LLMs in medical imaging[28], there is a lack of research on their feasibility in addressing reasoning tasks associated with medical diagnoses based on breast-specific reference standards, such as pathology. In this study, we utilized BI-RADS and standardized pathological reporting as reference standards, integrating LLMs into the diagnostic process to emphasize the benefits of combining ChatGPT-4 with image-to-text technologies. We validated the potential application of ChatGPT-4 in breast nodule ultrasound diagnostics, using pathological results as the benchmark to evaluate their performance in standardized reporting and diagnostic consistency. The results showed that ChatGPT-4 performed exceptionally well in BI-RADS classification and benign-versus-malignant predictions of breast nodules. Its diagnostic performance was significantly superior to that of less experienced radiologists. This lays a foundation for the practical application of LLMs in enhancing the standardization and accuracy of medical imaging diagnostics while providing evidence to support further exploration of their clinical utility.

ChatGPT-4 achieved an overall accuracy of 96.87% in BI-RADS classification and an AUC of 0.82 for benign-versus-malignant predictions (AUC = 0.82 [95% CI: 0.79–0.85]). Its accuracy, sensitivity, and specificity were 80.63% (541/671), 90.56% (259/286), and 73.51% (283/385), respectively. Notably, ChatGPT-4 outperformed less experienced radiologists in BI-RADS categories 4A, 3, and 4C. This indicates that ChatGPT-4 can deliver highly consistent and accurate diagnostic results, reducing variability caused by limited experience. Integrating LLMs into breast condition classification can minimize diagnostic errors and ultimately improve patient outcomes significantly. LLMs can be trained to identify and interpret individual variations, ultimately assisting in the development of new classification standards. This approach may address the lack of universally accepted breast disease classifications. Furthermore, integrating image-to-text conversion by radiologists with ChatGPT-4 improved the AUC, accuracy, sensitivity, and specificity for all four physicians, with notable performance enhancements observed among less experienced radiologists. This underscores that while AI has made significant advancements in medical imaging and diagnostics, human expertise remains indispensable.

While ChatGPT-4 demonstrated strong diagnostic capabilities, it is important to compare its performance with existing AI-driven ultrasound diagnostic models. Prior deep-learning models have shown comparable or even superior performance in specific ultrasound tasks. For example, convolutional neural networks (CNNs) and transformer-based models trained on large annotated datasets have achieved AUCs exceeding 0.90 in distinguishing benign from malignant breast lesions[29,30]. These models directly analyze imaging data, extracting pixel-level features for classification. In contrast, ChatGPT-4 processes structured text inputs rather than images, highlighting a key trade-off between interpretability and diagnostic precision. While CNNs and other deep-learning models operate as "black-box" systems with high accuracy, LLMs like ChatGPT-4 provide enhanced transparency and reasoning, allowing clinicians to trace the logic behind their decisions. Thus, future studies should explore hybrid approaches that integrate deep-learning image analysis with LLM-driven reasoning to maximize diagnostic performance.

Our study, grounded on pathological findings as the reference standard, provides substantial evidence that using pathology-based benchmarks to evaluate LLMs in ultrasound diagnostic reasoning and information processing differs significantly from previous approaches. Earlier studies primarily relied on expert questionnaires and subjective judgments to assess LLMs' text comprehension, summarization, and generation abilities in the context of medical imaging reports or inquiry responses[31,32]. The results showed that ChatGPT-4, when integrated with clinician-led image-to-text conversion strategies, achieved optimal performance. This demonstrates that LLMs can effectively assist radiologist, particularly less experienced radiologists, thereby improving overall diagnostic quality. Compared to senior radiologists, ChatGPT-4 demonstrated sensitivity (90.56%) in benign-versus-malignant diagnosis of breast nodules that closely matched that of two senior radiologists (91.61% and 86.71%). However, its specificity was slightly lower (73.51% vs. 77.40% and 76.36%). These findings suggest that ChatGPT-4's diagnostic reasoning, when provided with structured textual inputs, approaches that of senior radiologists. However, its performance remains constrained by its reliance on human-generated descriptions, rather than direct image interpretation.

Unlike black-box deep-learning models, ChatGPT-4 offers a transparent reasoning process, enhancing trust in AI-assisted diagnostics. This feature is particularly valuable in clinical practice and medical education, as it enables junior radiologists to follow structured diagnostic logic rather than relying on uninterpretable AI-generated outputs. By standardizing ultrasound diagnostic reasoning, ChatGPT-4 could improve the consistency and quality of radiological assessments, especially among less experienced practitioners.

This study demonstrates the advantages of LLMs in breast ultrasound diagnosis. First, LLMs standardize the diagnostic process, significantly reducing inconsistencies caused by variations in physician experience, making them particularly beneficial for less experienced radiologists. Second, they offer high efficiency in processing large image datasets, alleviating the workload of radiologists, particularly in breast cancer screening programs. Furthermore, LLMs can be incorporated into standardized reporting frameworks such as BI-RADS, providing real-time decision support to optimize clinical workflows. Rather than replacing CNN-based models, ChatGPT-4 functions as a complementary tool that interprets structured radiological descriptions, offering an alternative diagnostic strategy. Most traditional diagnostic support software focuses on extracting imaging features rather than providing comprehensive diagnostic suggestions. Our study paves the way for applying LLMs in medical image-based diagnostic reasoning. Future research should explore multimodal AI models that integrate ultrasound, CT, and MRI data to further enhance diagnostic accuracy. Additionally, refining image-to-text conversion methods and reducing reliance on manual input will be critical for advancing fully automated diagnostic systems. Validating these findings on multicenter datasets is essential to ensure generalizability.

Limitations

Despite demonstrating the potential of LLMs in ultrasound diagnostics, this study has certain limitations. First, it did not utilize the latest LLM versions, and ChatGPT-4 lacks the capability to directly interpret medical images. Its diagnostic process depends on structured text inputs provided by physicians, limiting its feasibility for fully automated diagnostics. Therefore, its current role is best suited as a decision-support tool rather than an independent diagnostic system. Second, the study was based on BI-RADS standards and did not account for complex imaging features or multimodal data, potentially restricting the model's applicability to challenging cases. Third, the dataset was derived from a single institution, introducing potential selection bias. Additionally, the patient cohort was drawn from a specific geographic and demographic background, which may not fully represent broader populations. Future studies should include diverse, multiethnic patient groups to improve external validity. Finally, while pathological results were used as the gold standard, they are not always available in routine clinical practice. Investigating alternative validation methods, such as long-term follow-up outcomes, could further enhance the clinical applicability of LLM-assisted diagnostics.

Conclusion

This study systematically evaluates the application of LLMs in breast ultrasound diagnosis. The results show that ChatGPT-4 achieves diagnostic performance comparable to senior radiologists while significantly improving the accuracy of less experienced radiologists. These findings highlight its potential to enhance diagnostic consistency and standardization. As AI and imaging technologies advance, LLMs are expected to further improve diagnostic accuracy and efficiency. Future research should focus on multicenter validation and multimodal imaging integration to further assess its clinical applicability.

References

- [1] Patel SB, Lam K. ChatGPT: the future of discharge summaries? *Lancet Digit Health* 2023;5(3):e107–e108.
- [2] Mukherjee P, Hou B, Lanfredi RB, Summers RM. Feasibility of using the privacy-preserving large language model Vicuna for labeling radiology reports. *Radiology* 2023;309(1):e231147.
- [3] Rau A, Rau S, Zoeller D, et al. A context-based chatbot surpasses radiologists and generic ChatGPT in following the ACR appropriateness guidelines. *Radiology* 2023;308(1):e230970.
- [4] Moy L. Guidelines for use of large language models by authors, reviewers, and editors: considerations for imaging journals. *Radiology* 2023;309(1):e239024.
- [5] Will ChatGPT transform healthcare? *Nat Med* 2023;29(3):505–506.
- [6] Gertz RJ, Bunck AC, Lennartz S, et al. GPT-4 for automated determination of radiological study and protocol based on radiology request forms: a feasibility study. *Radiology* 2023;307(5):e230877.
- [7] Moon JH, Lee H, Shin W, Kim YH, Choi E. Multi-modal understanding and generation for medical images and text via vision-language pretraining. *IEEE J Biomed Health Inform* 2022;26(12):6070–6080.
- [8] Bhayana R, Krishna S, Bleakney RR. Performance of ChatGPT on a radiology board-style examination: insights into current strengths and limitations. *Radiology* 2023;307(5):e230582.
- [9] Ayers JW, Poliak A, Dredze M, et al. Comparing physician and artificial intelligence chatbot responses to patient questions posted to a public social media forum. *JAMA Intern Med* 2023;183(6):589–596.
- [10] Mihalache A, Popovic MM, Muni RH. Performance of an artificial intelligence chatbot in ophthalmic knowledge assessment. *JAMA Ophthalmol* 2023;141(6):589–597.
- [11] Rahsepar AA, Tavakoli N, Kim GHJ, Hassani C, Abtin F, Bedayat A. How AI responds to common lung cancer questions: ChatGPT vs Google Bard. *Radiology* 2023;307(5):e230922.
- [12] Kulkarni PA, Singh H. Artificial intelligence in clinical diagnosis: opportunities, challenges, and hype. *JAMA* 2023;330(4):317–318.

- [13] Fink MA, Bischoff A, Fink CA, et al. Potential of ChatGPT and GPT-4 for data mining of free-text CT reports on lung cancer. *Radiology* 2023;308(3):e231362.
- [14] Siegel RL, Miller KD, Fuchs HE (2021) Jemal A (2021) Cancer statistics. *CA Cancer J Clin* 71(1):7–33.
- [15] Hassani H, Avazzadeh Z, Agarwal P, Mehrabi S, Ebadi MJ, Dahaghin MS, Naraghirad E (2023) A study on fractional tumor-immune interaction model related to lung cancer via generalized laguerre polynomials. *BMC Med Res Methodol* 23(1):189.
- [16] Gorgzadeh A, Hheidari A, Ghanbarikondori P, Arastonejad M, Goki T, Aria M, Allahyartorkaman A, Moazzam F (2023) Investigating the properties and cytotoxicity of cisplatin-loaded nano polybutylcyanoacrylate on breast cancer cells. *Asian Pac J Cancer Biol* 8:345–350.
- [17] Yousefi O, Azami P, Sabahi M, Dabecco R, Adada B, Borghei-Razavi H (2022) Management of optic pathway glioma: A systematic review and meta-analysis. *Cancers* 14:4781.
- [18] Mohammadinezhad F, Talebi A, Allahyartorkaman M, Nahavandi R, Vesal M, Akbarzadeh A (2023) Preparation, characterization and cytotoxic studies of cisplatin-containing nanoliposomes on breast cancer cell lines. *Asian Pac J Cancer Biol* 8:155–159.
- [19] Duggan, C. *et al.* The breast health global initiative 2018 global summit on improving breast healthcare through resource-stratified phased implementation: Methods and overview. *Cancer* 126, 2339–2352 (2020).
- [20] He J, Chen W, Li N, et al. China guideline for the screening and early detection of female breast cancer (2021, Beijing). *Zhonghua Zhong Liu Za Zhi* 2021; 43(4):357–382.
- [21] Bevers TB, Helvie M, Bonaccio E, et al. Breast cancer screening and diagnosis, version 3.2018, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw*. 2018;16(11):1362-1389.
- [22] Rebolj M, Assi V, Brentnall A, Parmar D, Duffy SW. Addition of ultrasound to mammography in the case of dense breast tissue: systematic review and meta-analysis. *Br J Cancer*. 2018;118(12):1559-1570.
- [23] 03. Teichgraber DC, Guirguis MS, Whitman GJ. Breast cancer staging: updates in the *AJCC Cancer Staging Manual*, 8th Edition, and current challenges for radiologists, from the *AJR* special series on cancer staging. *AJR Am J Roentgenol*. 2021;217(2):278-290.
- [24] 04. Sprague BL, Ichikawa L, Eavey J, et al. Breast cancer risk characteristics of women undergoing whole-breast ultrasound screening versus mammography alone. *Cancer*. 2023;129(16):2456-2468.
- [25] L. Levy, M. Suissa, J.F. Chiche, G. Teman, B. Martin, BIRADS ultrasonography, *Eur. J. Radiol*. 61 (2) (2007) 202–211,
- [26] ChatGPT 4.0. OpenAI. <https://chat.openai.com>. Accessed June 18– 28, 2023.
- [27] Bhayana R, Krishna S, Bleakney RR. Performance of ChatGPT on a radiology board-style examination: insights into current strengths and limitations. *Radiology* 2023;307(5):e230582.
- [28] Wu, Shao-Hong., Tong, Wen-Juan., Li, Ming-De., Hu, Hang-Tong., Lu, Xiao-Zhou. , et al. Collaborative Enhancement of Consistency and Accuracy in US Diagnosis of Thyroid Nodules Using Large Language Models. *Radiology* 2024; 310(3):e232255.
- [29] Ji H, Zhu Q, Ma T, et al. Development and validation of a transformer-based CAD model for improving the consistency of BI-RADS category 3–5 nodule classification among radiologists: a multiple center study. *Quant Imaging Med Surg*. 2023;13(6):3907-3920. doi:10.21037/qims-22-1091.
- [30] Zheng X, Yao Z, Huang Y, et al. Deep learning radiomics can predict axillary lymph node status in early-stage breast cancer. *Nat Commun*. 2020;11(1):1236.
- [31] Tariq R, Malik S, Khanna S. Evolving landscape of large language models: an evaluation of ChatGPT and Bard in answering patient queries on colonoscopy. *Gastroenterology* 2024;166(1):220–221.

- [32] Lim ZW, Pushpanathan K, Yew SME, et al. Benchmarking large language models' performances for myopia care: a comparative analysis of ChatGPT-3.5, ChatGPT-4.0, and Google Bard. *EBioMedicine* 2023;95:104770.

Abbreviations

LLM□large language model

BI-RADS□Breast Imaging Reporting and Data System

ACR□American College of Radiology

AUC□area under the curve

CI□confidence intervals

OR□odds ratios

SD□standard deviation

CT□computed tomography

MRI□magnetic resonance imaging

CNN□convolutional neural network

Ethical Statement

This study was approved by the Institutional Review Board of Fujian Provincial Hospital. Informed consent was waived due to the retrospective design. No identifiable patient information was shared with the LLMs.

Funding

The author(s) declare that financial support was received for the research, authorship, and/or publication of this article. Fujian Science and Technology Innovation Joint Fund project climbing project (2023Y9332).

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Authors' Contributions

Su MiaoJiao and Liang Xia contributed equally to this work and are designated as co-first authors. Zeng Xiantao, Hong Zhiliang, and Cheng Sheng contributed as second, third, and fourth authors, respectively, by providing significant support in the study design, data collection, and analysis. Wu Songsong served as the corresponding author, overseeing the entire project, providing critical intellectual input, and ensuring the manuscript's quality and integrity. All authors have read and approved the final manuscript.

Acknowledgments

The authors are thankful to Fujian Provincial Hospital and Fujian Medical University for their management of our patient database.

The authors are thankful to Song-Song Wu for helping critically revise the manuscript for important intellectual content and helping collect data and design the study.

Supplementary Files

Untitled.


URL: <http://asset.jmir.pub/assets/bb737b2a4319bd79ef0afd773d8d0961.docx>

Untitled.


URL: <http://asset.jmir.pub/assets/755007441ffe729f9cf8f89adf6f0a89.docx>

Figures

Flow chart.



Study Design.



The screenshot demonstrates ChatGPT-4 (OpenAI; <https://chat.openai.com/>) analyzing a single breast nodule in image-to-text-LLM model. This response was recorded as a malignant diagnosis.

Do you know the ACR BI-RADS of breast ultrasound

Yes, I'm familiar with the ACR BI-RADS for breast ultrasound. The **ACR BI-RADS** (Breast Imaging Reporting and Data System) was developed by the American College of Radiology to standardize breast imaging reporting, improve diagnostic accuracy, and ensure clear communication among healthcare providers. This system is used across various breast imaging modalities, including ultrasound, mammography, and MRI, and provides a framework for assessing the likelihood of malignancy and guiding patient management. Here's a summary of the BI-RADS categories:

- BI-RADS 0:** Incomplete - Additional imaging evaluation is needed.
- BI-RADS 1:** Negative - No significant findings.
- BI-RADS 2:** Benign findings - Definitely benign.
- BI-RADS 3:** Probably benign - Follow-up is recommended, as the finding has a low likelihood of malignancy.
- BI-RADS 4:** Suspicious abnormality - Biopsy should be considered; this is subdivided into:
 - 4A:** Low suspicion for malignancy
 - 4B:** Moderate suspicion for malignancy
 - 4C:** High suspicion for malignancy
- BI-RADS 5:** Highly suggestive of malignancy - Biopsy or appropriate intervention is strongly recommended.

If you are a radiologist, here is an ultrasound image of a breast nodule with clear boundaries, lobular composition, marginal spicules, irregular shape, uneven internal echoes, visible punctate hyperechoic areas, no attenuation behind, and punctate color flow within. I would like you to diagnose this breast nodule as either benign or malignant.

Given the ultrasound features described:

- Clear boundaries and lobular composition** can be seen in both benign and malignant lesions.
- Marginal spicules** and an **irregular shape** are often indicative of malignancy.
- Uneven internal echoes** and **punctate hyperechoic areas** (which may represent calcifications) are also suspicious features.
- The absence of **posterior attenuation** could lean toward certain malignancies, though not exclusively.
- Punctate color flow** within the nodule suggests vascularity, which can often be seen in malignancies.


Based on these characteristics, the lesion would likely fall under **BI-RADS 4C or BI-RADS 5**, which indicates a high suspicion of malignancy and a recommendation for biopsy. Features such as spiculated margins, irregular shape, and internal vascularity tend to be more characteristic of malignancies than benign nodules. However, a biopsy would be essential for a definitive diagnosis.

The screenshot demonstrates ChatGPT-4 (OpenAI; <https://chat.openai.com/>) analyzing a single breast nodule in image interpretation model. This response was recorded as a malignant diagnosis.

Do you know the ACR BI-RADS of breast ultrasound

Yes, I'm familiar with the ACR BI-RADS for breast ultrasound. The **ACR BI-RADS** (Breast Imaging Reporting and Data System) was developed by the American College of Radiology to standardize breast imaging reporting, improve diagnostic accuracy, and ensure clear communication among healthcare providers. This system is used across various breast imaging modalities, including ultrasound, mammography, and MRI, and provides a framework for assessing the likelihood of malignancy and guiding patient management. Here's a summary of the BI-RADS categories:

- BI-RADS 0:** Incomplete - Additional imaging evaluation is needed.
- BI-RADS 1:** Negative - No significant findings.
- BI-RADS 2:** Benign findings - Definitely benign.
- BI-RADS 3:** Probably benign - Follow-up is recommended, as the finding has a low likelihood of malignancy.
- BI-RADS 4:** Suspicious abnormality - Biopsy should be considered; this is subdivided into:
 - 4A:** Low suspicion for malignancy
 - 4B:** Moderate suspicion for malignancy
 - 4C:** High suspicion for malignancy
- BI-RADS 5:** Highly suggestive of malignancy - Biopsy or appropriate intervention is strongly recommended.



This is an image of a breast ultrasound, whether you think it's benign or malignant

In examining this ultrasound image with color Doppler flow, several features suggest a high suspicion for malignancy:

- Increased Internal Vascularity:** The nodule exhibits significant internal blood flow, with both red and blue signals, indicating active blood supply within the lesion. Increased vascularity is commonly associated with malignant tumors, as they often induce angiogenesis to support rapid growth.
- Echotexture and Margins:** Although the margins are not completely clear in this view, the internal structure appears heterogeneous, which is often a concerning feature in breast imaging. Malignant nodules tend to have irregular or heterogeneous echotextures, as opposed to the homogeneous texture typically seen in benign lesions.

Based on these observations, this lesion would likely be classified as **BI-RADS category 4C or 5**, indicating a high suspicion of malignancy. A biopsy is strongly recommended to confirm the diagnosis.