

Virtual Reality for the Prevention and Cessation of Nicotine Vaping in Youth: Protocol for a Randomized Controlled Trial

Belinda Borrelli, Daniel Weinstein, Romano Endrighi, Nikki Ling, Kathleen Koval,
Lisa M. Quintiliani, Kaitlyn Konieczny

Submitted to: JMIR Research Protocols
on: February 03, 2025

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 29

0..... 29

0..... 29

Multimedia Appendixes 30

Multimedia Appendix 0..... 30

Virtual Reality for the Prevention and Cessation of Nicotine Vaping in Youth: Protocol for a Randomized Controlled Trial

Belinda Borrelli¹ PhD; Daniel Weinstein¹ MSc; Romano Endrighi¹ PhD; Nikki Ling¹ BS; Kathleen Koval¹ BS; Lisa M. Quintiliani² PhD; Kaitlyn Konieczny¹ BA

¹ Center for Behavioral Science Research Boston University, Henry M. Goldman School of Dental Medicine Boston US

² School of Medicine, Tufts University, Tufts Medical Center Boston US

Corresponding Author:

Belinda Borrelli PhD

Center for Behavioral Science Research
Boston University, Henry M. Goldman School of Dental Medicine
560 Harrison Ave, 3rd floor
Boston
US

Abstract

Background: Only a few trials target high-school aged teens with vaping interventions, typically focusing on prevention rather than cessation, with content limited to vaping harms and refusal skills. Given the co-occurrence and increasing incidence of vaping and mental health issues in teens, both must be simultaneously addressed by vaping interventions.

Objective: The aim of this randomized controlled trial is to evaluate the feasibility and preliminary effectiveness of a Virtual Reality (VR) intervention for teens that targets both determinants of vaping (e.g., mood and stress) and vaping prevention and cessation skills.

Methods: Participants are 150 students attending one of two racially and ethnically diverse Boston-area high schools. Health classes are randomly assigned to VR or Control (assessment only). Students participate if they meet eligibility criteria, assent, and are not opted out by a parent. While Control classes have their regular classroom activity, VR classes are provided with Oculus Meta Quest 2 headsets and engage with the intervention during three class periods. Players solve a mystery with a vaping sub-plot and play five mini-games to teach emotional regulation, coping strategies, and building resilience. A mobile app connects to the game, where players text with their in-game friend, providing a home-based experience to reinforce skills. VR and Control classes complete baseline and post-intervention surveys on the same schedule. Primary outcome measures include vaping awareness, knowledge, attitudes and intention to avoid or quit vaping, engagement with the intervention, and satisfaction with game play. Secondary outcomes include vaping quit attempts, 7-and 30-day point prevalence abstinence, motivation and self-efficacy to quit or avoid vaping, positive and negative affect, emotional regulation and coping, and motion sickness.

Results: Data collection began in April 2024 and concluded in June 2024. Data cleaning has been completed, and analyses are ongoing. Results are expected to be submitted for publication in June 2025.

Conclusions: The multi-sensory VR experience (visual, auditory, and tactile) is ideal for practicing skills, thus facilitating memory and increasing the likelihood of transfer to real world settings. The immersion of VR promotes engagement with the intervention, rather than outside digital temptations. Our study is innovative due to the cluster randomized design, focus on both prevention and cessation, integration of mood management, and implementation among racially and ethnically diverse teens. Incorporation into school-based health curriculums allows for proactive reach. Clinical Trial: NCT06003439

(JMIR Preprints 03/02/2025:71961)

DOI: <https://doi.org/10.2196/preprints.71961>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.
Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org/>

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in <https://www.jmir.org/>



Original Manuscript

Virtual Reality for the Prevention and Cessation of Nicotine Vaping in Youth:
Protocol for a Randomized Controlled Trial

Borrelli, B., Weinstein, D., Endrighi, R., Ling, N. Koval, K., Quintiliani, L., Konieczny, K.

BB., DW., RE., KK., KK, NL. Boston University, Henry M. Goldman School of Dental Medicine,
Center for Behavioral Science Research, Boston, MA 02118.

LQ: Tufts University School of Medicine, Boston MA 02111

Address correspondence to: Belinda Borrelli, PhD, Boston University, Center for Behavioral Science
Research, Henry M. Goldman School of Dental Medicine, 560 Harrison Avenue, 3rd Floor, Boston,
MA 02118. (617) 358-3358; belindab@bu.edu

Word Count: 4,381

Abstract

Background: Only a few trials target high-school aged teens with vaping interventions, typically focusing on prevention rather than cessation, with content limited to vaping harms and refusal skills. Given the co-occurrence and increasing incidence of vaping and mental health issues in teens, both

must be simultaneously addressed by vaping interventions.

Objective: The aim of this randomized controlled trial is to evaluate the feasibility and preliminary efficacy of a Virtual Reality (VR) intervention for teens that targets both determinants of vaping (e.g., mood and stress) and vaping prevention and cessation skills.

Methods: Participants are 150 students attending one of two racially and ethnically diverse Boston-area high schools. Health classes are randomly assigned to VR or Control (assessment only). Students are eligible if not opted out by parent, provided assent, and did not have health issues that preclude participation. While Control classes (and ineligible students within them) have their regular classroom activity, VR classes are provided with Oculus Meta Quest 2 headsets and engage with the intervention during three class periods, once per week. Players solve a mystery with a vaping subplot and play a series of mini-games to teach emotional regulation, coping strategies, and building resilience. A mobile app connects to the VR game, providing a home-based experience to reinforce skills learned. VR and Control classes complete baseline and post-intervention surveys on the same schedule. Primary outcome measures include feasibility, satisfaction and engagement with game play, and vaping awareness, knowledge, attitudes, and intentions to avoid or quit vaping. Secondary outcomes include vaping quit attempts, 7-and 30-day point prevalence abstinence, motivation and self-efficacy to quit or avoid vaping, positive and negative affect, emotional regulation and coping, and motion sickness.

Results: Data collection began in April 2024 and concluded in June 2024. Data cleaning has been completed, and analyses are ongoing. Results are expected to be submitted for publication in August of 2025.

Conclusions: Because VR is visual, auditory and tactile, it is ideal for practicing skills, facilitating memory, and increasing the likelihood of transfer to real world settings. The immersion of VR promotes engagement with the intervention, rather than external digital temptations. Our study is innovative due to the cluster randomized design, focus on both prevention and cessation, integration

of mood management, and implementation among racially and ethnically diverse teens. Incorporation into school-based health curriculums allows for proactive reach.

Clinical Trial Registration: Clinical Trials.gov: NCT 06003439

Keywords: Vaping; teens; virtual reality; prevention; nicotine; coping; mood management; depression

Introduction

The prevalence of e-cigarette or vaping product use among teenagers is a substantial concern, with 10% of middle and high school students in the United States self-reporting current e-cigarette use in 2023, which is approximately 2.8 million students [1]. Moreover, 34.7% of current e-cigarette users report use on at least 20 of the past 30 days [1]. Though many teens believe e-cigarettes are a safer alternative to combustible cigarettes [2], studies suggest e-cigarette use has negative physiological, cognitive, and behavioral effects for teenagers [3]. E-cigarette use is associated with a variety of mental health issues, including depression, internalizing problems, perceived stress and suicidality [4-6]. Moreover, the percent of high school students experiencing persistent feelings of sadness or hopelessness (for at least two consecutive weeks) increased from 28% in 2011 to 42% in 2021 [7, 8]. Therefore, e-cigarette interventions should include emotional regulation and coping in addition to typical approaches that involve teaching vaping refusal skills and managing cravings.

Several intervention approaches have been developed and tested with older adolescents, such as text messaging [9], contingency management [10] and social media approaches [11], but few studies have tested vaping interventions for high school aged teens [12]. One trial among high school age teens randomly assigned participants to watch FDA 'Real Cost' advertisements (on either health harms of vaping or addiction potential) or a control group (neutral videos about vaping) with results showing that both intervention groups had lower self-reported likelihood of future vaping, less positive attitudes towards vaping and vaped fewer days per week compared to those in the control group [13]. In a small pilot study (n=69), text messages regarding the harms of e-cigarettes were sent to teens for 6 days. Outcomes were measured on day 7, with participants reporting increased knowledge and worry regarding the risks of e-cigarette use [14].

Another intervention approach used with high-school aged children is Virtual Reality (VR). VR is an immersive, computer-generated simulation which enables users to navigate within a 3D environment and interact with objects within the environment. One study used a pre-post design to test a VR program that focused on teaching vaping refusal skills (n=47) during two play sessions [15]. The results demonstrated a significant increase in knowledge and perceptions of e-cigarette harm, and a significant decrease in perceived likelihood of using e-cigarettes in the future, but there were no significant changes in attitudes toward e-cigarettes, social norm perceptions, or refusal self-efficacy. However, this study was not randomized and 38 out of the 47 study participants were boys. A follow-up study that randomized students by class, tested the efficacy of the program in 285 middle school students aged 11 to 14 [16]. Results showed that the VR program significantly increased e-cigarette knowledge and perceived addictiveness of e-cigarette at the 6-month follow-up, but there were no differences in e-cigarette behaviors between the control group and the VR group [16].

There are several advantages to virtual reality interventions. VR is a 360-degree experience so distractions by external stimuli are limited (e.g., temptations to check social media). The

immersive nature of VR also promotes learning and behavior change by increasing attention [17] and by creating meaningful emotions and experiences [18]. VR is interactive so that skills are not only learned auditorily and visually but also through tactile interaction. These multiple modes of skill acquisition increase memory retention. Finally, VR is an engaging medium for teens [16].

We describe the protocol for a randomized controlled trial that evaluates the feasibility, engagement, satisfaction, and acceptability of a VR program aimed at the prevention and cessation of vaping among high schoolers, delivered as part of their regular health curriculum over the course of three classes (Clinical Trial Registration#: NCT 06003439). A secondary aim is to assess the preliminary impact of the intervention on the determinants of vaping and vaping behavior. Distinct from the prior VR studies that focused solely on refusal skills for vaping [15], the intervention developed for this present study focuses on the determinants of vaping (emotional regulation, coping strategies, resilience) in addition to prevention and cessation skills (refusal skills, craving management, cessation resources). A mobile app was also developed for use by students at home in order to reinforce the skills learned during the VR experience, and to continue interactions with game characters in between VR sessions. Our study also expands upon prior literature by including a randomized controlled trial design, and by including an ethnically and racially diverse sample. Our overall aim was to test the feasibility, acceptability, and preliminary efficacy of a VR based prevention and cessation intervention program for high school age students. We hypothesize that the intervention will be feasible (e.g., high rate of student assent), satisfactory to students, and that the majority of students will engage with the intervention (e.g., complete all three sessions). Although the trial is not fully powered, we further hypothesize that the intervention will have promising effects in the desired direction on vaping attitudes and behaviors.

Methods

Participants

Eligible participants are assenting students enrolled at one of two high schools in the Greater Boston Area of Massachusetts. Exclusion criteria are: (1) prone to seizures or severe motion sickness; (2) visual impairments that would prevent them from playing VR or using the VR device; (3) parent or guardian opt out; (4) indication by school staff that participation is inappropriate (e.g.,

heart condition, visual or auditory disability, psychiatric disorder, pacemaker, hearing aid, implanted medical device, or any other medical reasons).

Opt-out forms were distributed to eligible students' parent(s) or guardian(s) that detailed the study procedures, potential risks, as well as how to opt their child out of the study. The opt-out form was translated into languages spoken by the families of the children who attended the schools and distributed approximately three weeks prior to obtaining assent from students. Parents or guardians were able to contact either the school or research staff to opt out their child. During their health class, electronic assent forms are given to all students who have not been opted out. Human subjects approval was obtained from our institutional review board (H-43102) and subsequent protocol modifications were submitted to the institutional review board. After informed consent, students complete a baseline survey and data are stored in REDCap (Research Electronic Data Capture), a secure, web-based app designed to support data capture and features to support cleaning, storage and analysis.

Study Design and Random Assignment

Health education classes in the two high schools were randomized to either Intervention (VR) or Control (Assessment only) by a statistician external to the study, using a 1:3 threshold probability. Three classes were randomized in one high school, and 4 classes in the other high school. Regarding the latter high school, two classes were 'yoked' during randomization due to small class sizes. After completing a baseline survey, classes randomized to VR engage with the VR intervention once per week for 3-4 weeks, followed by a follow-up survey. Classes randomized to Control complete baseline and follow-up surveys only, on the same schedule as the VR group. All study procedures take place during regular health classes. Students in Control classes as well as ineligible and non-assenting students in the VR classes engage in alternative assignments regarding vaping or substance use from their respective teachers while study procedures are conducted. Only eligible participants who assented to be in the study complete pre- and post-assessments. Because randomization occurs

at the class-level, teachers cannot be masked to condition. Because of the nature of the intervention, study staff and participants were also not masked to condition.

Equipment

Participants in the VR condition are provided with Meta Quest 2 VR headsets, which do not need to be tethered to computers. Each headset has adjustable head straps and lens spacing, and 3D positional audio. The headset is equipped with cameras to track the movement of the head and body and read hand gestures, so that hand controllers are not needed for navigation through the digital world, to execute game play, or to interact with the digital environment (e.g., grabbing or throwing objects). Head straps and the facial interfaces are disinfected with anti-bacterial wipes after each use.

Virtual Reality Intervention

Our prior mixed methods studies informed the structure and content of the intervention, in which we conducted focus groups and surveys to identify: teens' intrinsic and extrinsic motivators to initiate and maintain vaping; the methods teens currently use to quit vaping and perception of extant vaping cessation treatments; and intervention delivery methods and components that are acceptable to teens [19, 20]. We also conducted usability testing with teens that consisted of playing progressive iterations of the VR program and interviewing them about their play experiences. This helped to identify bugs, ensure functionality, and make content changes.

VR play session procedure:

There are three VR play sessions and each is 30-40 minutes in duration (approximately one class period) and played while seated. Four class periods are reserved for game play so that absent students can 'make up' missed sessions. The VR sessions occur in large areas that allow for entire classes of students to use the VR headsets simultaneously (i.e., school gym or open space in library) without injury (e.g, chairs that are not attached to desks).

Students download the study's mobile app onto their personal devices in order to obtain a login for the VR program and also to engage with game content at home. Students who do not have

compatible devices can be logged into the VR game via study devices. Students are assisted with donning the VR headsets, including adjustment of the head straps and the inter-pupillary distance (IPD) of the lenses, and providing spacers for eyeglasses, if needed. Students are given a brief overview of VR, including how to navigate through a digital environment (e.g., teleporting), how to interact with objects in the environment (e.g., grab, place, and throw objects), how to interact with characters in the environment (e.g., waving to initiate communication with characters) and an overview of the minigames.

VR play sessions:

The three play sessions are connected by a narrative arc that has a main story line (the player solves a mystery at a fictional high school) and a vaping sub-plot. During the vaping subplot, the player chooses a friend at the high school and the player discovers that the friend has a history of getting caught for vaping at school. During the story, the friend experiences triggers to vape, struggles with the decision of whether or not to quit vaping, tries to quit unsuccessfully, and finally applies a constellation of coping and motivational strategies to successfully quit. The player tries to help his or her friend quit vaping through various means, including playing five mini-games to learn about how to cope with negative moods, reduce the friend's craving to vape, and identify treatments for vaping cessation. The mini-games also go beyond vaping to teach general coping skills and building resilience, so that skills have broad applicability to a wide range of risk behaviors and mood problems. At the end of each play session, the player and friend go to a secret cabin in the woods behind the high school which has an "art wall" with additional intervention strategies for at-home practice. A friendly robot, ARAL, guides players through the overall game. A mobile app is connected to the game, where players can view their scores and text message with their in-game friend, providing a home-based experience (outside of class-time) to reinforce skills learned in the classroom-based VR game. No new intervention strategies are presented in the mobile app, rather the text message conversations relate to the events occurring in the VR game, as well as topics discussed

with their fictional friend during game play (e.g., the friend fighting urges to vape and what coping skill they used). Due to technical issues, actual use of the mobile app was not measured.

VR Intervention Content: VR Mini-games and Art Wall Experiences

Rhythm Game: The player learns that physical activity helps reduce stress and cravings to vape. The objective is to hit boxes that fly towards the player to the beat of high energy music while making the correct gesture (thumbs up, thumbs down, or OK symbol) with the correct hand, with game play becoming progressively more difficult over the course of three rounds. *Lunch Line Game:* The player identifies ways to cope with cravings to vape and also helps their friend reduce their cravings to vape (higher points lead to lower cravings on their friend's craving meter). The player grabs the correct coping strategy from the cafeteria shelf to destroy different cravings as they appear on lunch trays that are continuously moving with increasing speed down the lunch line conveyor belt. *Popcorn Game:* The player identifies their own triggers (for vaping, and for negative moods and stress) and learns coping skills (e.g., refusal skills, options for quitting vaping, stress management). While the player and friend are in a movie theater to search for a suspect, ARAL shrinks them down into a popcorn machine. Kernels pop and each one is labeled with a trigger. The player grabs each kernel and throws it into one of two popcorn buckets labeled, "Applies to me" or "Does not apply to me." In the next round, three popcorn buckets appear, each labeled with a coping strategy, and the player must throw the kernel with trigger on it into the bucket with the correct coping strategy, before the 'shot clock' runs out. *Vision Board Game:* Players identify their personal goals (how they envision their future selves) by grabbing and placing them on a vision board, and then they learn how vaping may hinder those goals because a vape cloud appears and moves around the board with increasing speed attempting to remove the goals. Players must use a spray hose to remove the vape cloud before it erases their goals and then they must grab and quickly replace the goals that have been destroyed. *Art Walls:* Intervention strategies include Behavioral Activation [21] (planning and scheduling activities, even if you feel sad), developing a Crisis Survival Kit [22],

engaging in a Deep Breathing exercise with ARAL, and learning about treatment options for vaping. Intervention strategies are delivered by having the player interact with objects (e.g., grabbing, placing) and/or practice them in real time (e.g., Deep Breathing).

Control Classes

These classes complete baseline assessments and post-assessments on the same schedule as the classes randomized to VR, but do not receive the VR intervention or vaping prevention or cessation. Instead, they receive their regular health education curriculum from their teachers, which includes required reading and in-class assignments regarding general substance abuse and vaping was included as a topic.

Measures

Measures are administered via RedCap during class time, using Chromebooks. The follow-up measure is given to both groups approximately one week after the classes randomized to VR complete their final session of the game.

Primary Outcome Measures

Feasibility is assessed as the percent of students who assented, the percent who withdrew from playing the VR game, and the percent who withdrew from the study. Engagement is objectively assessed via the VR headset and includes the number of participants who complete the VR game, the amount of time that the game was played by each participant, and the number of game chapters completed (1 – 10). Satisfaction is assessed with several self-report measures: 1). an overall star rating of the VR program (1 – 5 stars), 2). 7-item scale of gameplay experience (e.g., “I liked the way the game looked”; “I felt connected to my character in the game”), each item rated on a 4-point scale (strongly disagree (1) to strongly agree (4)) [15], and, 3). after each VR game session, participants rate the degree to which they liked each play session and each mini-game within that play session on a four-point scale (“I liked it a lot”; “I liked it”, “Meh”, “Thumbs down”). An adapted measure of perceived program impact is used to assess how much participants agree (1= Strongly disagree to 5=

Strongly agree) with a series of statements regarding the effects of the program on their vaping (awareness, knowledge, attitudes, behavior, intention to change, help-seeking, ability to resist peer pressure), and ability to cope with stress, improve their mood, and avoid use of alcohol and other substances [23].

Secondary Outcome Measures

Vaping-related behaviors are assessed, including attempts to quit vaping since starting the VR program, utilization of quit vaping methods (behavioral, pharmacological, and digital), 7-and 30-day point-prevalence abstinence, and frequency of current vaping (not at all, some days, most days, every day). Motivation to quit vaping (1-10 scale; 1= not at all to 10= very much) and intentions to quit vaping (e.g., Intention to stop in the next 30 days, 6 months, more than six months, no intentions of stopping) are also assessed [24], as well as motivation to use evidence-based vaping cessation resources (“How much do you want to use at least one recommended treatment for stopping vaping (such as nicotine patches, nicotine gum, nicotine lozenges or counseling?)”) [19, 25].

Vaping prevention is assessed with two items regarding the degree of motivation and confidence to avoid vaping (1-10 scales; 1= not at all to 10= very much), one item on intentions to try vaping (e.g., I intend to vape within the next 30 days; 6 months; I may vape at some point in the future but not in the next 6 months) [26], 6-items on self-efficacy to resist vaping across a variety of high-risk situations (e.g., “You feel agitated or tense. Are you confident you will not vape”; Definitely, Maybe, Neutral, Maybe not and Definitely not) [27], and 7-items assessing refusal skills and perceived enjoyability of vaping (“If one of your friends were to offer you a vape, how easy would it be to say no to your friend?”; 4 point scale 1=very easy, 4= very hard) [26].

Determinants of vaping are also measured. Coping is measured with the 16-item Adolescent-Coping Orientation for Problem Experiences scale (A-COPE; e.g., ‘when you face difficulties, how often do you...try to think of the good things in your life; never, hardly ever sometimes, often, most of the time) [28, 29]. Resilience is measured with 6-items regarding the frequency of different

situations (e.g., small problems got me very upset; never, a little, sometimes, often) [30]. Positive affect is assessed with the 8-item PROMIS Pediatric Positive Affect Scale [31], and negative affect is assessed with the 10-item negative affect scale from the Positive and Negative Affect Scale for Children [32].

Tolerability of the VR game is assessed with the 9-item Virtual Reality Sickness Questionnaire [33], which measures the degree to which participants are affected by a variety of symptoms (e.g., eyestrain, headache, fatigue, or dizziness) while playing the VR game.

Data Management

Participants' data are kept confidential according to our institutional regulations. Participants are identified by ID numbers, with all data stored and managed in REDCap in a secure, HIPAA-compliant server. Unanticipated problems and adverse events are recorded on electronic case report forms (eCRF) and reported to our institutional review board in accordance with regulations. Because the study is no more than minimal risk, a data safety and monitoring board was deemed not necessary.

Sample Size

Because this is a pilot feasibility study, a formal power calculation was not conducted. The sample size was determined based on the need to include a sufficient number of students exposed to the intervention, the availability of students in health classes at the participating public schools, and the timeframe allowed by school administrators. This approach allows us to assess our primary outcomes (e.g. feasibility, acceptability, engagement and satisfaction), as well as preliminary impact on relevant vaping-related constructs. We planned on including approximately 80 students in intervention classes and 40 students in control classes. This is based on an average of 25 – 35 students per class. We conservatively estimated that approximately 15 - 20% of students will either not provide assent to be in the study or drop out post-assent. Therefore, we aimed to enroll

approximately 150 participants.

Analytic Plan

Data will be inspected for outliers and descriptive analyses of baseline differences in sociodemographic and vaping-related characteristics between the VR group and the control group will be conducted using t-tests or chi-square tests, as applicable. We will examine differences in key baseline characteristics (e.g., gender distribution between groups) so that they may be covaried in analyses. Feasibility, acceptability, engagement and satisfaction (primary outcomes) in the VR group will be summarized as means and standard deviations for continuous variables, or number and proportion of participants for categorical variables, as appropriate. We will then report the number and proportion of participants who progressed through each of the three play sessions (chapter 1 – 10) of the VR game. Lastly, game ratings obtained after each play session will be reported as number and proportion of participants rating each category ("I liked it a lot"; "I liked it", "Meh", "Thumbs down").

Data on secondary outcomes will be summarized as means and standard deviations for continuous variables, or number and proportion of participants for categorical variables as appropriate and compared between the VR group and the control group (baseline and follow-up). If low cell sizes occur, data might be aggregated to support statistical stability in the models. Differences in secondary outcomes between the groups will be evaluated by computing effect sizes (e.g., Cohen's d or odds ratio) as appropriate.

We will conduct analyses on treatment completers (the subsample of participants who completed the entire VR game) as well intention to treat (as randomized). In addition, we will perform analyses on the entire sample (English learners and English proficient students) as well as conduct analyses separately by English proficiency status. This will allow us to gain invaluable knowledge to inform future research regarding the feasibility of gamified health interventions for students with different level of English proficiency.

Results

Data collection began in April 2024 and ended in June 2024 and 150 participants were recruited (98 intervention, 52 control). Although freshmen and sophomore classes were targeted in the study, there were several juniors and seniors (n=5) in these classes due to school transfers and the need for them to take the mandatory curriculum. Data cleaning and analysis are underway, and results are expected to be submitted for publication August, 2025.

Discussion

Both vaping and mental health issues are on the rise among teens. Given the relationship between vaping and mental health, both must be simultaneously addressed by vaping prevention and cessation interventions. We developed a virtual reality intervention that focuses not only on the skills needed for the prevention and cessation of vaping, but also skills that are needed to cope with common teen mental health issues, such as stress, negative moods, and building resilience. Focusing only on vaping without attention to mental health is not in line with what teens want in a vaping intervention [19, 20, 34] and ignores critical determinants of vaping behavior [35, 36]. Moreover, our virtual reality app is transdiagnostic, in that vaping prevention and cessation skills can be transferred to other risky behaviors, and the general coping skills can be applied to several mental health issues, including depressed mood and anxiety.

The choice of using a VR to deliver the intervention was driven by several factors. First, as VR technology has improved in both graphics and functionality, it is emerging as an engaging and effective method to treat a variety of mental health [37, 38] and physical health issues [39, 40]. Second, VR allows for visual and auditory intervention delivery, and interactivity with simulated objects and people. This multi-sensory delivery is ideal for practicing skills, supporting visual and tactile memory, and increasing the likelihood of transfer to the real world. In our study, we took the additional step of developing a companion mobile app that allowed for continued interaction with the characters from the game between VR sessions (while at home) to further reinforce the skills learned

in the VR sessions. Third, data has shown that teens enjoy games and VR [15, 16, 41] making it an acceptable platform for intervention. Fourth, given the 'cool' factor of many of the vaping device designs and functions, interventions are needed that are equally intriguing to teens. Finally, the immersive nature of VR allows teens to focus exclusively on the intervention, instead of checking their phones, scrolling social media, or engaging with other digital temptations.

Our study is innovative because it randomizes high schoolers at the class-level, targets both prevention and cessation of vaping in teens, integrates mental health coping skills into a vaping prevention and cessation program, and tests VR in racially and ethnically diverse teens. The school-based setting and integration into the regular health curriculum allows for proactive reach to teens, with minimal burden on the school. Because this is a pilot feasibility study, the conclusions will be limited given the small sample size and lack of power to detect effects. In addition, our control group does not control for attention. Using an attention control in VR is costly, requiring additional development or an off the shelf product that would sufficiently match all of the user mechanics (e.g., throwing, grabbing, etc) to ensure a true matched-control. For example, if effective, initial integration into schools would be a financial investment. However, the price of VR headsets continues to decline. Moreover, ten years ago there were no Chrome books integrated into classrooms and now every student has one. Another limitation to VR is that students with visual or auditory disabilities would not be able to fully experience VR, and, as with other learning platforms, accommodations would need to be made for these students. On the other hand, because VR has visual, auditory and tactile learning components, it may be easier for English Learners to learn concepts and skills through VR, rather than written material. Another advantage of VR is remote learning. If students need extra support for a particular lesson, they could take the VR headset home as homework, or if there is another pandemic, VR could be a platform for remote learning, particularly with a multi-player platform (other teachers and students could interact in the same simulated environment).

Importantly, our VR program focuses on both prevention among teens who do not currently vape and quitting among teens who do vape. Therefore, it can be broadly applied in school settings to the general student population and not solely used among those who were ‘caught’ vaping. Ideally, the VR program would be used in conjunction with other school-wide and community-wide policies that discourage vaping for maximum effect on reduction in teenage vaping and associated health outcomes.

Abbreviations

CTRL: Control group

INT: Intervention group

RA: Research assistant

REDCap: Research Electronic Data Capture

VR: Virtual reality

Acknowledgments

The authors would like to express their gratitude to the more than 100 teens who gave their opinion and tested various iterations of this VR game. This research is supported an award from the American Heart Association (20YVNR35500014). The trial sponsors address is: American Heart Association, Research Administration, 7272 Greenville Avenue, Dallas TX USA. This funding agency had no role in the design of this study and will not have any role during its execution, analyses, interpretation of the data, or decision to submit the results. The authors are solely responsible for the design of the research and for the preparation of the manuscript.

Authors' Contributions

BB: Principal Investigator responsible for the conception, overall design and conduct of the trial, principally responsible for developing the intervention content, co-wrote the introduction, methods, measures, and principally wrote the discussion. Reviewed and edited all sections of the manuscript.

DW: Project Manager; co-wrote the methods section. RE: Co-Investigator; co-wrote the measures section, and wrote the data management, sample size and analytic plan, and reviewed and edited all sections of the manuscript. NL, KK: Research assistants who assisted with literature review and

writing the introduction. KK: Research assistant who assisted with writing the methods section. LQ: Co-Investigator; assisted with discussion section. All authors contributed to the current manuscript through review and editing and have approved the final manuscript.

Availability of data and materials

The datasets generated and/or analyzed during the current study will be available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was reviewed and approved by the Boston University Institutional Review board on 10/2/2023 (IRB# 43102). Assent will be obtained from participants via REDcap and parents will be sent an opt-out form, as per school protocol. All data will be kept confidential and stored in a secured location on an institutional server, which is backed up daily. Only certain members of the study staff or others designated by the PI will have access to the study data. Protocol modifications will be reviewed and approved by our Institutional Review Board and communicated to the American Heart Association.

Consent for publication

Not applicable because we will not collect or videos of participants in this study. Additionally, no details on individuals are reported within the manuscript. We do not intend to use professional writers.

Conflicts of Interest

This research is supported an award from the American Heart Association (20YVNR35500014) to Belinda Borrelli, Principal Investigator. The funder did not have a role in the study design, collection, management, analysis or interpretation of data, writing of this report, or decision to submit the report. The authors declare that they have no other competing interests.




References

1. Birdsey J, Cornelius M, Jamal A, Park-Lee E, Cooper MR, Wang J, Sawdey MD, Cullen KA, Neff L: Tobacco Product Use Among U.S. Middle and High School Students - National Youth Tobacco Survey, 2023. *MMWR Morb Mortal Wkly Rep* 2023, 72(44):1173-1182. PMID: 37917558
2. Hilton S, Weishaar H, Sweeting H, Trevisan F, Katikireddi SV: E-cigarettes, a safer alternative for teenagers? A UK focus group study of teenagers' views. *BMJ open* 2016, 6(11):e013271. PMID: 27852721
3. Livingston JA, Chen CH, Kwon M, Park E: Physical and mental health outcomes associated with adolescent E-cigarette use. *J Pediatr Nurs*. 2022, 64:1-17. PMID: 35121206
4. Baiden P, Szlyk HS, Cavazos-Rehg P, Onyeaka HK, Peoples JE, Kasson E: Use of electronic vaping products and mental health among adolescent high school students in the United States: The moderating effect of sex. *J Psychiatr Res*. 2022, 147:24-33. PMID: 35007808
5. Becker TD, Arnold MK, Ro V, Martin L, Rice TR: Systematic Review of Electronic Cigarette Use (Vaping) and Mental Health Comorbidity Among Adolescents and Young Adults. *Nicotine Tob Res*. 2021, 23(3):415-425. PMID: 32905589
6. Lechner WV, Janssen T, Kahler CW, Audrain-McGovern J, Leventhal AM: Bi-directional associations of electronic and combustible cigarette use onset patterns with depressive symptoms in adolescents. *Prev Med*. 2017, 96:73-78. PMID: 28024859
7. CDC. Youth Risk Behavior Survey: data summary & trends report: 2011-2021. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention; 2023. [cdc.gov/healthyyouth/data/yrbs/pdf/YRBS_Data-Summary-Trends_Report2023_508.pdf](https://www.cdc.gov/healthyyouth/data/yrbs/pdf/YRBS_Data-Summary-Trends_Report2023_508.pdf)
8. Mpofu JJ, Underwood JM, Thornton JE, Brener ND, Rico A, Kilmer G, Harris WA, Leon-Nguyen M, Chyen D, Lim C et al: Overview and Methods for the Youth Risk Behavior Surveillance System - United States, 2021. *MMWR Suppl* 2023, 72(1):1-12. PMID: 37104281
9. Graham AL, Amato MS, Cha S, Jacobs MA, Bottcher MM, Papandonatos GD: Effectiveness of a Vaping Cessation Text Message Program Among Young Adult e-Cigarette Users: A Randomized Clinical Trial. *JAMA Intern Med*. 2021, 181(7):923-930. PMID: 33999133
10. Palmer AM, Tomko RL, Squeglia LM, Gray KM, Carpenter MJ, Smith TT, Dahne J, Toll BA, McClure EA: A pilot feasibility study of a behavioral intervention for nicotine vaping cessation among young adults delivered via telehealth. *Drug Alcohol Depend*. 2022, 232:109311. PMID: 35123362
11. Evans WD, Bingenheimer J, Cantrell J, Kreslake J, Tulsiani S, Ichimiya M, D'Esterre AP, Gerard R, Martin M, Hair EC: Effects of a Social Media Intervention on Vaping Intentions: Randomized Dose-Response Experiment. *J Med Internet Res*. 2024, 26:e50741. PMID: 38470468
12. Barnes C, Turon H, McCrabb S, Hodder RK, Yoong SL, Stockings E, Hall AE, Bialek C, Morrison JL, Wolfenden L: Interventions to prevent or cease electronic cigarette use in children and adolescents. *Cochrane Database Syst Rev*. 2023, 11(11):Cd015511. PMID: 37965949
13. Noar SM, Gottfredson NC, Kieu T, Rohde JA, Hall MG, Ma H, Fendinger NJ, Brewer NT: Impact of Vaping Prevention Advertisements on US Adolescents: A Randomized Clinical Trial. *JAMA Netw Open*. 2022, 5(10):e2236370. PMID: 36227597
14. Noar SM, Rohde JA, Horvitz C, Lazard AJ, Cornacchione Ross J, Sutfin EL: Adolescents' receptivity to E-cigarette harms messages delivered using text messaging. *Addict Behav*.

- 2019, 91:201-207. PMID: 29960716
15. Weser VU, Duncan LR, Pendergrass TM, Fernandes CS, Fiellin LE, Hieftje KD: A quasi-experimental test of a virtual reality game prototype for adolescent E-Cigarette prevention. *Addict Behav.* 2021, 112:106639. PMID: 32920455
 16. Weser VU, Duncan LR, Sands BE, Schartmann A, Jacobo S, François B, Hieftje KD: Evaluation of a virtual reality E-cigarette prevention game for adolescents. *Addict Behav.* 2021, 122:107027. PMID: 34225030
 17. Di Serio Á, Ibáñez MB, Kloos CD: Impact of an augmented reality system on students' motivation for a visual art course. *Comput Educ.* 2013, 68:586-596. doi:10.1016/j.compedu.2012.03.002
 18. Huang T-C, Chen C-C, Chou Y-W: Animating eco-education: To see, feel, and discover in an augmented reality-based experiential learning environment. *Comput Educ.* 2016, 96:72-82. doi:10.1016/j.compedu.2016.02.008
 19. Jones E, Endrighi R, Weinstein D, Jankowski A, Quintiliani LM, Borrelli B: Methods used to quit vaping among adolescents and associations with perceived risk, addiction, and socio-economic status. *Addict Behav.* 2023, 147:107835. PMID: 37598643
 20. Weinstein D, Jones E, Endrighi R, Hart JL, Walker KL, Borrelli B, Quintiliani LM: "I knew the effects the whole time." Examination of facilitators and barriers to quit vaping among adolescents. *Addict Behav.* 2024, 157:108087. PMID: 38870591
 21. Alber CS, Krämer LV, Rosar SM, Mueller-Weinitschke C: Internet-Based Behavioral Activation for Depression: Systematic Review and Meta-Analysis. *J Med Internet Res.* 2023, 25:e41643. PMID: 37227760
 22. Rathus JH, Miller AL: DBT skills manual for adolescents: Guilford Publications; 2014.
 23. Stoyanov SR, Hides L, Kavanagh DJ, Zelenko O, Tjondronegoro D, Mani M: Mobile app rating scale: a new tool for assessing the quality of health mobile apps. *JMIR mHealth uHealth.* 2015, 3(1):e27. PMID: 25760773
 24. DiClemente CC, Prochaska JO, Fairhurst SK, Velicer WF, Velasquez MM, Rossi JS: The process of smoking cessation: an analysis of precontemplation, contemplation, and preparation stages of change. *J Consult Clin Psychol.* 1991, 59(2):295-304. PMID: 2030191
 25. Borrelli B, Busch AM, Trotter DRM: Methods Used to Quit Smoking by People with Physical Disabilities. *Rehabil Psychol.* 2013, 58(2):117-123. PMID: 23437992
 26. McCauley DM, Baiocchi M, Cruse S, Halpern-Felsher B: Effects of a short school-based vaping prevention program for high school students. *Prev Med Rep.* 2023, 33:102184. PMID: 37223577
 27. Spek V, Lemmens F, Chatrou M, van Kempen S, Pouwer F, Pop V: Development of a smoking abstinence self-efficacy questionnaire. *Int J Behav Med.* 2013, 20(3):444-449. PMID: 22350635
 28. Patterson JM, H.I. (ed.): Adolescent-Coping Orientation for Problem Experiences (A-COPE). Madison: University of Wisconsin System; 1983.
 29. Patterson JM, McCubbin HI: Adolescent coping style and behaviors: conceptualization and measurement. *J Adolesc.* 1987, 10(2):163-186. PMID: 3611466
 30. Brown LK, Houck C, Lescano C, Donenberg G, Tolou-Shams M, Mello J: Affect regulation and HIV risk among youth in therapeutic schools. *AIDS Behav* 2012, 16(8):2272-2278. PMID: 22669595
 31. Forrest CB, Ravens-Sieberer U, Devine J, Becker BD, Teneralli R, Moon J, Carle A, Tucker CA, Bevans KB: Development and Evaluation of the PROMIS(®) Pediatric Positive Affect Item Bank, Child-Report and Parent-Proxy Editions. *J Happiness Stud.* 2018, 19(3):699-718. PMID: 29760578
 32. Laurent J, Catanzaro SJ, Joiner Jr TE, Rudolph KD, Potter KI, Lambert S, Osborne L,

- Gathright T: A measure of positive and negative affect for children: Scale development and preliminary validation. *Psychol Assess.* 1999, 11(3):326-338. doi: 10.1037/1040-3590.11.3.326
33. Kim HK, Park J, Choi Y, Choe M: Virtual reality sickness questionnaire (VRSQ): Motion sickness measurement index in a virtual reality environment. *Appl Ergon.* 2018, 69:66-73. PMID: 29477332
 34. Bold K, Kong G, Cavallo D, Davis D, Jackson A, Krishnan-Sarin S: School-based E-cigarette cessation programs: What do youth want? *Addict Behav.* 2022, 125:107167. PMID: 34753093
 35. Leventhal AM, Strong DR, Sussman S, Kirkpatrick MG, Unger JB, Barrington-Trimis JL, Audrain-McGovern J: Psychiatric comorbidity in adolescent electronic and conventional cigarette use. *J Psychiatr Res.* 2016, 73:71-78. PMID: 26688438
 36. North C, Loukas A: Depressive symptoms and vaping: The mediating effect of outcome expectancies among young adults. *Drug Alcohol Depend.* 2023, 247:109870. PMID: 37084509
 37. Carl E, Stein AT, Levihn-Coon A, Pogue JR, Rothbaum B, Emmelkamp P, Asmundson GJG, Carlbring P, Powers MB: Virtual reality exposure therapy for anxiety and related disorders: A meta-analysis of randomized controlled trials. *J Anxiety Disord.* 2019, 61:27-36. PMID: 30287083
 38. Katz AC, Norr AM, Buck B, Fantelli E, Edwards-Stewart A, Koenen-Woods P, Zetocha K, Smolenski DJ, Holloway K, Rothbaum BO et al: Changes in physiological reactivity in response to the trauma memory during prolonged exposure and virtual reality exposure therapy for posttraumatic stress disorder. *Psychol Trauma.* 2020, 12(7):756-764. PMID: 32338946
 39. Ahmadpour N, Randall H, Choksi H, Gao A, Vaughan C, Poronnik P: Virtual Reality interventions for acute and chronic pain management. *Int J Biochem Cell Biol.* 2019, 114:105568. PMID: 31306747
 40. Miguel-Rubio A, Rubio MD, Salazar A, Moral-Munoz JA, Requena F, Camacho R, Lucena-Anton D: Is Virtual Reality Effective for Balance Recovery in Patients with Spinal Cord Injury? A Systematic Review and Meta-Analysis. *J Clin Med.* 2020, 9(9). PMID: 32899665
 41. Ridout B, Kelson J, Campbell A, Steinbeck K: Effectiveness of Virtual Reality Interventions for Adolescent Patients in Hospital Settings: Systematic Review. *J Med Internet Res.* 2021, 23(6):e24967. PMID: 34185015

Figure 1. Player Journey in the VR Intervention



This image depicts the player's journey throughout the intervention. The VR app is divided into three play sessions. Each session ends at an 'art wall' that reviews the content learned in that session and how to bring coping skills into the 'real world.' Between VR sessions, there is a home-based component where the player can interact with a mobile app that is connected to the game so they can see their scores and interact with their friend from the game, reinforcing the skills learned in the game. The first VR session introduces the player to ARAL the robot, has the player select a friend and play the first mini-game together, and introduces the main plot (the mystery) and the subplot (the friend uses vapes). In the second VR play session, the player and friend try to solve the mystery and interview suspects in different locations (school cafeteria, movie theater) and two more mini-games are played to further develop coping skills and build resilience. Play Session 3 delivers content via two more mini-games, the players solve the mystery and also engage in an art wall activity that involves discovery of different options for vaping cessation.

Supplementary Files

Untitled.

URL: <http://asset.jmir.pub/assets/4a628664977fd47717512be84057fbdf.docx>

Untitled.

URL: <http://asset.jmir.pub/assets/e50caf9627f26df9b5f33d62b9b636df.pdf>

Multimedia Appendixes

Untitled.

URL: <http://asset.jmir.pub/assets/fd6e1269a2af2753dadd465dc7a14453.pdf>