

Optimization of Internet-Delivered Cognitive Behavioral Therapy for Canadian Leaders within Public Safety: Qualitative Study

Jill AB Price, Hugh C McCall, Sam A Demyen, Shaylee M Spencer, Benjamin MW Katz, Alyssa P Clairmont, Heather D Hadjistavropoulos

Submitted to: Journal of Medical Internet Research
on: February 07, 2025

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 24

 TOC/Feature image for homepages 25

 TOC/Feature image for homepage 0..... 26

Optimization of Internet-Delivered Cognitive Behavioral Therapy for Canadian Leaders within Public Safety: Qualitative Study

Jill AB Price^{1, 2, 3} PhD; Hugh C McCall^{1, 2, 3} PhD; Sam A Demyen^{1, 2, 3} BA; Shaylee M Spencer^{1, 2, 3} MSW; Benjamin MW Katz^{1, 2, 3} BA; Alyssa P Clairmont^{1, 2, 3} BA; Heather D Hadjistavropoulos^{1, 2, 3} PhD

¹University of Regina Canadian Institute for Public Safety Research and Treatment Regina CA

²University of Regina PSPNET Regina CA

³University of Regina Department of Psychology Regina CA

Corresponding Author:

Jill AB Price PhD

University of Regina

Canadian Institute for Public Safety Research and Treatment

3737 Wascana Parkway

Regina

CA

Abstract

Background: Canadian public safety personnel (PSP) report high rates of mental health concerns and barriers to treatment. PSPNET is a clinical research unit that offers internet-delivered cognitive behavioral therapy (ICBT) that is free, confidential, and developed with and for PSP. Treatment outcomes are promising with clinically significant symptom improvement (eg, anxiety, depression, posttraumatic stress) and favorable treatment satisfaction. While these results are promising, research has yet to explore ways to optimize therapist-guided ICBT for leaders within public safety. Optimizing ICBT for leaders is particularly important given their widespread organizational impact that can affect other PSP.

Objective: This study included 10 clients who self-identified as being in a supervisory or leadership position within their public safety organization and completed either the therapist-guided PSP Wellbeing Course or PSP PTSD Course. We used descriptive statistics to analyze demographics, mental health symptoms, treatment engagement, and treatment satisfaction. We used a reflexive thematic analysis of semi-structured interview transcripts to assess leaders' course perceptions and feedback.

Methods: Qualitative data was collected via 10 semi-structured interviews with clients who self-identified as being in a supervisory or leadership position within their public safety organization and completed either the therapist-guided PSP Wellbeing Course or PSP PTSD Course. These interviews assessed leaders' mental health stressors and course perceptions.

Results: Canadian leaders within public safety reported occupational and non-occupational stressors and enrolled in ICBT to support their own or colleagues' mental health. Most clients enrolled in the PSP Wellbeing Course, accessed 4 of 5 lessons (n=7;70%), engaged with therapist support (n=7;70%), and identified as employed (n=8;80%), White (n=8;80%), and men (n=7;70%) with an average age of 45 years. At pre-treatment, 80% of clients endorsed clinically significant symptoms of one or more disorder; most often depression (n=7;70%) and/or anger (n=6;60%). Clients reported favorable attitudes toward the ICBT courses with most reporting that they were satisfied with the course (n=9;90%). Feedback to further optimize ICBT content for leaders included development of a leader case story (n=6;60%) and new resources to help leaders apply skills learned in ICBT within the context of their leadership roles (n=4;40%). Leaders also recommended optimizing ICBT delivery by improving the platform technology and incorporating more multimedia.

Conclusions: Canadian leaders within public safety perceived therapist-guided ICBT developed with and for PSP as a suitable treatment option for their needs and identified ways to further optimize its content and delivery. Future research should investigate the impacts of these efforts and explore optimizing ICBT for other groups of clients. Clinical Trial: [Clinicaltrials.gov](https://clinicaltrials.gov) NCT04127032; <https://www.clinicaltrials.gov/ct2/show/NCT04127032>; [Clinicaltrials.gov](https://clinicaltrials.gov) NCT04335487; <https://clinicaltrials.gov/study/NCT04335487>

(JMIR Preprints 07/02/2025:72321)

DOI: <https://doi.org/10.2196/preprints.72321>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

Please make my preprint PDF available to anyone at any time (recommended).

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

✓ **No, I do not wish to publish my submitted manuscript as a preprint.**

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [http://www.jmir.org/](#)

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Original Paper

Jill AB Price, PhD, jill.price@uregina.ca^{1,2,3}

Hugh C McCall, PhD, hugh.mccall@uregina.ca^{1,2,3}

Sam A Demyen, BA, sam.demyen@uregina.ca^{1,2,3}

Shaylee M Spencer, shaylee.spencer@uregina.ca^{1,2}

Benjamin MW Katz, BSc, bmwkatz@gmail.com^{1,2,3}

Alyssa P Clairmont, BA, alyssapclairmont@gmail.com^{1,2,3}

Heather D Hadjistavropoulos, PhD, heather.hadjistavropoulos@uregina.ca^{1,2,3}

¹Canadian Institute for Public Safety Research and Treatment, Regina, SK, Canada

²PSPNET, University of Regina, Regina, SK, Canada

³Department of Psychology, University of Regina, Regina, SK, Canada

Optimization of Internet-Delivered Cognitive Behavioral Therapy for Canadian Leaders within Public Safety: Qualitative Study

Abstract

Background: Canadian public safety personnel (PSP) report high rates of mental health concerns and barriers to treatment. PSPNET is a clinical research unit that offers internet-delivered cognitive behavioral therapy (ICBT) that is free, confidential, and developed with and for PSP. Treatment outcomes are promising with clinically significant symptom improvement (eg, anxiety, depression, posttraumatic stress) and favorable treatment satisfaction. While these results are promising, research has yet to explore ways to optimize therapist-guided ICBT for leaders within public safety. Optimizing ICBT for leaders is particularly important given their widespread organizational impact that can affect other PSP.

Objectives: This study investigated (1) the perceived mental health stressors of Canadian leaders within public safety; (2) the degree to which leaders perceived existing therapist-guided ICBT courses tailored for PSP (ie, *PSP Wellbeing Course* and *PSP PTSD Course*) as suitable for their needs; and (3) ways to further optimize therapist-guided ICBT for public safety leaders.

Methods: This study included 10 clients who self-identified as being in a supervisory or leadership position within their public safety organization and completed either the therapist-guided *PSP Wellbeing Course* or *PSP PTSD Course*. We used descriptive statistics to analyze demographics, mental health symptoms, treatment engagement, and treatment satisfaction. We used a reflexive thematic analysis of semi-structured interview transcripts to assess leaders' course perceptions and feedback.

Results: Canadian leaders within public safety reported occupational and non-occupational stressors and enrolled in ICBT to support their own or colleagues' mental health. Most clients enrolled in the *PSP Wellbeing Course*, accessed 4 of 5 lessons ($n=7;70\%$), engaged with therapist support ($n=7;70\%$), and identified as employed ($n=8;80\%$), White ($n=8;80\%$), and men ($n=7;70\%$) with an average age of 45 years. At pre-treatment, 80% of clients endorsed clinically significant symptoms of one or more disorder; most often depression ($n=7;70\%$) and/or anger ($n=6;60\%$). Clients reported favorable attitudes toward the ICBT courses with most reporting that they were satisfied with the course ($n=9;90\%$). Feedback to further optimize ICBT content for leaders included development of a leader case story ($n=6;60\%$) and new resources to help leaders apply skills learned in ICBT within the context of their leadership roles ($n=4;40\%$). Leaders also recommended optimizing ICBT delivery by improving the platform technology and incorporating more multimedia.

Conclusions: Canadian leaders within public safety perceived therapist-guided ICBT developed with and for PSP as a suitable treatment option for their needs and identified ways to further optimize its

content and delivery. Future research should investigate the impacts of these efforts and explore optimizing ICBT for other groups of clients.

Trial **Registrations:** Clinicaltrials.gov NCT04127032;
<https://www.clinicaltrials.gov/ct2/show/NCT04127032>; Clinicaltrials.gov NCT04335487;
<https://clinicaltrials.gov/study/NCT04335487>

Keywords: first responders; public safety personnel; leadership; leaders; mental health; internet interventions; digital mental health interventions; internet-delivered cognitive behavioral therapy

Introduction

Background

Public safety personnel (PSP) are individuals whose work protects, supports, and secures the safety of citizens within border services, communications (eg, dispatch), corrections, emergency managers, fire (career and volunteer), intelligence, paramedics, policing (self-administered, municipal, provincial, and federal), and search and rescue, among others [1]. PSP report high rates of mental health challenges; in a survey of 5,813 Canadian PSP, 44.5% reported clinically significant symptoms of one or more mental disorders [2]. PSP also experience barriers to accessing mental healthcare, including structural barriers (eg, financial constraints, limited availability of services, distance from services, inconvenience of seeking treatment) and attitudinal barriers (eg, concerns about stigma, low perceived need for help) [3,4].

Occupational Stressors

Occupational stressors refer to workplace challenges that impact mental health [1]. PSP's occupational stressors have been categorized into operational stressors, which are directly tied to tasks in the workplace (eg, potentially psychologically traumatic events [PPTEs], job-related injuries, infringement on other parts of life, excessive demands) and organizational stressors, which are associated with the setting or context of the job (eg, staff shortages, lack of training and resources, inconsistent leadership) [1,5–8]. PPTEs refer to events that have the potential to induce symptoms of posttraumatic stress disorder (PTSD) or related disorders (eg, events involving death, serious injury, sexual violence) [1]. Among Canadian PSP, PPTEs are both common and predictive of symptoms of various mental health challenges [5].

Leaders' Mental Health

The term “leader” refers to an individual in a supervisory or management position in their workplace [9]. Despite perceptions that leaders are more resilient than non-leaders [10], have greater access to mental health supports [10,11], have greater work-life balance, and have more control over their work schedule and demands [12], mental health problems appear to be highly prevalent among leaders in the general public [13,14]. Campbell and colleagues (2007) found that 88% of leaders reported that their job is the main source of stress in their lives, 79% reported increased stress since taking on a supervisory or management position, and over 60% claimed that their organization was not providing sufficient mental health support [13]. Research similarly shows that workplace leaders experience greater occupational stressors than staff in non-leadership roles [15]. Compared with non-leaders, leaders are also more likely to get inadequate amounts of sleep (less than 6 hours) and self-report consuming more alcohol at work, both of which can predict disruptive and negative leadership behaviors and worsen pre-existing symptoms of mental disorders [10].

Despite the high rates of mental health concerns among PSP and the unique stressors faced by organizational leaders, relatively little is known about the unique challenges faced by PSP in leadership roles. This is an important gap in the research literature because mental health concerns among PSP leaders are not only inherently distressing but also have important implications for the

mental health and wellbeing of PSP in non-leadership roles. Research shows that mental health challenges among leaders—within the general public or public safety—can have widespread organizational impacts that affect non-leaders' job performance and satisfaction [10,14,16–18]. Mental health challenges can negatively impact leaders' behavioral functioning (eg, aggression, sleep) and cognitive functioning (eg, concentration, decision-making), and there is evidence that emotional difficulties among leaders in the general population (eg, anger, anxiety, depression) predict abusive management practices and unreasonable expectations of staff [10]. PSP further report that their experiences of distress and burnout are impacted by the quality of relationships with their leaders [17]. It is important to note that leaders' mental health is also affected by the mental health and morale of their staff [10], highlighting a need for mental health support among PSP in leadership and non-leadership roles alike.

Internet-Delivered Cognitive Behavioral Therapy

Internet-delivered cognitive behavioral therapy (ICBT) is an evidence-based treatment for various mental health symptoms (eg, anger, anxiety, depression, posttraumatic stress) that has demonstrated excellent outcomes among the general public [19,20] and Canadian PSP [21,22]. ICBT typically consists of self-directed learning of evidence-based treatment principles in online modules, often accompanied by therapist support via email or phone. This online, largely self-directed format affords privacy and convenience, can allow therapists to treat more clients at a given time, is cost-effective, and can, in turn, help overcome several barriers to treatment (eg, distance from services, cost of treatment, privacy concerns) [23,24].

PSPNET is a clinical research unit located at the University of Regina in Saskatchewan, Canada that was founded in response to the Government of Canada's National Action Plan on Posttraumatic Stress Injuries [25]. PSPNET provides free, confidential, and evidence-based ICBT developed with and for Canadian PSP. PSPNET offers a range of ICBT courses, including courses with or without optional therapist support, PTSD-specific and transdiagnostic courses (ie, courses designed to treat symptoms of various mental disorders), courses in English and French, and courses for both PSP and their spouses or significant others. PSPNET's research has shown that Canadian PSP, in general, report favorable perceptions of ICBT [3,18,26,27] as well as good treatment engagement, symptom reduction, and treatment satisfaction [22,28–31]. Although there is some evidence that leaders within public safety have favorable perceptions of ICBT [18,32], little is known about their unique mental health stressors and how ICBT might be further optimized to address those stressors. Interviews were invaluable in PSPNET's early efforts to tailor ICBT for PSP, but to date, interviews have not been conducted specifically with leaders to gain insight into their unique challenges, experiences, and treatment preferences [26].

Objectives and Hypotheses

The current study explored ways to optimize ICBT for Canadian leaders within public safety by investigating: (1) leaders' mental health stressors; (2) the degree to which leaders perceived existing therapist-guided ICBT programs (ie, *PSP Wellbeing Course* and *PSP PTSD Course*, both described further below) tailored for PSP as suitable for their unique needs; and (3) leaders' feedback on how to further optimize therapist-guided ICBT to better meet their treatment needs. We made three hypotheses. First, consistent with research showing high rates of various mental health problems among PSPNET clients [21,22] and Canadian PSP more broadly [2], we hypothesized that PSP leaders would report a variety of mental health concerns. Second, consistent with research showing positive perceptions of ICBT among PSPNET clients [21,22,28,30], Canadian PSP leaders [18,32], and Canadian PSP at varying levels of leadership [3,18,26], we hypothesized that participants in the current study would report largely positive perceptions of therapist-guided ICBT. Third, consistent with our past research using qualitative analysis of interviews with PSP to identify

areas for improvement in ICBT [26,28,33,34], we hypothesized that participants would provide tangible feedback to further optimize therapist-guided ICBT.

Methods

Ethical Considerations

This qualitative study was approved by the Research Ethics Board at the University of Regina (#2019-157) and was carried out within the context of a longitudinal single-group open observational trial (NCT04127032) and open cohort preference trial (NCT04335487) registered on Clinicaltrials.gov. Clients provided informed consent after being made aware of the study details and the potential benefits and risks of participation. Clients were provided access to the *PSP Wellbeing Course* or *PSP PTSD Course* but were not offered other incentives to encourage participation. Client data were stored on a secure server and were de-identified prior to analyses.

Study Design

Prospective clients signed up for a PSPNET account, provided informed consent to participate, and completed eligibility screening questionnaires that collected demographic, clinical, and occupational information. This online screening was followed by a telephone interview, after which eligible clients were enrolled in an ICBT course appropriate to their symptoms and preferences. The current study only included clients enrolled in the English versions of the therapist-guided *PSP Wellbeing Course* or *PSP PTSD Course*.

Intervention

The *PSP Wellbeing Course* and *PSP PTSD Course* are therapist-guided ICBT courses designed to be completed in eight weeks. Each are based on ICBT courses, called the *Wellbeing Course* and *PTSD Course*, that were originally developed by the eCentre Clinic at Macquarie University in Australia and have since shown excellent outcomes in Australia [35] and Canada [30]. The *PSP Wellbeing Course* was initially tailored for PSP following a series of interviews, focus groups, and questionnaires exploring their unique needs [3,26] and has since been continuously evaluated and iteratively improved based on client feedback (eg, [28]). The *PSP Wellbeing Course* is a transdiagnostic course that is designed to help with symptoms of anxiety, depression, and posttraumatic stress, whereas the *PSP PTSD Course* is a disorder-specific course that focuses on symptoms of posttraumatic stress. Both courses include psychoeducation and coping skills rooted in cognitive behavioral therapy and delivered via a combination of text, diagrams, videos, audio files, downloadable practice worksheets and activities, illustrative case stories, frequently asked questions, additional resources addressing issues not addressed in the core course content (eg, problems with anger, alcohol use, and sleep), and optional therapist support via phone or secure email. Each of these courses are described in greater detail elsewhere (see [21] for the *PSP Wellbeing Course* and [22] for the *PSP PTSD Course*).

Measures

During online screening, prospective clients completed a Participant Information Questionnaire that solicited demographic (eg, age, gender, ethnicity, and community size) and occupational data (eg, PSP sector, years of experience, leadership role). Clients also completed a series of questionnaires assessing symptoms of various mental disorders, which are presented in Table 1 below. Finally, clients completed a bespoke Treatment Satisfaction Questionnaire at 8-weeks post-enrollment. Detailed information about these measures—and several others not germane to the purposes of the current study—can be found in previous publications [21,36] or our open observational trial (NCT04127032) and open cohort preference trial (NCT04335487) registered on

Clinicaltrials.gov.

Eligibility Criteria

In order to be included in either the *PSP Wellbeing Course* or the *PSP PTSD Course* and, in turn, the current study, clients were required to (a) be at least 18 years of age; (b) be a current or former PSP; (c) have internet access; and (d) reside in a Canadian province where PSPNET offers therapist-guided ICBT services (ie, New Brunswick, Nova Scotia, Ontario, Prince Edward Island, Québec, or Saskatchewan). Additional details regarding eligibility criteria are described in previous papers [21,22] and our open observational trial (NCT04127032) and open cohort preference trial (NCT04335487) registered on Clinicaltrials.gov. Further to these criteria, clients were only included in the current study if they indicated that they were in a supervisory or leadership role within their public safety organization. Semi-structured interviews took place between January and July 2024.

Semi-Structured Interview

Clients were invited via email to participate in a semi-structured interview at six weeks post-enrollment, which allowed two weeks to schedule interviews for as early as possible after their completion of the 8-week ICBT programs. Of the 23 clients invited, 10 leaders within public safety completed the interview. Author JABP conducted interviews with the first 10 clients interested via phone or video call, with each lasting 20 to 30 minutes. Interview questions were piloted with the first client and then discussed with author HDH to ensure clarity and effectiveness before proceeding. Clients provided verbal consent for the interviews and their recording. First, clients were asked to describe stressors that impact their wellbeing. Clients were then asked questions assessing their perceptions of and feedback on ICBT: (1) course material; (2) case stories; (3) resources; (4) therapist support; and (5) overall impressions. Given the semi-structured nature of the interview, clients were asked follow-up questions where appropriate to gain further insight.

Course material questions assessed which topics and skills were most and least beneficial as a leader in public safety and then solicited suggestions for new topics and skills that would be relevant and beneficial for clients in leadership positions. Case stories questions explored whether leaders accessed at least one of the stories and, if so, whether case stories were perceived as authentic and relatable experiences, beneficial to leaders (ie, informing, comforting, modeling, persuading, and engaging) [37], and whether a leader-specific story is recommended. Resource questions assessed whether leaders accessed at least one additional resource and, if so, which resources were perceived as the most and least beneficial for leaders and whether a new leader-specific resource is recommended. Therapist support questions assessed whether leaders accessed the optional therapist support and, if so, preferred modality of therapist support (ie, email or phone), whether therapist support met their expectations, and general feedback. Lastly, overall impression questions assessed clients' satisfaction with their selected therapist-guided ICBT program, perceived benefits and negative impacts of participation, likelihood of recommending therapist-guided ICBT to other PSP in both leadership and non-leadership roles, and additional feedback not covered in previous questions.

Clients

Demographic data was collected via the Participant Information Questionnaire and analyzed using SPSS (version 28; IBM Corp). Results showed that eligible clients predominantly enrolled in the *PSP Wellbeing Course* ($n=9;90\%$) and self-identified as employed ($n=8;80\%$), White ($n=8;80\%$), and men ($n=7;70\%$) with an average age of 45 years. Half of the clients reported working within policing ($n=5;50\%$) and half in various other public safety sectors. Clients resided in communities with less than ($n=6;60\%$) and greater than 100,000 citizens ($n=4;40\%$) across various Canadian provinces. At pretreatment, 80% of clients endorsed clinically significant symptoms of one or more mental disorders, with 50% endorsing clinically significant symptoms of three or more mental

disorders. In total, 70% of clients endorsed clinically significant symptoms of depression, 60% endorsed clinically significant symptoms of anger, 50% endorsed clinically significant symptoms of generalized anxiety, 30% endorsed clinically significant symptoms of posttraumatic stress, 10% endorsed clinically significant problems with alcohol and drug use. Further details surrounding sample characteristics are presented in Table 1.

Table 1. Sample characteristics.

Characteristic		Clients (N=10)
Age (years), mean (SD)		45.3 (6.0)
Gender, n (%)		
	Women	3 (30.0)
	Men	7 (70.0)
Ethnicity, n (%)		
	White	8 (80.0)
	Other	2 (20.0)
Community size, n (%)		
	< 100,000 citizens	6 (60.0)
	> 100,000 citizens	4 (40.0)
Employment status, n (%)		
	Employed	8 (80.0)
	Unemployed	2 (20.0)
PSP^a sector, n (%)		
	Police	5 (50.0)
	Other	5 (50.0)
Years in PSP field, mean (SD)		17.2 (6.4)
Clinically significant symptoms at pre-treatment, n (%)		
	Depression (Patient Health Questionnaire-9; PHQ-9; cut-off score ≥ 10) [38]	7 (70.0)
	Anger (Dimensions of Anger Reactions-5; DAR-5; cut-off score ≥ 12) [39]	6 (60.0)
	Anxiety (Generalized Anxiety Disorder-7; GAD-7; cut-off score ≥ 10) [40]	5 (50.0)
	Posttraumatic stress (PTSD Checklist for DSM-5; PCL-5; cut-off score ≥ 33) [41]	3 (30.0)
	2-item Panic Disorder Severity Scale – Self-Report (2-item PDSS-SR; cut-off score ≥ 3) [42]	3 (30.0)
	Social anxiety (Mini-Social Phobia Inventory; Mini-SPIN; cut-off score ≥ 6) [43]	2 (20.0)
	Alcohol use (Alcohol Use Disorder Identification Test;	1 (10.0)

	AUDIT; cut-off score ≥ 20) [44]	
	Drug use (Drug Use Disorder Identification Test; DUDIT; cut-off score ≥ 25) [45]	1 (10.0)
	Clinically significant symptoms on at least one measure	8 (80.0)
	Clinically significant symptoms on at least three measures	5 (50.0)

^aPSP: public safety personnel.

Analyses

Qualitative data ($N=10$) were sourced from open-ended questions via the semi-structured interviews which were de-identified and transcribed by author SAD. Transcriptions were then analyzed using the NVivo 20 analysis software [46] using a reflexive thematic approach. The initial codebook was created by the primary coder, author JABP, to align with questions posed during the semi-structured interviews. These themes were then re-evaluated to identify, analyze, and report on any subthemes [47]. The authors acknowledge that we hold certain preconceived attitudes and beliefs about ICBT and recognize that, as non-PSP, our understanding of PSP experiences is limited. To help minimize bias, themes were reviewed by author SAD to identify and discuss any conflicts. Themes were then discussed among all other co-authors for diverse perspectives. Upon consensus of coding, each theme was assessed for semantic meaning and given an appropriate name [48]. Client demographics were pulled from Data from the closed-ended survey questions were imported into IBM SPSS Statistics v.26 [42], and descriptive statistics were computed

Results

Stressors and Reasons for Seeking Treatment

Qualitative analyses identified two reasons why leaders within public safety enrolled in the *PSP Wellbeing Course* ($n=9$) and *PSP PTSD Course* ($n=1$): (1) to seek support for their mental health ($n=6$;60%); and (2) to review the course for their members ($n=4$;40%). Leaders cited two themes of stressors that impact their mental health and wellbeing: (1) occupational ($N=10$; 100%); and (2) non-occupational ($n=3$;30%). Occupational stressors were further categorized into operational (ie, related to tasks in the workplace; $n=7$;70%) or organizational stressors (ie, related to the job setting or context; $N=10$;100%). The operational stressor most reported was workload ($n=7$;70%). One client said that “The challenge I find as a leader is I work a lot of extra hours” while another stated “I’m on call 24 hours a day”. Additionally, clients reported operational stress due to PPTE exposures ($n=5$;50%). One client explained, “in leadership positions...you get exposed to a lot of things” while another client elaborated, “I’ve been through riots... finding dead bodies.”

Clients further identified a variety of organizational stressors that impact their wellbeing. Most clients highlighted the continued presence and impacts of mental health stigma ($n=7$;70%) within their organizations. One client said, “I’ve been in the service for 29 years and the only place it’s safe to have a meltdown is at home or behind closed doors.” while other clients clarified, “Creating a safe space for a leader is really, really difficult because you don’t know who you can trust.” and “If you come out and say you have problems, you’re damaged goods in their eyes. So, you keep that to yourself.” Many clients reported lack of support within their organization ($n=6$;60%), stating, “My officers know they can come to me at any time with any issues whether its work or personal. I don’t get that from my leaders” and “There’s no [union] representation of being in a management position whereas frontline workers have representation.” Clients indicated concerns related to staffing ($n=5$;50%) with one stating, “Most people who would merit that type of a position don’t want the job” and another reporting that “My staff are stretched fairly thin.” Clients reported the impacts of guilt because of their job ($n=4$;40%) with one explaining that “You are the ones who are setting up

the plans and the ones who are directing people to act and so it's going to be a lot of guilt and trauma.” Another client elaborated that “You’re always worried about the safety of the people under you, so you always carry that burden. You want to make sure they’re happy, cared for, looked after, their families are looked after, so you carry that a little bit more.” Clients also highlighted concerns related to interpersonal difficulties ($n=4;40\%$), “I don’t know how to handle him anymore. I’m beyond being able to deal with him... Dealing with him on a day-to-day basis left me with no energy to deal with stuff at home.” and middle management ($n=3;30\%$), “The stressors come from below as well as above...it comes uphill and it goes downhill and we’re sort of the in between.” Lastly, clients identified the impact of resources ($n=3;30\%$), “I’m burning through cash...I’ve been addressing some of the staffing issues we have, which has been great, but I’ve been doing it double time and there’s only a finite amount of money.” Non-occupational stressors were recognized by many, but not all, clients ($n=3;30\%$) with all identifying familial concerns specific to the death of a loved one ($n=1;33\%$), interpersonal violence ($n=1;33\%$), or marital problems ($n=1;33\%$). More information about these occupational and non-occupational stressors are discussed in Table 2.

Table 2. Occupational and non-occupational stressors.

Themes	Example quotes	Count, n (%)
Occupational stressors: operational stressors		7 (10.0)
Workload	“The challenge I find as a leader is I work a lot of extra hours.”	7 (70.0)
PPTes	“I’ve been through riots...finding dead bodies.”	5 (50.0)
Occupational stressors; organizational stressors		10 (100.0)
Mental Health Stigma	“I’ve been in the service for 29 years and the only place it’s safe to have a meltdown is at home or behind closed doors.”	7 (70.0)
Lack of Support	“My officers know they can come to me at any time with any issues whether its work or personal. I don’t get that from my leaders.”	6 (60.0)
Staffing Challenges	“Most people who would merit that type of a position don’t want the job.”	5 (50.0)
Guilt	“You are the ones who are setting up the plans and the ones who are directing people to act and so it's going to be a lot of guilt and trauma.”	4 (40.0)
Interpersonal Difficulties	“I’m beyond being able to deal with him...Dealing with him on a day-to-day basis left me with no energy to deal with stuff at home.”	4 (40.0)
Middle Management	“The stressors come from below as well as above...it comes uphill and it goes downhill and we’re sort of the in between”	3 (30.0)
Lack of Resources	“I’m burning through cash...I've been addressing some of the staffing issues we have, which has been great, but I've been doing it double time and there's only a finite amount of money.”	3 (30.0)
Non-Occupational Stressors		3 (30.0)

	Familial	“My wife and I were struggling back and forth, and it has started to affect me, and us, and our kids...and just everything in general.”	3 (30.0)
--	----------	---	----------

Positive Feedback

Clients generally engaged well with ICBT, as most accessed at least the first 4 of 5 lessons in their selected course ($n=7/10;70\%$), completed post-treatment mental health measures ($n=9/10;90\%$), and utilized optional therapist support ($n=7/10;70\%$) via phone ($n=3/7;42\%$), email ($n=1/7;14\%$), or both ($n=3/7;42\%$). Clients' perceptions of ICBT were largely positive. Most clients reported in the treatment satisfaction questionnaire that the course was worth their time ($n=8/10;80\%$), that they were satisfied with the course overall ($n=9;90\%$), and that they would recommend the course to other PSP ($N=10/10;100\%$). One client stated that “Every lesson has given me something as a first responder leader”, while another recognized that “It is changing my life. I feel less angry and less frustrated.” Of those who had contact with their clinician ($n=7/10;70\%$), most clients reported that their therapist support met or exceeded their expectations ($n=6/7;86\%$). One client explained “I think I can be helpful to other leaders if they find themselves in this position.”

Skills most commonly cited as helpful were controlled breathing ($n=6/10;60\%$) followed by challenging unhelpful thoughts ($n=4/10;40\%$) and graduated exposure ($n=4/10;40\%$). Many clients reported positive perceptions of the course delivery ($n=6/10;60\%$), highlighting the program's flexibility ($n=4/6;67\%$) structure ($n=2/6;33\%$), free-of-cost availability ($n=2/6;33\%$), confidentiality ($n=1/6;17\%$), and audio features ($n=1/6;17\%$). For example, one client stated, “You don't have to go out somewhere to meet with somebody or those kinds of things like for an introvert like me, it's perfect that it's like this”.

All clients accessed at least one case story ($N=10/10;100\%$), and most accessed at least one additional resource ($n=6/10;60\%$). Clients perceived the case stories as authentic representations of PSP ($n=9/10;90\%$) that were relatable to their own experiences ($n=9/10;90\%$). One client explained, “I've seen my staff go through some of the things that they were talking about...So, I could sort of go through the story and empathize with it and possibly be able to use that to help my staff.” Most clients agreed that the case stories modeled ($n=9/10;90\%$) and motivated use of targeted behaviour learned in the course ($n=9/10;90\%$). One client stated that, “Just reading it...was one thing but then to kind of take those stories and put them into a real-life situation, that absolutely helps.” Clients also described stories as helping to normalize mental health symptoms ($n=8/10;80\%$), with one client highlighting that “Knowing it's so common makes it much easier to start facing things.” Moreover, most clients recognized that the case stories increased understanding of and engagement with the course material ($n=7/10;70\%$). One client stated, “[The stories] put it into perspective what the material was teaching us so like okay I can see how this applies to a situation or to real life.” Some clients also recognized the case stories as a source of mental health information ($n=4/10;40\%$).

Constructive Feedback

Case Stories

Clients provided tangible feedback on ways to optimize therapist-guided ICBT for leaders via the case stories, additional resources, and delivery. Most clients ($n=6;60\%$) recommended a new story tailored for leaders within public safety. One client stated, “Yeah, it could be beneficial. Especially if...you're in that middle or senior position.” while another stated “It could be an interesting avenue if you wanted to do a leadership version of this. So, the story is not about you. The story is about somebody on your team.” Two clients were unsure if a leadership story is necessary with one elaborating, “We do have unique issues but there's also similarities and I think...how they deal with them is going to be the same. So, it's just that we're more isolated.” One client recommended removing the stories entirely, stating, “Get rid of them...It makes you feel like – well

this person has it worse than you”.

Additional Resources

Clients ($n=4;40\%$) provided topic suggestions for the additional resources. Three clients recommended a leadership-specific resource with two clients recommending ways to apply the skills learned to a leadership position. Another client requested tools to build confidence, explaining, “For leadership...the huge thing is confidence, self-awareness, and being able to project that confidence and your abilities and skills.” One client also recommended a resilience-based resource that provided tools and skills to maintain mental health. Lastly, one client suggested an unconscious bias resource, explaining, “Unconscious biases seems to be like a real cancer because everybody has a different one and you don’t know what it is. So, that means everybody’s discriminated against in some way and it makes moving forward in any capacity hard.”

Delivery

Additionally, clients ($n=7;70\%$) provided recommendations for tailoring the course delivery. Three clients highlighted technological challenges navigating the online platform. One client stated that “I don't know if this was myself or the system. The site would load fairly slowly. And it would crash a lot.” while another said, “As an older person, technology's not my forte...This isn't just with your course it's with everything.” Two clients suggested incorporating more audiovisuals, with one explaining, “There is sections of the course that are narrated, and they do have someone talking to you and kind of walking you through it, and I liked those parts the best. That’s where I absorbed the most information, I don’t absorb the most if I’m just reading a screen.” Two clients also revealed challenges with the pace of the course, with one explaining that “You kind of feel like oh, I should get through this lesson quick because I have another one.” and another stating, “It would sure be nice if I didn't have to give the survey on the module I hadn't done yet.”

Discussion

Principal Results

Overview

ICBT is an effective and convenient mental health treatment among the general population [19,20]. To increase awareness of and access to ICBT, PSPNET was developed with and for PSP across Canada. Previous results show favorable mental health outcomes, treatment engagement, and treatment satisfaction [21,30]. Leaders previously recognized the need and suitability of PSPNET for Canadian PSP and provided advice to optimize implementation generally [32]. This study expands on these findings by exploring ways to optimize implementation of therapist-guided ICBT specifically among Canadian PSP in leadership roles.

Stressors and Reasons for Seeking Treatment

Leaders reported concerns with occupational and non-occupational stressors impacting their wellbeing. Occupational stressors included both operational stressors (ie, workload and PPTes) and organizational stressors (ie, mental health stigma, lack of support, staffing challenges, guilt, interpersonal difficulties, middle management, and lack of resources) [16]. While these findings are broadly consistent with previous research showing the prevalence and diversity of occupational stressors within public safety organizations, it is notable that leaders in the current study described a higher proportion organizational stressors relative to operational stressors than did a general sample of PSP who were not necessarily leaders in a past study [33]. PSP’s most salient stressors may then shift from operational to organizational in nature as they progress into leadership roles. Some leaders

also reported non-occupational stressors specific to their family. Leaders enrolled in PSPNET most frequently described seeking support for their mental health, with some enrolling instead to review the course for other employees in their organizations. Therefore, despite leadership stereotypes [49], the current study shows that Canadian PSP in leadership role are susceptible to occupational and non-occupational stressors and may have interest in seeking therapist-guided ICBT to support their mental health (among other reasons).

Positive Feedback

Leaders in this sample showed good treatment engagement and satisfaction, including satisfaction with therapist support, which is consistent with findings of prior research on ICBT tailored for PSP at varying levels of leadership [21,22]. Self-managing mental health symptoms appears to be a priority among leaders just as it is among PSP more broadly [34], although several leaders indicated that they were reviewing the course before recommending it to colleagues. Many leaders in the current study cited controlled breathing, challenging unhelpful thoughts, and graduated exposure as helpful skills learned in PSPNET's courses.

Case Stories

Previous research found that many (ie, 69%;[34]), but not all, PSP accessed at least one case story in ICBT compared to all leaders in the current study. Leaders, in general, report increased social isolation [14,15,50] and decreased social supports [15] compared to non-leaders. Perhaps leaders are then more likely to engage in ICBT case stories as a result. Nevertheless, most leaders indicated that the case stories represented authentic and relatable experiences [34,51], but still recommended the development of a new story tailored to their needs. Research supports tailoring case stories to current and prospective clientele as it may improve story interest and engagement [52,53]. Therefore, the current study highlights the need for and potential benefits of a leadership-based case story in the *PSP Wellbeing Course*. Specifically, content could include: (1) mental health stigma; (2) lack of support; (3) staffing challenges; and (4) guilt, among other topics provided in Table 2.

Additional Resources

Leaders similarly recommended the development of a new resource that provides information on application of the content and skills learned in the course to address their unique stressors. Offering a variety of additional resources provides a flexible way for clients to access psychoeducation and strategies relevant to their needs, without requiring them to seek external support or engage with content irrelevant to their needs [54]. Additional resources can also improve treatment engagement and satisfaction by allowing clients to personalize treatments [54]. This recommendation may then offer promising outcomes given that additional resources are perceived as a valued component of ICBT even though previous research [34] and the current study show that not all clients access them. A leadership-specific resource could support leaders in applying cognitive behavioral therapy skills to manage the unique issues they identified (eg, how to use thought challenging to address unhelpful thoughts that induce feelings of guilt in leaders, how to use assertive communication skills to manage interpersonal difficulties with staff).

Delivery

Leaders continue to recognize the benefits of delivering cognitive behavioral therapy via an online platform. The delivery feature most appreciated by leaders in this study was flexibility, allowing clients to work at their own pace in the privacy of their home or office, with optional

therapist support. This finding is consistent with previous research highlighting the need for flexible mental health support for PSP [26,28,32]. For example, McCall and colleagues (2021) found that PSP prefer therapist-guided ICBT that is delivered within a flexible timeframe for course completion and with a choice of communication modes with their assigned therapist. Some leaders also remarked on their appreciation that PSPNET is free and confidential with the accessibility option of audio narratives. To further optimize ICBT delivery, leaders generally recommended improving the platform technology and adding audiovisual content. Technological barriers are a common concern with internet interventions [55] because they play a critical role in treatment engagement and satisfaction [56]. Audiovisual content can also improve treatment engagement and satisfaction by improving accessibility options [57]. Yet, little is known about PSP dropout rates in ICBT and whether they mirror the general population.

Limitations and Future Directions

The current study has several limitations that can help inform future research directions. First, most clients self-identified as White, men, and working in policing. In the largest study of Canadian PSP we are aware of ($N = 5,813$), 92% of the sample identified as White, 67% reported being male, and 52% identified as either municipal/provincial police or Royal Canadian Mounted Police personnel [2], suggesting that our sample is relatively representative of the demographic makeup of Canadian PSP.¹ Nevertheless, our total sample size was very small, and the number of participants who were not White, men, and/or working in policing was particularly small, limiting the diversity and representativeness of responses and perspectives. Second, sampling bias may be a risk, as clients with positive perceptions of PSPNET may have been more likely to complete questionnaires and agree to participate in interviews. As such, clients with negative perceptions of PSPNET may not have been included in the current study. Additionally, this study did not consider the role of mental health history (symptoms or services) nor type of leadership role (eg middle or upper management) on leaders' perceptions of ICBT. Future research could include a larger sample size to improve sample diversity with a particular focus on sector, gender, ethnicity, mental health history, and types of leadership roles. Quantitative analyses could then explore the role that demographics play in leaders' preconceived beliefs of ICBT. A follow-up thematic reflexive analysis is also recommended to investigate the impact of a leader-specific case story and additional resource on ICBT engagement, mental health outcomes, and satisfaction.

Conclusion

This study provides insight on the suitability and optimization of therapist-guided ICBT for Canadian leaders within public safety. Most leaders reported enrolling in PSPNET for support with occupational and non-occupational stressors impacting their wellbeing and symptoms of depression, anger, anxiety, posttraumatic stress, and other concerns. Some leaders reported enrolling in PSPNET to review the course for their staff. Nevertheless, most leaders held positive perceptions of therapist-guided ICBT and provided tangible feedback to further optimize its implementation, focusing on the case stories, additional resources, and delivery. Overall, these findings are promising given the important role that leaders play in the promotion, implementation, and sustainability of programs and services [9,58].

Acknowledgements

The authors would like to acknowledge the contributions of the eCentre Clinic at Macquarie University, Australia, who helped develop the original version of the *Wellbeing Course* and *PTSD*

¹ These statistics were not directly reported by the authors but were calculable based on data provided in Table 1 of their paper.

Course; and the Online Therapy Unit for their support in setting up PSPNET. The authors would also like to acknowledge all working groups, whose contributions are integral to the development of this course; all first responders and other PSP who participated in this study; and all PSPNET staff for their support with this project.

Funding Statement:

PSPNET is part of the Canadian Institute for Public Safety Research and Treatment (CIPSRT), which is funded by the Canadian Government's Ministry of Public Safety and Emergency Preparedness. The funder was not involved in the study design or in the collection, analysis, or interpretation of the data.

Conflicts of Interest

None declared.

Abbreviations

ICBT: internet-delivered cognitive behavioral therapy

PPTE: potentially psychologically traumatic event

PSP: public safety personnel

PTSD: posttraumatic stress disorder

Data Availability

The data described in this study are not publicly available due our commitment to our participants to securely store their data and refrain from sharing them with anyone outside of our research team.

References

1. Heber A, Testa V, Groll D, et al. Glossary of terms: A shared understanding of the common terms used to describe psychological trauma, version 3.0. *Health Promot Chronic Prev Can.* 2023;43(10-11):S1-S999. doi:10.24095/hpcdp.43.10/11.09
2. Carleton RN, Afifi TO, Turner S, et al. Mental disorder symptoms among public safety personnel in Canada. *Can J Psychiatry.* 2018;63(1):54-64. doi:10.1177/0706743717723825
3. McCall HC, Sison AP, Burnett JL, Beahm JD, Hadjistavropoulos HD. Exploring Perceptions of Internet-Delivered Cognitive Behaviour Therapy among Public Safety Personnel: Informing Dissemination Efforts. *Int J Environ Res Public Health.* 2020;17:1-17. doi:10.3390/ijerph17176026
4. Ricciardelli R, Carleton RN, Mooney T, Cramm H. "Playing the system": Structural factors potentiating mental health stigma, challenging awareness, and creating barriers to care for Canadian public safety personnel. *Health (N Y).* 2020;24(3):259-278. doi:10.1177/1363459318800167
5. Carleton RN, Afifi TO, Taillieu T, et al. Exposures to potentially traumatic events among public safety personnel in Canada. *Can J Behav Sci Rev Can Sci Comport.* 2019;51(1):37-52. doi:10.1037/cbs0000115
6. Finney C, Stergiopoulos E, Hensel J, Bonato S, Dewa CS. Organizational stressors associated with job stress and burnout in correctional officers: a systematic review. *BMC Public Health.*

2013;13(1):82. doi:10.1186/1471-2458-13-82

7. Ricciardelli R, Carleton RN, Groll D, Cramm H. Qualitatively Unpacking Canadian Public Safety Personnel Experiences of Trauma and Their Well-Being. *Can J Criminol Crim Justice*. 2018;60(4):566-577. doi:10.3138/cjccj.2017-0053.r2
8. Waters JA, Ussery W. Police stress: history, contributing factors, symptoms, and interventions. Burke R, ed. *Polic Int J Police Strateg Manag*. 2007;30(2):169-188. doi:10.1108/13639510710753199
9. Knaak S, Luong D, McLean R, Szeto A, Dobson KS. Implementation, Uptake, and Culture Change: Results of a Key Informant Study of a Workplace Mental Health Training Program in Police Organizations in Canada. *Can J Psychiatry*. 2019;64(1_suppl):30S-38S. doi:10.1177/0706743719842565
10. Barling J, Cloutier A. Leaders' mental health at work: Empirical, methodological, and policy directions. *J Occup Health Psychol*. 2017;22(3):394-406. doi:10.1037/ocp0000055
11. Hobfoll SE. Social and Psychological Resources and Adaptation. *Rev Gen Psychol*. 2002;6(4):307-324. doi:10.1037/1089-2680.6.4.307
12. Sherman GD, Lee JJ, Cuddy AJC, et al. Leadership is associated with lower levels of stress. *Proc Natl Acad Sci*. 2012;109(44):17903-17907. doi:10.1073/pnas.1207042109
13. Campbell M, Baite JI, Martin A, Meddings K. *The Stress of Leadership*. Center for Creative Leadership; 2007:1-16. <https://eminencecoach.com/wp-content/uploads/2016/04/StressofLeadership.pdf>
14. Octavia J, Prihadi KD, Yeoh HC, Purwaningtyas EK. They can handle it, they are leaders: a look into organizational leaders' mental health. *Int J Public Health Sci IJPHS*. 2022;11(4):1439. doi:10.11591/ijphs.v11i4.21719
15. Wallis A, Robertson J, Bloore RA, Jose PE. Differences and similarities between leaders and nonleaders on psychological distress, well-being, and challenges at work. *Consult Psychol J Pract Res*. 2021;73(4):325-348. doi:10.1037/cpb0000214
16. Carleton RN, Afifi TO, Taillieu T, et al. Assessing the Relative Impact of Diverse Stressors among Public Safety Personnel. *Int J Environ Res Public Health*. 2020;17(4):1234. doi:10.3390/ijerph17041234
17. Harms PD, Credé M, Tynan M, Leon M, Jeung W. Leadership and stress: A meta-analytic review. *Leadersh Q*. 2017;28(1):178-194. doi:10.1016/j.leaqua.2016.10.006
18. Landry CA, Beahm JD, McCall HC, Hadjistavropoulos HD. Sustaining and Expanding Internet-Delivered Cognitive Behavioral Therapy (ICBT) for Public Safety Personnel across Canada: A Survey of Stakeholder Perspectives. *Int J Environ Res Public Health*. 2023;20(8):5592. doi:10.3390/ijerph20085592
19. Andersson G, Carlbring P, Titov N, Lindefors N. Internet Interventions for Adults with Anxiety and Mood Disorders: A Narrative Umbrella Review of Recent Meta-Analyses. *Can J Psychiatry Rev Can Psychiatr*. 2019;64(7):465-470. doi:10.1177/0706743719839381

20. Lewis C, Roberts NP, Simon N, Bethell A, Bisson JJ. Internet-delivered cognitive behavioural therapy for post-traumatic stress disorder: systematic review and meta-analysis. *Acta Psychiatrica Scand.* 2019;140(6):508-521. doi:10.1111/acps.13079
21. Hadjistavropoulos HD, McCall HC, Dear BF, Beahm JD, Carleton RN, Titov N. Outcomes of transdiagnostic internet-delivered cognitive behavioural therapy tailored to public safety personnel: A longitudinal observational study. *J Anxiety Disord.* 2024;104:1-12. doi:10.1016/j.janxdis.2024.102861
22. McCall HC, Dear BF, Landry C, et al. Internet-delivered cognitive behavioural therapy for symptoms of PTSD among public safety personnel: Initial outcomes of an open cohort preference trial of transdiagnostic and disorder-specific therapy. *Internet Interv.* 2023;33:100656. doi:10.1016/j.invent.2023.100656
23. Christensen H, Hickie IB. Using e-health applications to deliver new mental health services. *Med J Aust.* 2010;192(S11). doi:10.5694/j.1326-5377.2010.tb03695.x
24. Richards D, Duffy D, Blackburn B, et al. Digital IAPT: the effectiveness & cost-effectiveness of internet-delivered interventions for depression and anxiety disorders in the Improving Access to Psychological Therapies programme: study protocol for a randomised control trial. *BMC Psychiatry.* 2018;18(1):59. doi:10.1186/s12888-018-1639-5
25. Public Safety Canada . *Supporting Canada's Public Safety Personnel: An Action Plan on Post-Traumatic Stress Injuries.* Government of Canada; 2019. <https://www.publicsafety.gc.ca/cnt/rsrscs/pblctns/2019-ctn-pln-ptsj/index-en.aspx>
26. McCall HC, Beahm JD, Fournier AK, Burnett JL, Carleton RN, Hadjistavropoulos HD. Stakeholder perspectives on internet-delivered cognitive behavioural therapy for public safety personnel: A qualitative analysis. *Can J Behav Sci Rev Can Sci Comport.* 2021;53(3):232-242. doi:10.1037/cbs0000242
27. McCall HC, Landry CA, Ogunade A, Carleton RN, Hadjistavropoulos HD. Why Do Public Safety Personnel Seek Tailored Internet-Delivered Cognitive Behavioural Therapy? An Observational Study of Treatment-Seekers. *Int J Environ Res Public Health.* 2021;18(22):1-14. doi:10.3390/ijerph182211972
28. Beahm JD, McCall HC, Carleton RN, Titov N, Dear B, Hadjistavropoulos HD. Insights into internet-delivered cognitive behavioural therapy for public safety personnel: Exploration of client experiences during and after treatment. *Internet Interv.* 2021;26:1-13. doi:10.1016/j.invent.2021.100481
29. Hadjistavropoulos HD, Reiser SJ, Beahm JD, et al. Internet-Delivered Cognitive Behavioral Therapy Tailored to Spouses and Significant Others of Public Safety Personnel: Formative Evaluation Study. *JMIR Form Res.* 2023;7:1-20. doi:10.2196/51088
30. Hadjistavropoulos HD, McCall HC, Thiessen DL, et al. Initial Outcomes of Transdiagnostic Internet-Delivered Cognitive Behavioral Therapy Tailored to Public Safety Personnel: Longitudinal Observational Study. *J Med Internet Res.* 2021;23(5):1-13. doi:10.2196/27610
31. McCall HC, Hadjistavropoulos HD. Impact of an Online Discussion Forum on Self-Guided Internet-Delivered Cognitive Behavioral Therapy for Public Safety Personnel: Randomized

Trial. *J Med Internet Res*. 2024;26:1-23. doi:10.2196/59699

32. Beahm JD, McCall HC, Carleton RN, Jones N, Hadjistavropoulos HD. Examining how organizational leaders perceive internet-delivered cognitive behavioural therapy for public safety personnel using the RE-AIM implementation framework. *Internet Interv*. 2024;35:1-9. doi:10.1016/j.invent.2024.100718
33. Beahm JD, Landry CA, McCall HC, Carleton RN, Hadjistavropoulos HD. Understanding and Addressing Occupational Stressors in Internet-Delivered Therapy for Public Safety Personnel: A Qualitative Analysis. *Int J Environ Res Public Health*. 2022;19(8):4744. doi:10.3390/ijerph19084744
34. Price JA, Gregory J, McCall HC, Landry CA, Beahm JD, Hadjistavropoulos HD. Client Perspectives of Case Stories in Internet-Delivered Cognitive Behavioral Therapy for Public Safety Personnel: Mixed Methods Study. *JMIR Form Res*. 2024;8:e64454. doi:10.2196/64454
35. Titov N, Dear BF, Staples LG, et al. MindSpot Clinic: An Accessible, Efficient, and Effective Online Treatment Service for Anxiety and Depression. *Psychiatr Serv*. 2015;66(10):1043-1050. doi:10.1176/appi.ps.201400477
36. Landry CA, McCall HC, Beahm JD, et al. Web-Based Mindfulness Meditation as an Adjunct to Internet-Delivered Cognitive Behavioral Therapy for Public Safety Personnel: Mixed Methods Feasibility Evaluation Study. *JMIR Form Res*. 2024;8:1-18. doi:10.2196/54132
37. Shaffer VA, Zikmund-Fisher BJ. All Stories Are Not Alike: A Purpose-, Content-, and Valence-Based Taxonomy of Patient Narratives in Decision Aids. *Med Decis Making*. 2013;33(1):4-13. doi:10.1177/0272989X12463266
38. Kroenke K, Spitzer RL, Williams JBW, Löwe B. The Patient Health Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: a systematic review. *Gen Hosp Psychiatry*. 2010;32(4):345-359. doi:10.1016/j.genhosppsych.2010.03.006
39. Forbes D, Alkemade N, Mitchell D, et al. Utility of the dimensions of anger reactions-5 (Dar-5) scale as a brief anger measure. *Depress Anxiety*. 2014;31(2):166-173. doi:10.1002/da.22148
40. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing Generalized Anxiety Disorder: The GAD-7. *Arch Intern Med*. 2006;166(10):1092. doi:10.1001/archinte.166.10.1092
41. Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and Initial Psychometric Evaluation: Posttraumatic Stress Disorder Checklist for DSM-5. *J Trauma Stress*. 2015;28(6):489-498. doi:10.1002/jts.22059
42. Forsell E, Kraepelien M, Blom K, et al. Development of a very brief scale for detecting and measuring panic disorder using two items from the Panic Disorder Severity Scale-Self Report. *J Affect Disord*. 2019;257:615-622. doi:10.1016/j.jad.2019.07.057
43. Connor KM, Kobak KA, Churchill LE, Katzelnick D, Davidson JRT. Mini-SPIN: A brief screening assessment for generalized social anxiety disorder. *Depress Anxiety*. 2001;14(2):137-140. doi:10.1002/da.1055

44. Saunders JB, Aasland OG, Babor TF, De La Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction*. 1993;88(6):791-804. doi:10.1111/j.1360-0443.1993.tb02093.x
45. Berman AH, Bergman H, Palmstierna T, Schlyter F. Evaluation of the drug use disorders identification test (DUDIT) in criminal justice and detoxification settings and in a Swedish population sample. *Eur Addict Res*. 2005;11(1):22-31. doi:10.1159/000081413
46. Lumivero. Nvivo Qualitative Data Analysis Software for Windows, Version 14. 2023;14.
47. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006;3(2):77-101. doi:10.1191/1478088706qp063oa
48. Salminen J, Santos JM, Kwak H, An J, Jung S gyo, Jansen BJ. Persona Perception Scale: Development and Exploratory Validation of an Instrument for Evaluating Individuals' Perceptions of Personas. *Int J Hum-Comput Stud*. 2020;141:102437. doi:10.1016/j.ijhcs.2020.102437
49. Cloutier AE, Barling J. Expectations of Leaders' Mental Health. *J Leadersh Organ Stud*. 2023;30(3):276-296. doi:10.1177/15480518231178637
50. Quick JC, Gavin JH, Cooper CL, Quick JD. Executive health: Building strength, managing risks. *Acad Manag Perspect*. 2000;14(2):34-44. doi:10.5465/ame.2000.3819304
51. Hadjistavropoulos HD, Hill TG, Philipp-Muller AE, Dear B, Titov N. Perceptions of client stories in internet-delivered cognitive behaviour therapy: A mixed-methods evaluation. *Internet Interv*. 2023;34:100692. doi:10.1016/j.invent.2023.100692
52. Salminen J, Nielsen L, Jung SG, An J, Kwak H, Jansen BJ. "Is More Better?": Impact of Multiple Photos on Perception of Persona Profiles. In: *Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems*. ACM; 2018:1-13. doi:10.1145/3173574.3173891
53. Nielsen L. *Personas - User Focused Design*. Springer London; 2019. doi:10.1007/978-1-4471-7427-1
54. Hadjistavropoulos HD, Peynenburg V, Sapkota RP, Titov N, Dear BF. Evaluation of additional resources used in therapist-assisted transdiagnostic internet-delivered cognitive behaviour therapy. *Internet Interv*. 2024;37:100758. doi:10.1016/j.invent.2024.100758
55. Smoktunowicz E, Barak A, Andersson G, et al. Consensus statement on the problem of terminology in psychological interventions using the internet or digital components. *Internet Interv*. 2020;21(1):1-8. doi:10.1016/j.invent.2020.100331
56. Moskalenko MY, Hadjistavropoulos HD, Katapally TR. Barriers to patient interest in internet-based cognitive behavioral therapy: Informing e-health policies through quantitative analysis. *Health Policy Technol*. 2020;9(2):139-145. doi:10.1016/j.hlpt.2020.04.004
57. Henka A, Zimmerman G. Personas Based Accessibility Testing: Towards User-Centererd Accessibility Evaluation. In: Vol 435. ; 2014:226-231. doi:10.1007/978-3-319-07854-0

58. Milliard B. Utilization and Impact of Peer-Support Programs on Police Officers' Mental Health. *Front Psychol.* 2020;11:1686. doi:10.3389/fpsyg.2020.01686



Supplementary Files

TOC/Feature image for homepages

Staff of the Office of Public Safety working in the Operations Centre in Charlottetown.

