

Leveraging Aging Service Providers to Support iCBT for Depression in Homebound Older Adults: Protocol for a Type 1 Hybrid Effectiveness-Implementation Trial

Xiaoling Xiang, Samson Ash, Skyla Turner, Xinyin Zhang, Elyse Narbut, Ruopeng An, Jennifer N. Jester, Sunggeun Park, Joseph A. Himle

Submitted to: JMIR Research Protocols
on: February 21, 2025

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript..... 5

Supplementary Files..... 31

 Figures 32

 Figure 1..... 33

 Multimedia Appendixes 34

 Multimedia Appendix 1..... 35

 TOC/Feature image for homepages 36

 TOC/Feature image for homepage 0..... 37

Leveraging Aging Service Providers to Support iCBT for Depression in Homebound Older Adults: Protocol for a Type 1 Hybrid Effectiveness-Implementation Trial

Xiaoling Xiang¹ MSW, PhD; Samson Ash¹ MSW; Skyla Turner¹ MSW; Xinyin Zhang²; Elyse Narbut²; Ruopeng An³ PhD; Jennifer N. Jester¹ PhD; Sunggeun Park¹ PhD; Joseph A. Himle¹ PhD, MSW

¹ School of Social Work University of Michigan–Ann Arbor Ann Arbor US

² University of Michigan–Ann Arbor Ann Arbor US

³ Silver School of Social Work New York University New York US

Corresponding Author:

Xiaoling Xiang MSW, PhD

School of Social Work
University of Michigan–Ann Arbor
1080 S University
Ann Arbor
US

Abstract

Background: Homebound older adults face a high burden of depression and significant barriers to accessing mental health treatments. Few interventions address their specific needs. Empower@Home, an internet-based cognitive behavioral therapy (iCBT) program, was co-designed with stakeholders and tailored to older adults. The program includes nine self-paced online sessions to be completed within 12 weeks, augmented by telephone-based human support. Efficacy studies have demonstrated its acceptability and effectiveness in reducing depression when supported by trained research staff. However, its real-world effects and feasibility for integration into community aging service settings remain unknown.

Objective: This study aims to assess the real-world effectiveness of Empower@Home integrated into aging services, using non-clinician agency staff as coaches to support older adults. A secondary objective is to evaluate the implementation process through a qualitative process evaluation.

Methods: The study is a Type 1 Hybrid Effectiveness-Implementation Trial with a two-arm randomized controlled design. A total of 256 homebound older adults will be recruited from three community aging service agencies, and agency staff will be trained as coaches to support iCBT use. Participants in the treatment group will receive Empower@Home immediately, while the control group will receive biweekly friendly calls and enhanced care as usual, including the provision of psychoeducational materials and usual care. Outcomes will be assessed at baseline, post-intervention (12 weeks), and at two follow-up points (24 and 36 weeks). The primary outcome is change in depressive symptoms as measured by the Patient Health Questionnaire-9 (PHQ-9). Secondary outcomes include changes in social isolation, health-related quality of life, disability burden, pain intensity, and anxiety symptoms.

Results: Institutional Review Board approval was obtained in August 2024, and recruitment began in October 2024. Recruitment is expected to conclude by April 2028. Upon trial completion, the effectiveness of Empower@Home on primary and secondary outcomes will be analyzed.

Conclusions: This study will provide critical insights into the real-world effectiveness of Empower@Home. If successful, it offers a scalable, cost-effective model for integrating technology-assisted mental health treatments into community aging services, enhancing access to care for an underserved, hard-to-reach population. Clinical Trial: ClinicalTrials.gov. NCT06584422. Registered on August 30, 2024. <https://clinicaltrials.gov/study/NCT06584422>

(JMIR Preprints 21/02/2025:72953)

DOI: <https://doi.org/10.2196/preprints.72953>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in [JMIR Publications](#)

No. Please do not make my accepted manuscript PDF available to anyone.

Original Manuscript

Leveraging Aging Service Providers to Support iCBT for Depression in Homebound Older Adults: Protocol for a Type 1 Hybrid Effectiveness-Implementation Trial

Xiaoling Xiang, Ph.D., MSW, Samson Ash, MSW, Skyla Turner, MSW, LMSW, Xinyin Zhang, Elyse Narbut, Ruopeng An, PhD, Jennifer N. Jester, PhD, Sunggeun Park, PhD, Joseph A. Himle, Ph.D., MSW

Corresponding Author:

Xiaoling Xiang <https://orcid.org/0000-0002-4926-4707>

Address: 1080 S University Ave, Ann Arbor, MI, 48109

Phone: (734) 763-6581; Fax: (734) 763-3372

Email: xiangxi@umich.edu

Abstract

Background:

Homebound older adults face a high burden of depression and significant barriers to accessing mental health treatments. Few interventions address their specific needs. Empower@Home, an internet-based cognitive behavioral therapy (iCBT) program, was co-designed with stakeholders and tailored to older adults. The program includes nine self-paced online sessions to be completed within 12 weeks, augmented by telephone-based human support. Efficacy studies have demonstrated its acceptability and effectiveness in reducing depression when supported by trained research staff. However, its real-world effects and feasibility for integration into community aging service settings remain unknown.

Objective:

This study aims to assess the real-world effectiveness of Empower@Home integrated into aging services, using non-clinician agency staff as coaches to support older adults. A secondary objective is to evaluate the implementation process through a qualitative process evaluation.

Methods:

The study is a Type 1 Hybrid Effectiveness-Implementation Trial with a two-arm randomized controlled design. A total of 256 homebound older adults will be recruited from three community aging service agencies, and agency staff will be trained as coaches to support iCBT use. Participants in the treatment group will receive Empower@Home immediately, while the control group will receive biweekly friendly calls and enhanced care as usual, including the provision of psychoeducational materials and usual care. Outcomes will be assessed at baseline, post-intervention (12 weeks), and at two follow-up points (24 and 36 weeks). The primary outcome is change in depressive symptoms as measured by the Patient Health Questionnaire-9 (PHQ-9). Secondary outcomes include changes in social isolation, health-related quality of life, disability burden, pain intensity, and anxiety symptoms.

Results:

Institutional Review Board approval was obtained in August 2024, and recruitment began in October 2024. Recruitment is expected to conclude by April 2028. Upon trial completion, the effectiveness of Empower@Home on primary and secondary outcomes will be analyzed.

Conclusion:

This study will provide critical insights into the real-world effectiveness of Empower@Home. If successful, it offers a scalable, cost-effective model for integrating technology-assisted mental health treatments into community aging services, enhancing access to care for an underserved, hard-to-reach population.

Keywords: depression; older adults; homebound; eHealth; mHealth; digital mental health; internet intervention; iCBT; implementation effectiveness; aging services

Trial Registration: ClinicalTrials.gov. [NCT06584422](https://clinicaltrials.gov/study/NCT06584422). Registered on August 30, 2024. <https://clinicaltrials.gov/study/NCT06584422>

Protocol: iCBT for Older Adults

Introduction

Background

Depression among homebound older adults is a critical yet often overlooked public health issue [1,2]. Research has shown that 50% of homebound older adults experience clinically significant depression. Of these, 14% face major depression, a rate seven times higher than that of their non-homebound counterparts [3,4]. Regardless of its severity, untreated and undertreated depression has serious health consequences for older adults. It can increase the risk of suicide, physical disabilities, cognitive impairment, hospitalizations, and premature mortality [5–8]. Compounding these challenges, homebound older adults often face substantial access barriers to mental health services due to socio-economic disadvantages and physical disabilities [9–11].

Late-life depression can be effectively treated with antidepressants, psychotherapies, or a combination of both [12,13]. However, the aging process is often associated with an increased risk of chronic illness, functional limitations, social isolation, and reduced coping resources. Psychotherapies can be particularly useful as they not only address symptoms but also teach coping skills to better address these challenges [10,14]. Unfortunately, few older adults receive psychotherapy treatments due to stigma, a shortage of trained mental health providers, and systemic barriers, including inadequate insurance coverage for subthreshold depression [15–17]. Without proper treatment, depression and physical disability can reinforce each other, ultimately worsening both health and overall quality of life. [6].

Aging Network agencies have unmatched reach to the target population, covering nearly every community in the US and serving over 8 million older Americans annually. Consequently, integrating mental health treatment into the Aging Network holds significant potential to increase access to mental health services for homebound older adults [10]. Previous research has already demonstrated that non-clinician aging service providers can be effectively trained to deliver a range of depression interventions, including those based on behavioral activation, psychoeducation, problem-solving, and depression self-management [18–25]. For example, Choi et al. showed that aging service providers could effectively administer behavioral activation via teleconference [26,27]. However, depending on these providers to directly administer structured treatments poses several challenges, such as limited expertise in depression care, heavy caseloads, and frequent turnover.

Leveraging digital mental health interventions (DMHIs), particularly internet-based cognitive behavioral therapy (iCBT), is a promising solution to these challenges. When paired with human support, iCBT has been shown to be as effective as face-to-face CBT [28–30]. Through packaged online lessons, iCBT consistently delivers core therapeutic techniques to patients. Implementing iCBT in aging service agencies shifts providers from direct treatment delivery to supportive roles—thereby reducing training demands and time commitments without compromising CBT fidelity. This makes iCBT a more feasible and scalable approach.

Despite their potential, iCBT programs face significant challenges with user engagement, particularly among older adults. Completion rates have been low, with an average of 17% in self-guided iCBT [31] and 65% in programs with human support [32]. For homebound older adults, engagement barriers are often exacerbated by usability challenges, including text-heavy,

Protocol: iCBT for Older Adults

academic content and interfaces that are not designed with older users in mind. For example, in our evaluation of an evidence-based iCBT program, only one in four homebound older adults completed the program, even with weekly in-home support [33]. These findings underscore the urgent need for iCBT programs tailored to older adults.

Although several iCBT programs have been designed to address older adults' needs, none are widely available to U.S. consumers [34–36]. To address this gap, we developed Empower@Home, an iCBT program explicitly tailored for older adults. Created in collaboration with key stakeholders, the program uses plain language, integrates text with visuals, incorporates age-relevant examples, and features a simplified user interface. Preliminary studies found *Empower@Home* to have superior usability compared to other iCBT programs. Feasibility trials also demonstrated exceptional program completion rates and significant reductions in depressive symptoms when participants were supported by lay coaches [37–40]. However, it remains unclear whether *Empower@Home* will be equally effective when integrated into real-world community aging services and supported by aging service providers.

Objective

The primary aim of this study is to evaluate the real-world effectiveness of Empower@Home, an iCBT program tailored to older adults with depression, when supported by aging service providers. A mixed-methods approach will be used, incorporating structured surveys and semi-structured interviews to examine intervention outcomes and underlying mechanisms. The secondary aim is to identify implementation barriers and facilitators through a qualitative process evaluation. This study builds on previous efficacy trials demonstrating the acceptability and effectiveness of Empower@Home. It takes a critical next step by transitioning the intervention from an academic setting into community-based aging services, thereby assessing its real-world impact and the feasibility of training aging service providers to act as coaches or human supporters for iCBT.

Intervention Model

The design of Empower@Home incorporates user-centered and community-participatory research principles [37]. The intervention was designed with input from homebound older adults, therapists, and over 30 aging service providers, undergoing multiple iterative refinements based on testing and feedback [37–39]. Empower@Home consists of nine sequenced online lessons, each lasting 20–25 minutes, featuring didactic content delivered through short videos, voice-over instructions, interactive exercises, and motivational quotes. These lessons are grounded in evidence-based CBT manuals [41–43] and focus on core CBT skills: "Doing Tools" for behavioral activation and problem-solving, "Thinking Tools" for cognitive restructuring, "Feeling Tools" for relaxation and mood monitoring, and "Communication Tools" for effective communication [41].

The content is specifically tailored to the unique needs of homebound older adults, incorporating age-relevant examples informed by stakeholder input and prior research [44–46]. Each lesson integrates a mini animated story centered on a homebound older woman, Jackie, facing common health challenges within the target demographic. According to persuasive design principles,

Protocol: iCBT for Older Adults

interventions that are interactive, engaging, and immersive are more effective in promoting behavioral change [45]. Entertainment education theories further suggest that narratives can reduce resistance to behavioral modifications by fostering emotional engagement and personal reflection [46]. Jackie's story encourages users to identify with the character, adopt story-aligned behaviors, and improve treatment engagement. Qualitative interviews with past participants indicated mixed identification with the storyline but suggested that it enhanced engagement [40].

Aligned with best practices for age-friendly digital design [47], the program features a streamlined interface with large buttons, text-labeled icons, high-contrast colors, and intuitive navigation. Online lessons are complemented by a large-print workbook containing session summaries, additional psychoeducational materials, wellness resources, and home practice instructions. Designed with older adults in mind, the workbook features large text, images of diverse older adults, and a visually appealing layout. Prior studies highlighted the workbook's importance, noting its familiar format, aesthetic appeal, and utility as a convenient reference for completing and revisiting exercises and lessons [40].

To address real-time support needs and technological challenges, we introduced "Empower Coaches," laypersons who provide weekly support calls to enhance program personalization, guided by the Efficiency Model of Support [37]. Our target demographic favors this human interaction over purely automated systems. Supported iCBT programs consistently demonstrate better engagement and clinical outcomes than those without support [30–32]. While many iCBT programs rely on therapists with specialized mental health qualifications, utilizing aging service providers as iCBT supporters offers an innovative and practical solution. This strategy leverages aging service providers' existing relationship with the target population and aligns with the staffing structures of aging service agencies, ensuring widespread and accessible support. Prior research has successfully trained aging service providers to administer psychotherapy and multi-component interventions directly [18–25], suggesting that training aging service providers as coaches is also likely feasible.

Our approach focuses on equipping service providers to support iCBT—a role that does not require extensive specialized training and is less time-intensive than acting as lead interventionists. Our previous efficacy studies showed that Empower@Home supported by students, such as undergraduate and graduate students in psychology, social work, or related fields, was acceptable and efficacious based on both qualitative and quantitative outcomes [38,40]. However, to date, no studies have examined the feasibility of training aging service providers such as case workers and community health workers to support iCBT or integrating iCBT into aging service settings. This study aims to fill that gap.

Our team developed a structured coaching guide based on experiences from pilot studies. The guide outlines coaching tasks, provides tips, and includes CBT teaching points aligned with session content. Coaches conduct weekly check-ins, which can be completed either over the phone or in person, allowing flexibility and greater efficiency. For example, coaches with scheduled home visits can incorporate coaching sessions into the visit without requiring additional scheduling. Based on prior research and our experience, coaching support typically includes providing motivation, encouragement, homework review, assistance with applying tools, and technical support. In line with the efficiency model of support [48], coaches are

Protocol: iCBT for Older Adults

encouraged to personalize their approach for each client—for instance, offering to complete part of the program alongside an unmotivated participant. The length of coaching sessions can be adapted to meet participants' needs and preferences, ranging from 5 minutes to an hour.

Coach training includes reviewing all Empower@Home online sessions (approximately 4–5 hours), supplemented with didactic videos introducing each session and its techniques. Coaches also attend a live, in-person training workshop lasting about 4 hours, where they observe demonstrations of coaching sessions and receive clarification on operating logistics. The total training requirement is approximately 9–10 hours per coach. Biweekly coach support meetings are offered throughout the study to provide ongoing support. While previous studies limited the intervention period to 10 weeks, this project extends it to 12 weeks to accommodate the frailty of the target population, who may face additional barriers such as hospitalizations that could delay program completion.

Methods

Preliminary Study

The procedures outlined in this protocol, including recruitment, assessments, coaching call logistics, and device management, have been tested and refined through feasibility and efficacy studies conducted in 2022–2023. A pilot randomized controlled trial (RCT) with 70 participants, supported by trained research assistants, demonstrated that the program was significantly more effective in reducing depressive symptoms compared to biweekly friendly calls, with a medium between-group effect size (Cohen's $d = 0.7$) [38]. Program adherence was high, with approximately 90% of participants completing all sessions within 10 weeks [39]. Participants reported high satisfaction with the program, and themes from qualitative interviews supported the quantitative findings, further highlighting the program's positive effects [40]. However, positive findings from small-scale pilot studies may not always translate into comparable outcomes in real-world settings or larger studies. This trial evaluates whether these promising results can be replicated in a community-based, real-world implementation context.

Study Setting

The University of Michigan will be the primary study site, in collaboration with community agencies serving older adults. All participants will be 60 years old or over and are expected to be predominantly low-income. Participants will complete the intervention in their own homes, with all assessments and technological support provided remotely by phone.

Eligibility Criteria

To participate in the trial, individuals must be at least 60 years old, eligible to receive services at the participating community agency, and have mild or greater depressive symptoms, defined as a score of ≥ 8 on the Patient Health Questionnaire (PHQ-9). Participants will be excluded if they meet any of the following criteria: (a) Probable cognitive impairment, indicated by a score of ≥ 12 on the Blessed Orientation-Memory-Concentration Test [49]; (b) Non-English speakers; (c) Moderate to high suicide risk, as assessed by the Columbia-Suicide Severity Rating Scale [50];

Protocol: iCBT for Older Adults

(d) Terminal illness with a life expectancy of fewer than six months or unstable physical health; (e) Visual impairment that prevents the use of a screen, even with corrective measures (self-reported); (f) Current or recent substance use, as indicated by a score of ≥ 2 on the CAGE-AID Substance Abuse Screening Tool [51]; (g) Diagnosis of a psychotic disorder (self-reported); (h) Active psychotherapy (more than once a month) or initiation of new therapy within the last three months. The study will not exclude participants based on race or gender. While completing the intervention requires a tablet and internet access, participants are not required to have their own. Those without a tablet or internet access will be provided with both at no cost for the duration of the active intervention period.

Recruitment and Compensation

Participants will be recruited on a rolling basis, primarily through referrals from participating community agencies. Two of these agencies are Area Agencies on Aging (AAAs), which may refer existing clients enrolled in their programs, such as case management, caregiver services, or Meals on Wheels. Additionally, AAAs may refer new clients or individuals seeking information from the agency, as providing information is a key role of AAAs.

A 20-minute online training covering eligibility criteria and referral procedures is provided to participating agencies to support participant recruitment. Agencies are also supplied with a pre-screening form that includes the PHQ-2 and additional brief screening items designed by the research team to improve referral accuracy. Based on prior experience, approximately half of the individuals passing the pre-screening will qualify for the study. All pre-screening and referral forms are hosted online and optimized for efficiency.

After receiving referrals, study personnel will contact potential participants by phone to complete a more detailed eligibility screening. This initial screening typically takes 20 minutes but may take longer if a full suicidal ideation assessment is needed. Participants who agree to participate will provide recorded verbal consent, while those who decline will be referred back to their case managers or agency when feasible. Additionally, participants can self-refer for eligibility screening through advertisements on social media platforms and by word of mouth.

Qualified participants will be compensated up to \$140, including \$30 for each survey assessment and \$20 for the qualitative interview. Compensation is prorated based on completed assessments (e.g., participants completing only the baseline assessment will receive \$30).

Study Design and Randomization

Eligible and consented participants will be randomly assigned to either the treatment or control groups. Both groups will follow the same assessment schedule, including a comprehensive baseline assessment, a post-intervention/control assessment at 12 weeks, and follow-up assessments at 24 and 36 weeks. Both groups will continue to receive usual care from their referring agency. Participants in the control group will receive the intervention after completing their final follow-up assessment.


Block randomization, stratified by participating community agencies, will be used to assign

Protocol: iCBT for Older Adults

participants. Each site has nine blocks of 10 participants. The randomization sequence was created by the study statistician (JJ), and sealed envelopes were prepared by a research assistant who was not involved in the randomization process. The lead project coordinator will conduct the randomization using these sealed envelopes.

Due to the nature of the intervention and study design, blinding participants and evaluators to group assignments is not feasible. However, assessors will remain blinded to group allocation whenever possible.

Figure 1 Participant Flow



Protocol: iCBT for Older Adults

Control Condition

The attention control condition provides participants with a comparable amount of phone contact as the intervention group, including 15–25-minute companionship calls (“friendly calls”) every other week. These calls are conducted by members of the research team and focus on providing social interaction and support without introducing CBT-related content or structured therapeutic interventions. To ensure safety and align with the biweekly PHQ-9 assessments conducted in the intervention group as a part of the online program, participants in the control group will complete a telephone-administered PHQ-9 assessment every two weeks as a part of the friendly calls. Additionally, participants in the control group will continue receiving regular social services from the participating agency and will receive a psychoeducational handout containing information about depression and a list of community resources for further support.

Safety Monitoring and Discontinuation

Participant safety is a top priority in this study, with procedures in place to monitor and address any significant clinical changes. Biweekly assessments of depressive symptoms using the PHQ-9 serve as both clinical outcome measures and safety checks. A clinically significant worsening of symptoms, defined as a 5-point or greater increase in the PHQ-9 from baseline, triggers a multi-step review process. This process includes automated alerts in our electronic data-capturing system and in-depth risk assessments by the project coordinator, a licensed clinical social worker. While such occurrences are rare—less than 3% of participants experienced a 5+ point increase during pilot studies—persistent or worsening symptoms over four weeks may lead to discontinuation. Similarly, any new or increased endorsement of suicidal ideation on the PHQ-9’s 9th item prompts immediate follow-up using the Columbia-Suicide Severity Rating Scale (C-SSRS), following an established protocol to ensure participant safety.

Participants may also be withdrawn from the study if their physical health conditions worsen unrelated to the intervention, compromising their ability to participate safely (e.g., repeated hospitalizations). Such decisions are informed by input from the participant, agency coaches, and family members. In all cases of withdrawal, the study team collaborates with partnering community agencies to connect participants to appropriate treatment, including pharmacotherapy and locally available psychosocial resources. Final discontinuation decisions will be made by the core research team involved in the day-to-day management of human subjects, including the PI and the three project staff.

Data Management and Storage

All study data collected through surveys or structured observations will be entered directly by project staff into REDCap [41]. Agency coaches will also complete a structured coaching form hosted on REDCap. During the active intervention, participants will complete the PHQ-9 every other week as part of the online therapy lessons. These assessment data will be securely stored in a HIPAA-compliant database hosted by the University of Michigan. Recordings of qualitative interviews will be stored on Dropbox, in compliance with the University of Michigan’s safe

Protocol: iCBT for Older Adults

computing guidelines for handling sensitive research data. Additionally, meeting minutes and other program administrative records, which do not contain participant identifying information, will be stored on Google Drive.

Measures

Primary clinical outcomes

All key outcome measures are summarized in Table 1. The primary outcome is changes in depressive symptoms, assessed using the PHQ-9, a widely used screening tool in primary care, community agencies, and iCBT studies [52]. Each participant will complete the PHQ-9 up to a total of nine times: at screening (T0), post-intervention at 12 weeks (T2), and follow-up assessments at 24 weeks (T3) and 36 weeks (T4), along with up to five biweekly assessments between baseline (T1) and post-intervention (T2). For participants in the intervention group, PHQ-9 assessments will be embedded within the online program and completed during sessions 1, 3, 5, 7, and 9. Control group participants will complete these assessments via telephone at corresponding intervals. Additionally, the 7-item Hamilton Depression Rating Scale (HAMD-7) will be used as a secondary measure to evaluate the intervention's clinical effectiveness. The HAMD-7, an abbreviated version of the HAMD-17, is shown to be equally effective in assessing remission in patients with major depressive disorder [53]. It was chosen to reduce participant burden and to enhance the feasibility of telephone-based administration.

Table 1 Key Study measures and time points

Measurement Domain	Instrument	Baseline (T1)	Bi-weekly	Post (T2)	Follow-Up 1 (T3)	Follow-Up 2 (T4)
Primary Clinical Outcomes						
Depression	PHQ-9	*	✓	✓	✓	✓
Depression	HAMD-7	✓		✓	✓	✓
Secondary Clinical Outcomes						
Anxiety	GAD-7	✓		✓	✓	✓
Social Isolation	DSSI-10	✓		✓	✓	✓
Health-related Quality of Life	EQ-5D-5L	✓		✓	✓	✓
Disability Burden	WHODAS 2.0	✓		✓	✓	✓
Pain	PEG	✓		✓	✓	✓
Treatment Mechanisms						
CBT related factors						
CBT Skills Acquisition	Modified CBTSQ	✓		✓		
Behavioral Activation	BADS-SF	✓		✓		

Protocol: iCBT for Older Adults

Dysfunctional Attitudes	DAS-SF2	✓		✓		
Social Engagement	SEAQ	✓		✓		
Psychological Flexibility	AAQ-II	✓		✓		
<i>Entertainment or Engagement-Related Factors</i>						
Digital Alliance	D-WAI			✓		
Narrative Engagement	NES			✓		
Character Identification	EDI			✓		
<i>Coach-related factors</i>						
Coaching Fidelity	Study-specific measure			✓		
<i>Self-determination</i>						
Autonomy	BNSQ-S	✓		✓		
<i>Feasibility & Acceptability</i>						
Treatment Adherence	Session completion			✓		
Treatment Acceptability	TEI			✓		

*Note. The PHQ-9 was administered at the screening visit (T0) rather than at baseline (T1). This decision was made because the PHQ-9 assesses symptoms over the previous two weeks, and T0 and T1 assessments are typically scheduled within two weeks of each other. Participants generally complete their first bi-weekly PHQ-9 assessment within two weeks of the baseline assessment. The PHQ-9 is assessed at T0 but not repeated at T1 to minimize redundancy and reduce participant burden.

Secondary Clinical Outcomes

The secondary clinical outcomes, listed in Table 1, include changes in anxiety, social isolation, quality of life, disability burden, and pain severity, assessed at four time points (T1-T4). These outcomes were chosen for their significant impact on the quality of life of older adults with chronic illnesses and disabilities, aligning with prior research on geriatric depression treatments [20,54].

Anxiety will be assessed with General Anxiety Disorder-7 (GAD-7), which has been validated in older adults [33,55]. Participants answer 7 questions probing the frequency of their anxiety symptoms rated on a 4-point Likert scale ranging from “not at all” to “nearly every day.” The GAD-7 has demonstrated 89% sensitivity and 82% specificity in detecting generalized anxiety disorders in primary care patients [37].

Social isolation will be assessed using the Duke Social Support Index (DSSI) 10-item scale, which evaluates social interaction and satisfaction with social support in older adults [56].

Quality of Life will be measured using the EuroQol 5-Dimension 5-level (EQ-5D-5L) scale [57].

Protocol: iCBT for Older Adults

Participants rate their ability to perform tasks in five domains—mobility, self-care, usual activities, pain, and anxiety/depression—on a 5-point Likert scale ranging from "no problem" to "extreme problem or unable." Responses will be converted into utility values using a U.S. population-specific algorithm [58].

Disability burden will be assessed using the WHO Disability Assessment Schedule 2.0 (WHODAS 2.0) 12-item survey, which examines difficulty performing activities across six domains: cognition, mobility, self-care, getting along with people, life activities, and societal participation. Responses range from "none" to "extreme or cannot do" [59].

Pain severity will be evaluated with the PEG Scale, a 3-item measure that assesses pain intensity and its interference with enjoyment of life and general activity. Each item is rated on a scale from 0 to 10 [60].

Treatment Mechanisms

We hypothesize that the intervention operates through multiple mechanisms grounded in CBT, entertainment persuasion, and self-determination theories. CBT principles suggest that depression stems from maladaptive thoughts and behaviors; thus, improved CBT skills, behavioral activation, and reduced dysfunctional attitudes are expected to mediate treatment effects [61]. The animated Jackie storyline aims to enhance engagement, fostering a connection with the narrative and program [46]. Additionally, the coach-client rapport, aligned with the common factors theory, is hypothesized to mediate outcomes through the therapeutic bond and satisfaction with coaching [62]. Self-determination theory highlights the importance of autonomy, competence, and relatedness in fostering motivation and well-being [63].

CBT related factors

Jacob et al., [64] developed the 16-item Cognitive-Behavioral Therapy Skills Questionnaire (CBTSQ) to assess the use of CBT skills such as cognitive restructuring and behavioral activation. We modified the scale to 14 items by removing two: "Identify stressors that led me into treatment," as participants were not recruited from clinical settings, and "Catch myself when I jump to conclusions," as the pre-program assessment occurs before participants are introduced to this concept, enhancing reliability. Item wording for the remaining 14 items was simplified for readability, e.g., "Identify risk factors for relapse" was revised to "Identify warning signs that trigger bad mood, depression, or anxiety." Participants rate their frequency of activities like organizing plans, motivating themselves, and socializing on a 5-point Likert scale from "never" to "always."

The 9-item Behavioral Activation for Depression Scale–Short Form (BADS-SF) evaluates activation and avoidance behaviors consistent with Behavioral Activation theory [65]. Each item is rated on a 7-point scale from 0 ("not at all") to 6 ("completely"), with higher scores reflecting greater activation.

The 9-item Dysfunctional Attitudes Scale–Short Form 2 (DAS-SF2) is a self-report measure for assessing the presence and intensity of rigid or maladaptive beliefs [66]. We adapted the original

Protocol: iCBT for Older Adults

4-point Likert scale ("totally agree," "agree," "disagree," "totally disagree") to a 5-point Likert scale ranging from "strongly disagree" to "strongly agree" to be consistent with the other measures in the study.

The 10-item Social Engagement and Activities Questionnaire (SEAQ) was developed to assess social engagement in homebound older adults, accounting for physical and functional limitations. Participants report the frequency of activities such as socializing, volunteering, attending religious services, and participating in recreational or educational events over the past month. Responses are scored on a 6-point scale (0–5) for each item [67].

Entertainment or Engagement-Related Factors

The Digital Working Alliance Inventory (D-WAI) is a 6-item measure that asks participants to rank their experience using a digital mental health intervention [68]. Responses are scored using a 5-point Likert scale from "strongly agree" to "strongly disagree." The word "app" from the original D-WAI scale was modified to "program" to reflect that the intervention is accessed online rather than through a mobile app.

The 12-item Narrative Engagement Scale (NES) assesses the audience's mental models, particularly their enjoyment and understanding of a story, using a 5-point Likert scale ranging from "strongly agree" to "strongly disagree" [69]. The scale includes four domains, each measured by three items: narrative understanding, attentional focus, narrative presence, and emotional engagement. For this study, we retained all three items from the emotional engagement domain but selected only one item each from the narrative understanding, attentional focus, and narrative presence domains.

The Identification with Characters Scale (EDI) consists of 14 items that help identify different levels of relatedness and connectedness the viewer has to a film [70]. We adapted the original 5-point scale ("1 = not at all" to "5 = very much") to a 5-point scale ranging from "strongly disagree" to "strongly agree" for consistency with other questions. Additionally, "characters" was replaced with Jackie's name (e.g., "I felt like I was similar to Jackie").

Coach-related factors

Coaching fidelity (client report) will be assessed using a 9-item scale developed specifically for this study. Some items were adapted from the Therapist's Interpersonal Style Scale (ATIS) [71] to align with the study context, emphasizing aspects of coaching that reflect therapeutic alliance and its potential influence on treatment outcomes. Example items include: "The feedback I receive from my coach is helpful," "My coach is trustworthy," and "My coach makes the program more personalized and relevant to my situation and needs." Participants will rate their agreement with each statement on a 5-point Likert scale ranging from "strongly agree" to "strongly disagree."

Self-determination

The Basic Needs Satisfaction in General Scale (BNSQ-S) is a 16-item questionnaire designed to

Protocol: iCBT for Older Adults

assess satisfaction with three fundamental psychological needs: autonomy, competence, and relatedness. Autonomy pertains to an individual's desire to perceive their actions and subsequent consequences as self-determined, rather than being subject to external influences or control [72]. Competence refers to the inherent desire to experience a sense of efficacy and proficiency in executing tasks of varying complexities [73]. Relatedness refers to the inherent human desire for social connection, support, and caring from others [73]. According to the self-determination theory, all three needs must be fulfilled to achieve psychological well-being [72]. Consistent with the responses of other measures used in this study, participants respond on a 5-point scale ranging from "strongly agree" to "strongly disagree." [74].

Acceptability will be assessed by the number of sessions participants completed (i.e., adherence) and the Modified Treatment Evaluation Inventory (TEI) developed explicitly for evaluating geriatric depression treatment [75]. The TEI has 11 items, including eight positively worded items and three negatively worded items. We modified the TEI by rephrasing the original TEI items from questions to statements and reducing the responses from a 7-point Likert scale to a 5-point Likert scale to ease respondents' burden. The modified TEI yields total scores ranging from 11 to 55, where a score of ≥ 32 suggests positive attitudes toward the treatment.

Implementation fidelity monitoring

Implementation fidelity is essential for evaluating interventions delivered in real-world settings. While the online program ensures consistent delivery of core CBT concepts, coaching support naturally varies due to its personalized nature. We employed three methods to assess fidelity in this context: coach self-reports, in vivo observations of coaching calls, and fidelity check-in calls with clients.

Coach self-reports capture the coach's perspective on each session. Coaches document session duration, delivery mode (phone, virtual, or in-person), and types of support provided, selecting from a checklist including application, education, motivation, personalization, and technology assistance. Coaches also respond to close-ended questions assessing adherence to session tasks outlined in the coaching guide, such as "Did you review in-session exercises as outlined in the coaching guide?" with response options of All, Most, Some, or None.

In vivo observations involve trained research team members evaluating at least one coaching session per client, selected randomly using a predetermined random sequence. Observers join sessions via three-way calls to assess fidelity in real-time. A structured rating guide, developed specifically for this study, includes items adapted from the literature and validated scales such as the UK Alcohol Treatment Trial Process Rating Scale (UKATT PRS) [76] and the Therapist's Interpersonal Style Scale (ATIS) [71]. The guide comprises items across three domains: session management, adherence to the coaching guide, and coaching style. Key items under session management and adherence to the coaching guide include collaborative agenda-setting, reviewing exercises, checking goal progress, and providing personalized feedback. Coaching style is rated based on factors such as engagement, empathy, reflective listening, and collaboration. Each item is scored on a 4-point scale from Excellent to Poor. The instrument includes detailed instructions for rating each item. To ensure reliability, two research staff independently rated coaching call recordings from pilot studies, compared their ratings, and

Protocol: iCBT for Older Adults

resolved discrepancies to refine the scale and its instructions for consistency.

Qualitative Interviews

After completing the intervention, approximately half of the intervention group will take part in a semi-structured qualitative interview lasting about 30 minutes. These interviews will explore several domains, including participants' overall experience with the program, engagement barriers and facilitators, and their experience with the coaching component.

Additionally, all Empower Coaches, their immediate supervisors (who also serve as site principal investigators for IRB purposes), and other relevant agency staff involved in implementation will be invited to participate in qualitative interviews during the final year of data collection. These interviews will gather insights into how the intervention functioned, what modifications were or should be made, and how well it integrates into routine provider workflows. The updated Consolidated Framework for Implementation Research (CFIR) [77] will guide the design of qualitative interview protocols and data analysis.

Sample Size

All parameters for the power analysis were based on the pilot RCT conducted in 2023, with an original sample size of 70, with 35 treatment and 35 control participants. At the time of this analysis, a total of 440 data points across the 7 time points had been collected. Time 7 data had been collected from 63 participants, 33 in the treatment and 30 in the control group. A linear mixed model using lme from R package lme4 showed a treatment by time effect of -0.69, $p < .001$. This pilot data was used in the R package simr with the effect size conservatively set to approximately one-half of that found in pilot study, -0.35. With a sample size of 192, there was a power of 0.91 to detect the effect of -0.35 for the treatment by time parameter. Assuming 25% attrition, we will aim to recruit 256 participants.

Analytic Plan

Primary analyses

The primary analyses will use an intention-to-treat approach, including all individuals who completed the baseline interview and were randomized. If sufficient variability in participation or dropout is observed, secondary analyses will incorporate measures of intervention dosage (i.e., adherence) as covariates to account for differences in engagement with the intervention.

Linear mixed-effects models will be used to examine differences between the treatment and control groups over time. All outcome variables will be assessed for normality, and transformations will be applied if needed. Parameter estimates for the interaction between group membership and time will determine whether the treatment group demonstrates significantly different changes over time compared to the control group. Various covariance structures will be evaluated to identify the best-fitting model, and non-linear time functions (e.g., quadratic) will also be explored.

Protocol: iCBT for Older Adults

Mediation analysis

Latent growth curve modeling will be used to assess whether proposed mediators—CBT-related, engagement-related, and coach-related factors—explain the primary treatment effect [78]. These models will be estimated using Mplus software. According to Cheong [79], latent growth curve models with five measurement points show minimal bias and sufficient power to detect mediation effects, especially when the R^2 (the proportion of variance in mediator and outcome variables explained by growth factors) is moderate or high. In this study, seven measurement points will be used in mediation analysis (T1, T2, and 5 in-app assessments), making the estimates conservative as power increases and bias decreases with additional measurements. Based on guidance from Cheong [79], assuming a medium effect size and moderate R^2 , the sample size of 192 will yield power of 0.8 to detect the mediation effect, indicating adequate statistical power for the analysis.

Qualitative analysis

Third-party transcription services will be used to transcribe recorded qualitative interviews. Coding and theme extraction will follow Saldaña's methodology for qualitative analysis [80]. In the first coding cycle, a combination of a priori codes from preliminary studies and inductively developed new codes will be used to analyze transcripts. Two trained research staff will independently review a random subset of transcripts to ensure coding consistency. This process will continue until a satisfactory level of agreement is reached, at which point the finalized codebook will be applied to the remaining transcripts. During the second coding cycle, patterns and relationships among initial codes will be identified, with codes being grouped into broader themes or concepts through an iterative process. Data analysis will be conducted using Dedoose, which supports essential qualitative analysis tasks such as coding, creating memos, and diagramming.

Results

Funding for this project was awarded in July 2024. Institutional Review Board approval was obtained in August 2024. Enrollment began in October 2024 and is anticipated to be completed in April 2028. The trial is registered at ClinicalTrials.gov, NCT06584422.

Discussion

To our knowledge, this study represents the first Hybrid Type 1 implementation-effectiveness randomized controlled trial (RCT) of an aging service provider-supported internet-based cognitive behavioral therapy (iCBT) intervention for low-income homebound older adults with depression. We hypothesize that the Empower@Home intervention will reduce depressive symptoms and improve psychosocial functioning and health-related quality of life when supported by aging service providers.

Despite the availability of numerous iCBT programs, very few are explicitly designed for older adults, a population more sensitive to usability challenges. Furthermore, the real-world

Protocol: iCBT for Older Adults

effectiveness of these programs has often been limited by difficulties in achieving consistent and meaningful user engagement [81]. This study addresses these gaps by tailoring an iCBT program specifically for older adults and embedding it within community-based aging services, supported by trained aging service providers.

This study makes two significant contributions to the field. First, it provides crucial evidence on integrating iCBT with training community providers, such as aging service staff, to offer human support. This approach tests the feasibility and effectiveness of using providers already embedded in the community to deliver iCBT in real-world settings. While our prior efficacy trials demonstrated that layperson-supported iCBT can succeed in highly controlled academic environments, this study explores whether those results can be replicated in community settings. It will also address questions about treatment fidelity—whether aging service providers can consistently follow the intervention protocol—and whether their support will yield effects comparable to those achieved by student coaches in our previous studies. Interestingly, the intergenerational aspect of using student coaches in past trials may have played a unique role that indigenous providers with existing relationships with their clients might not replicate. By testing these variables, this study lays the groundwork for future large-scale implementation and scalability efforts.

Second, our previous efficacy trials showed exceptional program engagement, with 90% of participants completing the program—a stark contrast to the average completion rates reported in meta-analyses. For example, one meta-analysis of 40 iCBT studies found that 57% of participants dropped out [82]. More recent systematic reviews have shown that completion rates average 17% for self-guided digital mental health interventions [83] and 65% for therapist-supported interventions [32]. Additionally, usage rates often drop sharply when these interventions are implemented outside controlled study conditions [84,85]. While our earlier qualitative studies offered insights into high engagement, many factors contributing to this success remain unknown. In this study, we will examine engagement-related, coaching-related, and CBT-related mechanisms to better understand treatment pathways. These findings could provide broader insights into overcoming engagement challenges in digital mental health interventions.

The rapid evolution of digital mental health technologies, particularly with advancements in artificial intelligence, underscores the importance of addressing the digital divide. Without careful consideration, these innovations risk further marginalizing underserved populations, including low-income older adults. This study demonstrates how technology can be leveraged alongside existing service delivery infrastructures to address unmet mental health needs. It provides a critical data point on effectively integrating digital mental health interventions into community settings effectively.

Finally, this study is particularly timely given the long-term mental health consequences of the COVID-19 pandemic [86], which have disproportionately affected vulnerable populations such as homebound older adults. By addressing engagement and implementation challenges, this research contributes to the ongoing effort to close mental health disparities and ensure equitable access to effective treatments.

Protocol: iCBT for Older Adults

Strengths and Limitations

This study has several notable strengths. The intervention was iteratively developed, incorporating patient and provider feedback, and was tested using a rigorous randomized design. It employs laypersons as interventionists and is implemented in real-world community settings, enhancing its practical relevance. The study includes a 36-week follow-up period to assess longer-term effects and focuses on an underserved, hard-to-reach population with significant unmet mental health needs. [81]. However, the study also has limitations. While it is fully powered at the individual participant level, including only three community agencies limits the ability to thoroughly analyze differences at the agency or coach levels. Future studies could address these limitations with a greater focus on implementation.

Conclusion

This study offers some of the first evidence on integrating digital mental health interventions into aging services. If successfully implemented, this innovative approach could help bridge the digital divide and address mental health disparities among homebound older adults. Additionally, the findings have broader relevance for other digital mental health interventions and populations with physical disabilities or access barriers who continue to face significant barriers to mental health care.

Acknowledgments

Research reported in this publication was supported by the National Institute of Mental Health of the National Institutes of Health under award number R01MH137064. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflicts of Interest

None declared.

Abbreviations

MMSE: Mini Mental State Exam
BOMC: Blessed Memory and Orientation Scale
PHQ-9: Patient Health Questionnaire – 9
HAM-D-7: Hamilton Depression Rating Scale – 7
CAGE: CAGE Substance Use Screener
CSSRS: Columbia-Suicide Severity Rating Scale
GAD-7: GAD-7, Generalized Anxiety Disorder – 7
DSSI-10: Duke Social Support Index – 10
EQ-5D: EuroQol 5-Dimension 5-Level
WHODAS 2.0: WHO Disability Assessment Schedule 2.0
PEG: PEG Scale (Pain, Enjoyment, General Activity)

Protocol: iCBT for Older Adults

TEI: Treatment Evaluation Inventory

AAQ-II: Acceptance and Action Questionnaire

MARS-10: Medication Adherence Report Scale – 10

BADS-SF: Behavioral Activation for Depression Scale Short Form

DAS-SF2: Dysfunctional Attitudes Scale–Short Form 2

SEAQ: Social Engagement and Activities Questionnaire

CBTSQ: Cognitive-Behavioral Therapy Skills Questionnaire

D-WAI: Digital Working Alliance Inventory

NES: Narrative Engagement Scale

EDI: Identification with Characters Scale

Coaching Fidelity: Study-specific measure

BNSQ-S: Basic Needs Satisfaction in General Scale

MDPQ: Mobile Device Proficiency Questionnaire – mobile device basics subscale

Protocol: iCBT for Older Adults

References

1. Qiu WQ, Dean M, Liu T, et al. Physical and mental health of homebound older adults: An overlooked population. *J Am Geriatr Soc*. 2010;58(12):2423-2428. doi:10.1111/j.1532-5415.2010.03161.x.
2. Choi NG, McDougall GJ. Comparison of depressive symptoms between homebound older adults and ambulatory older adults. *Aging Ment Health*. 2007;11(3):310-322. doi:10.1080/13607860600844614.
3. Xiang X, Leggett A, Himle JA, Kales HC. Major Depression and Subthreshold Depression among Older Adults Receiving Home Care. *Am J Geriatr Psychiatry*. 2018;26(9):939-949. doi:10.1016/j.jagp.2018.05.001.
4. Bruce ML, McAvay GJ, Raue PJ, et al. Major depression in elderly home health care patients. *Am J Psychiatry*. 2002;159(8):1367-1374. doi:10.1176/appi.ajp.159.8.1367
5. Reynolds CF 3rd, Kupfer DJ. Depression and aging: a look to the future. *Psychiatr Serv*. 1999;50(9):1167-1172. doi:10.1176/ps.50.9.1167.
6. Xiang X, An R, Oh H. The Bidirectional Relationship Between Depressive Symptoms and Homebound Status Among Older Adults. *J Gerontol B Psychol Sci Soc Sci*. 2020;75(2):357-366. doi:10.1093/geronb/gbx180.
7. Alexopoulos GS. Depression in the elderly. *Lancet*. 2005;365(9475):1961-1970. doi:10.1016/S0140-6736(05)66665-2.
8. Meng R, Yu C, Liu N, et al. Association of Depression With All-Cause and Cardiovascular Disease Mortality Among Adults in China. *JAMA Netw Open*. 2020;3(2):e1921043. doi:10.1001/jamanetworkopen.2019.21043.
9. Xiang X, Danilovich MK, Tomasino KN, Jordan N. Depression prevalence and treatment among older home health services users in the United States. *Arch Gerontol Geriatr*. 2018;75:151-157. doi:10.1016/j.archger.2017.12.005.
10. Choi NG. The integration of social and psychologic services to improve low-income homebound older adults' access to depression treatment. *Fam Community Health*. 2009;32(1 Suppl):S27-S35. doi:10.1097/01.FCH.0000342837.97982.36.
11. Ornstein KA, Leff B, Covinsky KE, et al. Epidemiology of the Homebound Population in the United States [published correction appears in *JAMA Intern Med*. 2015 Aug;175(8):1426. doi: 10.1001/jamainternmed.2015.4022]. *JAMA Intern Med*. 2015;175(7):1180-1186. doi:10.1001/jamainternmed.2015.1849.
12. Alexopoulos GS. Mechanisms and treatment of late-life depression. *Transl Psychiatry*. 2019;9(1):188. doi:10.1038/s41398-019-0514-6.
13. Reynolds CF 3rd, Lenze E, Mulsant BH. Assessment and treatment of major depression in older adults. *Handb Clin Neurol*. 2019;167:429-435. doi:10.1016/B978-0-12-804766-8.00023-6.
14. Blazer DG. Depression in Late Life: Review and Commentary. *Focus*. 2009;7(1):118-136. doi:10.1176/foc.7.1.foc118.

Protocol: iCBT for Older Adults

15. Pepin R, Segal DL, Coolidge FL. Intrinsic and extrinsic barriers to mental health care among community-dwelling younger and older adults. *Aging Ment Health*. 2009;13(5):769-777. doi:10.1080/13607860902918231.
16. Kunik ME, Mills WL, Amspoker AB, et al. Expanding the geriatric mental health workforce through utilization of non-licensed providers. *Aging Ment Health*. 2017;21(9):954-960. doi:10.1080/13607863.2016.1186150.
17. Lavingia R, Jones K, Asghar-Ali AA. A Systematic Review of Barriers Faced by Older Adults in Seeking and Accessing Mental Health Care. *J Psychiatr Pract*. 2020;26(5):367-382. doi:10.1097/PRA.0000000000000491.
18. Bruce ML, Raue PJ, Reilly CF, et al. Clinical effectiveness of integrating depression care management into medicare home health: the Depression CAREPATH Randomized trial. *JAMA Intern Med*. 2015;175(1):55-64. doi:10.1001/jamainternmed.2014.5835.
19. Choi NG, Wilson NL, Sirrianni L, Marinucci ML, Hegel MT. Acceptance of home-based telehealth problem-solving therapy for depressed, low-income homebound older adults: Qualitative interviews with the participants and aging-service case managers. *Gerontologist*. 2014;54(4):704-713. doi:10.1093/geront/gnt083.
20. Choi NG, Caamano J, Vences K, Marti CN, Kunik ME. Acceptability and effects of tele-delivered behavioral activation for depression in low-income homebound older adults: in their own words. *Aging Ment Health*. 2021;25(10):1803-1810. doi:10.1080/13607863.2020.1783516.
21. Ciechanowski P, Wagner E, Schmalting K, et al. Community-Integrated Home-Based Depression Treatment in Older Adults. *JAMA*. 2004;291(13):1569. doi:10.1001/jama.291.13.1569.
22. Quijano LM, Stanley M a., Petersen NJ, et al. Healthy IDEAS: A Depression Intervention Delivered by Community-Based Case Managers Serving Older Adults. *J Appl Gerontol*. 2007;26(2):139-156. doi:10.1177/0733464807299354.
23. Gitlin LN, Harris LF, McCoy MC, Hess E, Hauck WW. Delivery Characteristics, Acceptability, and Depression Outcomes of a Home-based Depression Intervention for Older African Americans: The Get Busy Get Better Program. *The Gerontologist*. 2016;56(5):956-965. doi:10.1093/geront/gnv117.
24. Szanton SL, Thorpe RJ, Gitlin LN. Beat the Blues Decreases Depression in Financially Strained Older African-American Adults. *Am J Geriatr Psychiatry*. 2014;22(7):692-697. doi:10.1016/j.jagp.2013.05.008.
25. Smith ML, Steinman LE, Montoya CN, Thompson M, Zhong L, Merianos AL. Effectiveness of the Program to Encourage Active, Rewarding Lives (PEARLS) to reduce depression: a multi-state evaluation. *Front Public Health*. 2023;11:1169257. doi:10.3389/fpubh.2023.1169257.
26. Choi NG, Marti CN, Wilson NL, et al. Effect of Telehealth Treatment by Lay Counselors vs by Clinicians on Depressive Symptoms Among Older Adults Who Are Homebound: A Randomized Clinical Trial. *JAMA Netw Open*. 2020;3(8):e2015648. doi:10.1001/jamanetworkopen.2020.15648.
27. Chen GJ, Kunik ME, Marti CN, Choi NG. Cost-effectiveness of Tele-delivered behavioral activation by Lay counselors for homebound older adults with depression. *BMC Psychiatry*. 2022;22(1):648. doi:10.1186/s12888-022-04272-9.
28. Andersson G, Waara J, Jonsson U, Malmaeus F, Carlbring P, Ost LG. Internet-based self-help versus one-session exposure in the treatment of spider phobia: a randomized controlled trial.

Protocol: iCBT for Older Adults

- Cogn Behav Ther. 2009;38(2):114-120. doi:10.1080/16506070902931326.
29. Baumeister H, Reichler L, Munzinger M, Lin J. The impact of guidance on Internet-based mental health interventions - A systematic review. *Internet Interv.* 2014;1(4):205-215. doi:10.1016/j.invent.2014.08.003.
 30. Carlbring P, Andersson G, Cuijpers P, Riper H, Hedman-Lagerlöf E. Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: an updated systematic review and meta-analysis. *Cogn Behav Ther.* 2018;47(1):1-18. doi:10.1080/16506073.2017.1401115.
 31. Karyotaki E, Riper H, Twisk J, et al. Efficacy of Self-guided Internet-Based Cognitive Behavioral Therapy in the Treatment of Depressive Symptoms: A Meta-analysis of Individual Participant Data. *JAMA Psychiatry.* 2017;74(4):351. doi:10.1001/jamapsychiatry.2017.0044.
 32. van Ballegooijen W, Cuijpers P, van Straten A, et al. Adherence to Internet-based and face-to-face cognitive behavioural therapy for depression: a meta-analysis. *PLoS One.* 2014;9(7):e100674. doi:10.1371/journal.pone.0100674.
 33. Xiang X, Sun Y, Smith S, Lai PHL, Himle J. Internet-Based Cognitive Behavioral Therapy for Depression: A Feasibility Study for Home Care Older Adults. *Res Soc Work Pract.* 2020;30(7):791-801. doi:10.1177/1049731520927783.
 34. Dear BF, Zou JB, Ali S, et al. Clinical and cost-effectiveness of therapist-guided internet-delivered cognitive behavior therapy for older adults with symptoms of anxiety: a randomized controlled trial. *Behav Ther.* 2015;46(2):206-217. doi:10.1016/j.beth.2014.09.007.
 35. Titov N, Dear BF, Ali S, et al. Clinical and cost-effectiveness of therapist-guided internet-delivered cognitive behavior therapy for older adults with symptoms of depression: a randomized controlled trial. *Behav Ther.* 2015;46(2):193-205. doi:10.1016/j.beth.2014.09.008.
 36. Tomasino KN, Lattie EG, Ho J, Palac HL, Kaiser SM, Mohr DC. Harnessing Peer Support in an Online Intervention for Older Adults with Depression. *Am J Geriatr Psychiatry.* 2017;25(10):1109-1119. doi:10.1016/j.jagp.2017.04.015.
 37. Xiang X, Kayser J, Ash S, et al. Web-Based Cognitive Behavioral Therapy for Depression Among Homebound Older Adults: Development and Usability Study. *JMIR Aging.* 2023;6:e47691. doi:10.2196/47691.
 38. Xiang X, Kayser J, Turner S, Ash S, Himle JA. Layperson-Supported, Web-Delivered Cognitive Behavioral Therapy for Depression in Older Adults: Randomized Controlled Trial. *J Med Internet Res.* 2024;26:e53001. doi:10.2196/53001.
 39. Xiang X, Kayser J, Turner S, Zheng C. Layperson-supported internet-delivered cognitive behavioral therapy for depression among older adults. *Res Soc Work Pract.* 2024;34(7):725-735. doi:10.1177/10497315231184143.
 40. Xiang X, Turner S, Ruiz-Sierra S, et al. Older Adults Experience with a Layperson-Supported Digital Mental Health Intervention for Depression: Qualitative Insights on Engagement. *Clin Gerontol.* doi:10.1080/07317115.2024.2395890.
 41. Gallagher-Thompson D, Thompson LW. Treating Late Life Depression: A Cognitive-Behavioral Therapy Approach, Therapist Guide. Oxford: Oxford University Press; 2009. ISBN:0195383699
 42. Laidlaw K, Thompson LW, Dick-Siskin L, Gallagher-Thompson D. Cognitive Behaviour Therapy with Older People. doi:https://doi.org/10.1002/9780470713402.

Protocol: iCBT for Older Adults

43. Lejuez CW, Hopko DR, Acierno R, Daughters SB, Pagoto SL. Ten Year Revision of the Brief Behavioral Activation Treatment for Depression: Revised Treatment Manual. *Behav Modif.* 2011;35(2):111-161. doi:10.1177/0145445510390929.
44. Xiang X, Kayser J, Sun Y, Himle J. Internet-Based Psychotherapy Intervention for Depression Among Older Adults Receiving Home Care: Qualitative Study of Participants' Experiences. *JMIR Aging.* 2021;4(4):e27630. doi:10.2196/27630.
45. Lehto T. Designing persuasive health behavior change interventions. In: Wickramasinghe N, Bali R, Suomi R, Kirn S, eds. *Critical Issues for the Development of Sustainable E-health Solutions. Healthcare Delivery in the Information Age.* Boston, MA: Springer; 2012. doi:10.1007/978-1-4614-1536-7_11.
46. Moyer-Gusé E. Toward a Theory of Entertainment Persuasion: Explaining the Persuasive Effects of Entertainment-Education Messages. *Commun Theory.* 2008;18(3):407-425. doi:10.1111/j.1468-2885.2008.00328.x.
47. Johnson J, Finn K. *Designing User Interfaces for an Aging Population: Towards Universal Design.* Morgan Kaufmann Publishers; 2017.
48. Schueller SM, Tomasino KN, Mohr DC. Integrating human support into behavioral intervention technologies: The efficiency model of support. *Clin Psychol-Sci Pr* 2017;24(1):27-45. doi:10.1111/cpsp.12173.
49. Katzman R, Brown T, Fuld P, Peck A, Schechter R, Schimmel H. Validation of a short Orientation-Memory-Concentration Test of cognitive impairment. *Am J Psychiatry.* 1983;140(6):734-739. doi:10.1176/ajp.140.6.734.
50. Posner K, Brown GK, Stanley B, et al. The Columbia–Suicide Severity Rating Scale: Initial Validity and Internal Consistency Findings From Three Multisite Studies With Adolescents and Adults. *Am J Psychiatry.* 2011;168(12):1266-1277. doi:10.1176/appi.ajp.2011.10111704.
51. Hinkin CH, Castellon SA, Dickson-Fuhrman E, Daum G, Jaffe J, Jarvik L. Screening for Drug and Alcohol Abuse Among Older Adults Using a Modified Version of the CAGE. *Am J Addict.* 2001;10(4):319-326. doi:10.1111/j.1521-0391.2001.tb00521.x.
52. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606-613. doi:10.1046/j.1525-1497.2001.016009606.x.
53. McIntyre RS, Konarski JZ, Mancini DA, et al. Measuring the severity of depression and remission in primary care: validation of the HAMD-7 scale [published correction appears in *CMAJ.* 2006 Jan 17;174(2):207]. *CMAJ.* 2005;173(11):1327-1334. doi:10.1503/cmaj.050786.
54. Choi NG, Hegel MT, Marti CN, Marinucci ML, Sirrianni L, Bruce ML. Telehealth Problem-Solving Therapy for Depressed Low-Income Homebound Older Adults. *Am J Geriatr Psychiatry.* 2014;22(3):263-271. doi:10.1016/j.jagp.2013.01.037.
55. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7. *Arch Intern Med.* 2006;166(10):1092. doi:10.1001/archinte.166.10.1092.
56. Koenig HG, Westlund RE, George LK, Hughes DC, Blazer DG, Hybels C. Abbreviating the Duke Social Support Index for Use in Chronically Ill Elderly Individuals. *Psychosomatics.* 1993;34(1):61-69. doi:10.1016/S0033-3182(93)71928-3.
57. EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy.* 1990;16(3):199-208. doi:10.1016/0168-8510(90)90421-9.
58. Pickard AS, Law EH, Jiang R, et al. United States Valuation of EQ-5D-5L Health States

Protocol: iCBT for Older Adults

- Using an International Protocol. *Value Health*. 2019;22(8):931-941. doi:10.1016/j.jval.2019.02.009.
59. Measuring Health and Disability : Manual for WHO Disability Assessment Schedule (WHODAS 2.0). www.who.int. Published June 16, 2012. [https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-\(-whodas-2.0\)](https://www.who.int/publications/i/item/measuring-health-and-disability-manual-for-who-disability-assessment-schedule-(-whodas-2.0)).
 60. Krebs EE, Lorenz KA, Bair MJ, et al. Development and initial validation of the PEG, a three-item scale assessing pain intensity and interference. *J Gen Intern Med*. 2009;24(6):733-738. doi:10.1007/s11606-009-0981-1.
 61. Beck JS. *Cognitive Behavior Therapy: Basics and Beyond*. 2nd ed. Guilford Press; 2011.
 62. Cuijpers P, Reijnders M, Huibers MJH. The Role of Common Factors in Psychotherapy Outcomes. *Annu Rev Clin Psychol*. 2019;15:207-231. doi:10.1146/annurev-clinpsy-050718-095424.
 63. Ryan RM, Deci EL. A self-determination theory approach to psychotherapy: The motivational basis for effective change. *Can Psychol Psychol Can*. 2008;49(3):186-193. doi:10.1037/a0012753.
 64. Jacob KL, Christopher MS, Neuhaus EC. Development and validation of the cognitive-behavioral therapy skills questionnaire. *Behav Modif*. 2011;35(6):595-618. doi:10.1177/0145445511419254.
 65. Manos RC, Kanter JW, Luo W. The behavioral activation for depression scale-short form: development and validation. *Behav Ther*. 2011;42(4):726-739. doi:10.1016/j.beth.2011.04.004.
 66. Beevers CG, Strong DR, Meyer B, Pilkonis PA, Miller IR. Efficiently assessing negative cognition in depression: an item response theory analysis of the Dysfunctional Attitude Scale. *Psychol Assess*. 2007;19(2):199-209. doi:10.1037/1040-3590.19.2.199.
 67. Marti CN, Choi NG. Measuring Social Engagement among Low-Income, Depressed Homebound Older Adults: Validation of the Social Engagement and Activities Questionnaire. *Clin Gerontol*. 2022;45(3):548-561. doi:10.1080/07317115.2020.1753275.
 68. Henson P, Wisniewski H, Hollis C, Keshavan M, Torous J. Digital mental health apps and the therapeutic alliance: initial review. *BJPsych Open*. 2019;5(1):e15. doi:10.1192/bjo.2018.86.
 69. Busselle R, Bilandzic H. Measuring Narrative Engagement. *Media Psychol*. 2009;12(4):321-347. doi:10.1080/15213260903287259.
 70. Igartua J. Identification with characters and narrative persuasion through fictional feature films. *Communications*. 2010;35(4): 347-373. <https://doi.org/10.1515/comm.2010.019>.
 71. Hadziahmetović N, Alispahić S, Tuce Đ, Hasanbegović-Anić E. Therapist's interpersonal style and therapy benefit as the determinants of personality self-reports in clients. *Vojnosanit Pregl*. 2016;73(2):135-145. doi:10.2298/vsp140911141h.
 72. Deci EL, Ryan RM. The “what” and “why” of goal pursuits: Human needs and the self-determination of behavior. *Psychological Inquiry*. 2000;11(4):227-268. doi:https://doi.org/10.1207/S15327965PLI1104_01.
 73. Ryan RM, Deci EL. Overview of self-determination theory: An organismic-dialectical perspective. *Handbook of self-determination research*. Rochester, NY, US: University of Rochester Press; 2002. p. 3-33.
 74. Johnston MM, Finney SJ. Measuring basic needs satisfaction: Evaluating previous research and conducting new psychometric evaluations of the Basic Needs Satisfaction in General Scale. *Contemp Educ Psychol*. 2010;35(4):280-296. doi:10.1016/j.cedpsych.2010.04.003.


Protocol: iCBT for Older Adults

75. Landreville P, Guérette A. Psychometric Properties of a Modified Version of the Treatment Evaluation Inventory for Assessing the Acceptability of Treatments for Geriatric Depression. *Canadian Journal on Aging / La Revue canadienne du vieillissement*. 1998;17(4):414-424. doi:10.1017/S071498080001268X.
76. Tober G, Clyne W, Finnegan O, Farrin A, Team IR. Validation of a Scale for Rating the Delivery of Psycho-Social Treatments for Alcohol Dependence and Misuse: The UKATT Process Rating Scale (PRS). *Alcohol and Alcoholism*. 2008;43(6):675-682. doi:https://doi.org/10.1093/alcalc/agn064.
77. Damschroder LJ, Reardon CM, Widerquist MAO, Lowery J. The updated Consolidated Framework for Implementation Research based on user feedback. *Implement Sci*. 2022;17(1):75. doi:10.1186/s13012-022-01245-0.
78. O’Laughlin KD, Martin MJ, Ferrer E. Cross-Sectional Analysis of Longitudinal Mediation Processes. *Multivar Behav Res*. 2018;53(3):375-402. doi:10.1080/00273171.2018.1454822
79. Cheong J. Accuracy of Estimates and Statistical Power for Testing Mediation in Latent Growth Curve Modeling. *Struct Equ Model Multidiscip J*. 2011;18(2):195-211. doi:10.1080/10705511.2011.557334.
80. SALDANA J. Coding Manual for Qualitative Researchers. SAGE; 2025.
81. Webb CA, Rosso IM, Rauch SL. Internet-Based Cognitive-Behavioral Therapy for Depression: Current Progress and Future Directions. *Harv Rev Psychiatry*. 2017;25(3):114-122. doi:10.1097/HRP.0000000000000139.
82. Christensen H, Griffiths KM, Mackinnon AJ, Brittliffe K. Online randomized controlled trial of brief and full cognitive behaviour therapy for depression. *Psychol Med*. 2006;36(12):1737-1746. doi:10.1017/S0033291706008695.
83. Karyotaki E, Efthimiou O, Miguel C, et al. Internet-Based Cognitive Behavioral Therapy for Depression: A Systematic Review and Individual Patient Data Network Meta-analysis. *JAMA Psychiatry*. 2021;78(4):361–371. doi:10.1001/jamapsychiatry.2020.4364.
84. Baumel A, Muench F, Edan S, Kane JM. Objective User Engagement With Mental Health Apps: Systematic Search and Panel-Based Usage Analysis. *J Med Internet Res*. 2019;21(9):e14567. doi:10.2196/14567.
85. Baumel A, Edan S, Kane JM. Is there a trial bias impacting user engagement with unguided e-mental health interventions? A systematic comparison of published reports and real-world usage of the same programs. *Transl Behav Med*. 2019;9(6):1020-1033. doi:10.1093/tbm/ibz147.
86. Adorjan K, Stubbe HC. Insight into the long-term psychological impacts of the COVID-19 pandemic. *Eur Arch Psychiatry Clin Neurosci*. 2023;273(2):287-288. doi:10.1007/s00406-023-01599-6.

Supplementary Files

Figures

Participant flow CONSORT diagram.



Multimedia Appendixes

Peer review report.

URL: <http://asset.jmir.pub/assets/d7e54e93b6d1ff632ec6490bbbfe7932.pdf>

Preprint
JMIR Publications

TOC/Feature image for homepages

Empower@Home user mockup.

