

# **Unlocking the Secrets Beneath: Utilizing the Water Quality Index (WQI) to Assess Groundwater Contamination Near Dumpsites - A Case Study of Port Harcourt**

Tomquin Abiye, Morufu Olalekan Raimi

Submitted to: JMIR Preprints  
on: March 08, 2025

**Disclaimer:** © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

## ***Table of Contents***

---

|                                 |          |
|---------------------------------|----------|
| <b>Original Manuscript.....</b> | <b>5</b> |
|---------------------------------|----------|

Preprint  
JMIR Publications

# Unlocking the Secrets Beneath: Utilizing the Water Quality Index (WQI) to Assess Groundwater Contamination Near Dumpsites – A Case Study of Port Harcourt

Tomquin Abiye<sup>1\*</sup> PhD; Morufu Olalekan Raimi<sup>2\*</sup> PhD

<sup>1</sup>Department of Environmental management Faculty of Environmental Sciences, Rivers State University (RSU) Ogbia NG

<sup>2</sup>2Niger-Delta Institute for Emerging and Re-Emerging Infectious Diseases (NDIERID) Federal University Otuoke Ogbia NG

\*these authors contributed equally

## Corresponding Author:

Morufu Olalekan Raimi PhD

2Niger-Delta Institute for Emerging and Re-Emerging Infectious Diseases (NDIERID)

Federal University Otuoke

Otuoke

Ogbia

NG

## Abstract

**Background:** Groundwater contamination poses a significant public health risk, particularly in urban areas with inadequate waste management. Dumpsites serve as major sources of pollutants, including heavy metals, which infiltrate aquifers through leachate migration. Port Harcourt, Nigeria, faces increasing groundwater quality concerns due to the proliferation of uncontrolled waste disposal sites.

**Objective:** This study aims to evaluate the spatial and seasonal variations in groundwater quality around dumpsites in Port Harcourt and determine the suitability of groundwater for drinking based on WQI values. It also seeks to identify contamination patterns and assess the influence of rainfall on pollutant dispersion. Furthermore, the study compares findings with global research to establish broader implications for waste management and public health. By doing so, it provides a scientific basis for policy recommendations aimed at mitigating groundwater pollution.

**Methods:** Groundwater samples were collected from various locations around major dumpsites in Port Harcourt during dry and rainy seasons. Physicochemical parameters, including heavy metal concentrations, were analyzed to compute WQI values. Comparative analysis with previous studies was conducted to validate observed contamination trends. The impact of leachate migration on water quality was assessed using seasonal variations in WQI values.

**Results:** Findings reveal significant spatial and seasonal fluctuations in groundwater quality. While Choba exhibited excellent water quality, Sasun, Olumeni, and Epirikom recorded dangerously high WQI values, indicating unsuitability for drinking. Seasonal variations showed that rainfall exacerbated contamination levels, as seen in Eleme, where WQI increased from 56.362 in the dry season to 140.928 in the rainy season. The study aligns with previous research from India, China, and Ghana, demonstrating that landfill leachates and surface runoff are key contributors to groundwater degradation.

**Conclusions:** The study confirms that dumpsite leachates significantly impact groundwater quality, posing a major risk to public health. The high WQI values in several locations highlight the need for urgent interventions. Findings align with global research on groundwater contamination, emphasizing the critical role of effective waste management in reducing environmental pollution. To mitigate groundwater pollution from dumpsite leachates, it is essential to implement stringent waste management policies that regulate landfill operations and prevent leachate infiltration into aquifers. Establishing continuous groundwater monitoring programs can help detect contamination trends early and guide timely intervention measures. Additionally, promoting alternative potable water sources in highly contaminated areas is crucial to reducing health risks for affected communities. The adoption of modern landfill technologies, such as leachate treatment and containment systems, should be prioritized to minimize pollution and safeguard water resources for future generations. This study contributes to the growing body of research on groundwater contamination by providing empirical evidence of the impact of dumpsites in an urban African setting. The findings underscore the urgent need for improved waste management policies and public health interventions. By aligning with global research, this

study reinforces the importance of sustainable environmental practices to safeguard water resources and protect communities from the adverse effects of pollution.

(JMIR Preprints 08/03/2025:73643)

DOI: <https://doi.org/10.2196/preprints.73643>

## Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <http://www.jmir.org>

No. Please do not make my accepted manuscript PDF available to anyone.

## Original Manuscript

## Unlocking the Secrets Beneath: Utilizing the Water Quality Index (WQI) to Assess Groundwater Contamination Near Dumpsites – A Case Study of Port Harcourt

<sup>1</sup>**Tomquin Abiye** 0009-0002-5868-5025

<sup>1</sup>Department of Environmental management, Faculty of Environmental Sciences, Rivers State University (RSU) Rivers State, Nigeria. [tomquinabiye@ust.edu.ng](mailto:tomquinabiye@ust.edu.ng)

<sup>2</sup>**Morufu Olalekan Raimi** 

<sup>2</sup>Niger-Delta Institute for Emerging and Re-Emerging Infectious Diseases (NDIERID), Federal University Otuoke, Bayelsa State. Nigeria. [raimimo@fuotuoche.edu.ng](mailto:raimimo@fuotuoche.edu.ng)

### Abstract

**Rationale:** Groundwater contamination poses a significant public health risk, particularly in urban areas with inadequate waste management. Dumpsites serve as major sources of pollutants, including heavy metals, which infiltrate aquifers through leachate migration. Port Harcourt, Nigeria, faces increasing groundwater quality concerns due to the proliferation of uncontrolled waste disposal sites. **Objectives:** This study aims to evaluate the spatial and seasonal variations in groundwater quality around dumpsites in Port Harcourt and determine the suitability of groundwater for drinking based on WQI values. It also seeks to identify contamination patterns and assess the influence of rainfall on pollutant dispersion. Furthermore, the study compares findings with global research to establish broader implications for waste management and public health. By doing so, it provides a scientific basis for policy recommendations aimed at mitigating groundwater pollution. **Methods:** Groundwater samples were collected from various locations around major dumpsites in Port Harcourt during dry and rainy seasons. Physicochemical parameters, including heavy metal concentrations, were analyzed to compute WQI values. Comparative analysis with previous studies was conducted to validate observed contamination trends. The impact of leachate migration on water quality was assessed using seasonal variations in WQI values. **Results:** Findings reveal significant spatial and seasonal fluctuations in groundwater quality. While Choba exhibited excellent water quality, Sasun, Olumeni, and Epirikom recorded dangerously high WQI values, indicating unsuitability for drinking. Seasonal variations showed that rainfall exacerbated contamination levels, as seen in Eleme, where WQI increased from 56.362 in the dry season to 140.928 in the rainy season. The study aligns with previous research from India, China, and Ghana, demonstrating that landfill leachates and surface runoff are key contributors to groundwater degradation. **Conclusion:** The study confirms that dumpsite leachates significantly impact groundwater quality, posing a major risk to public health. The high WQI values in several locations highlight the need for urgent interventions. Findings align with global research on groundwater contamination, emphasizing the critical role of effective waste management in reducing environmental pollution. **Recommendation:** To mitigate groundwater pollution from dumpsite leachates, it is essential to implement stringent waste management policies that regulate landfill operations and prevent leachate infiltration into aquifers. Establishing continuous groundwater monitoring programs can help detect contamination trends early and guide timely intervention measures. Additionally, promoting alternative potable water sources in highly contaminated areas is crucial to reducing health risks for affected communities. The adoption of modern landfill technologies, such as leachate treatment and containment systems, should be prioritized to minimize pollution and safeguard water resources for future generations. **Significance Statement:** This study contributes to the growing body of research on groundwater contamination by providing empirical evidence of the impact of dumpsites in an urban African setting. The findings underscore the urgent need for improved waste management policies and public health interventions. By aligning with global research, this study reinforces the importance of sustainable environmental practices to safeguard water resources and protect communities from the adverse effects of pollution.

**Keywords:** Groundwater Quality; Water Quality Index (WQI); Dumpsites; Contamination;

## Port Harcourt

## 1. Introduction

Groundwater serves as a primary source of drinking water, agricultural irrigation, and industrial processes, making it essential for human life and economic development [1-5]. In urban areas of developing countries, groundwater faces significant contamination risks from human activities, particularly waste disposal practices [6-10]. Open dumpsites, characterized by unregulated solid waste accumulation, pose environmental and public health threats due to potential leachate infiltration into soil and groundwater [6-10]. Port Harcourt, Nigeria, exemplifies challenges associated with inadequate waste management. Rapid urbanization has led to numerous dumpsites within and around the city, increasing the risk of groundwater contamination [11-23]. Pollutants such as heavy metals, pathogens, and organic chemicals from these sources threaten safe drinking water supply, agricultural productivity, and ecosystem health [24-50]. However, a total of 45 studies conducted in Nigeria [1-5, 11-23, 51-77] were found to be significantly affects residents relying on groundwater for daily consumption, exposing them to health risks and potential financial costs of waterborne diseases. The problem has persisted for over a decade, with studies indicating a historical trend of increasing groundwater contamination correlating with urban expansion and waste generation [6-9]. The contamination is more pronounced in urban and peri-urban areas where dumpsites are prevalent, disproportionately impacting low-income communities lacking access to alternative water sources [56-68].

Existing studies have assessed groundwater contamination near dumpsites in Port Harcourt. For instance, Ugwoha and Emete [78] (2015) analyzed the impact of open dumpsite leachate on groundwater quality at the Alakahia dumpsite, finding elevated levels of contaminants. Similarly, Alaye *et al.* [79] conducted a health risk assessment of heavy metal contamination in groundwater near waste dumpsites, highlighting significant health risks. Additionally, Eseyin *et al.* [80] evaluated the effect of municipal solid waste leachate on groundwater quality, reporting contamination by various pollutants. Despite these studies, there remains a lack of comprehensive research integrating physicochemical and biological parameters to assess groundwater quality near dumpsites in Port Harcourt. Specifically, the application of the Water Quality Index (WQI) as a holistic assessment tool has not been extensively utilized in this context. Addressing this gap is crucial for informed decision-making in environmental management and public health protection. The relevance of this research extends beyond environmental science, intersecting with public health, urban planning, and socio-economic development. By providing empirical data on groundwater quality, the study informs public health initiatives aimed at reducing waterborne diseases. It also offers evidence-based recommendations for urban planners to design sustainable waste management systems. Furthermore, understanding the socio-economic implications of groundwater contamination can guide policies to mitigate financial burdens on affected communities. This study aims to apply the Water Quality Index (WQI) to evaluate groundwater quality near dumpsites in Port Harcourt. By analyzing both



physicochemical and biological parameters, the research seeks to: Determine the extent of groundwater contamination near selected dumpsites; Assess potential health risks associated with the consumption of contaminated groundwater and provide actionable insights for policymakers, environmental scientists, and public health experts to improve waste management practices and protect groundwater resources. Thus, this study addresses a critical environmental and public health issue by evaluating groundwater quality near dumpsites in Port Harcourt using the Water Quality Index. By filling existing research gaps, the findings aim to inform and influence policies and practices that safeguard groundwater resources, ensuring a safer and healthier future for the community.

## **2. Methodology**

### **2.1 Research Design**

This study employs a quasi-experimental research design to assess groundwater quality near dumpsites in Port Harcourt metropolis using the Water Quality Index (WQI). The quasi-experimental approach allows for the examination of groundwater contamination in a real-world setting without random assignment, providing insights into the impact of dumpsites on groundwater quality.

### **2.2 Study Area**

Port Harcourt is situated within the Niger Delta sedimentary basin, comprising three main formations: the Akata Formation (Paleocene), Agbada Formation (Eocene), and Benin Formation (Miocene to recent) [81]. The region's soils are a mix of marine and fluvial marine sediments,

mangrove swamp alluvial soils, and freshwater brown and sandy loams. Clay soils prevalent in the area hinder water percolation, potentially leading to flooding during heavy rainfall, which can impact groundwater quality.

### **2.3 Data Collection**

Data were collected through primary and secondary sources. Primary data involved direct field surveys, collecting groundwater samples at varying distances (200m, 400m, and 600m) from 10 dumpsites in Port Harcourt. A total of 60 samples were collected, with 30 during the rainy season and 30 during the dry season. Secondary data were sourced from published and unpublished reports, periodicals, and relevant scientific literature to supplement the primary data.

### **2.4 Sample Collection and Preservation**

Groundwater samples were collected following strict protocols to minimize contamination [82]. Sampling bottles were rinsed with the same groundwater before collection, properly labeled, and preserved in coolers before transportation to an approved laboratory for analysis. Sampling locations were recorded using GPS technology to ensure accuracy and facilitate spatial analysis. Sampling during both rainy and dry seasons assessed how seasonal variations might affect groundwater quality.

### **2.5 Laboratory Analysis**

Collected samples were analyzed for key physicochemical and microbial parameters, including pH, electrical conductivity, total organic carbon (TOC), total dissolved solids (TDS), chlorides, sulfates, nitrates, phosphates, total hardness, iron, manganese, lead, aluminum, copper,

zinc, and *E. coli*. These parameters were chosen based on their significance in identifying groundwater contamination, particularly from leachates near dumpsites. Testing procedures adhered to international best practices, specifically those outlined in the *Standard Methods for the Examination of Water and Wastewater*, ensuring reliability and accuracy [82].

## 2.6 Water Quality Index (WQI) Calculation

The WQI simplifies complex water quality data into a single, easily understandable metric, facilitating better decision-making for water resource management. This study adopts the weighted arithmetic water quality index method (WAWQIM) to assess groundwater contamination in Port Harcourt metropolis, following the method used by Morufu [83]. The quality rating scale was assigned to the parameters, weighted according to their relative importance in overall water quality. The WQI is calculated using the formula:

$$WQI = \frac{\sum (Q_i \times W_i)}{\sum W_i}$$

Where  $Q_i$  is the quality rating and  $W_i$  is the unit weight of each parameter. For this study, the following parameters were selected: electrical conductivity, pH, manganese, turbidity, lead, *E. coli*, aluminum, TOC, TDS, nitrate, sulfate, phosphate, total hardness, iron, copper, and zinc. The WQI rating by Morufu [83] is as follows:

- i. Above 100: Unsuitable for drinking
- ii. 76–100: Very Bad
- iii. 51–75: Poor

- iv. 26–50: Good
- v. 0–25: Excellent

### 3. Results

Table 1 shows the Water Quality Index (WQI) values for groundwater samples collected from various dumpsites in Port Harcourt metropolis reveal significant spatial and seasonal variations in water quality. Among the sampled locations, Choba stands out as the only site with consistently excellent water quality, with WQI values of 10.248 in the dry season and 10.655 in the rainy season. This indicates that groundwater in Choba is largely unaffected by dumpsite leachates and meets drinking water standards. In contrast, other locations such as Epirikom, Olumeni, and Sasun recorded alarmingly high WQI values, far exceeding the threshold for safe drinking water. For instance, Sasun had the highest contamination levels, with WQI values of 2553.23 in the dry season and 2634.939 in the rainy season, making it the most severely impacted site. Seasonal variations also play a crucial role in groundwater quality, as evident from the data. In many locations, WQI values tend to be higher during the rainy season, which suggests that increased rainfall and surface runoff may exacerbate contamination by facilitating the infiltration of leachates from the dumpsites into groundwater. This trend is particularly notable in sites like Eleme, where the WQI rises from 56.362 (poor) in the dry season to 140.928 (unfit for drinking) in the rainy season. A similar seasonal effect is observed at Agholu, where the WQI jumps from 124.855 to 165.791. However, in some locations, such as Eneka, seasonal fluctuations appear

negligible, with WQI values remaining stable at approximately 41, indicating that groundwater contamination is less influenced by seasonal changes in these areas.

The implications of these findings are profound, particularly for public health and water resource management in Port Harcourt. The high WQI values at several sites suggest that groundwater near many dumpsites is heavily contaminated and unsafe for consumption. Prolonged exposure to such polluted water sources can lead to severe health issues, including waterborne diseases, heavy metal poisoning, and other long-term health complications. Communities relying on these groundwater sources may face increased risks of gastrointestinal infections, kidney damage, and neurological disorders due to contaminants such as lead, nitrates, and microbial pathogens. The findings highlight the urgent need for interventions such as improved waste management practices, leachate treatment facilities, and public awareness campaigns to mitigate groundwater contamination in these areas. From a policy and environmental management perspective, this study underscores the necessity of stricter regulations and better enforcement of waste disposal practices in urban areas. The government and environmental agencies must implement measures such as controlled landfill engineering, groundwater monitoring programs, and the provision of alternative safe drinking water sources in affected communities. Additionally, long-term remediation strategies, including phytoremediation and bioremediation, could be explored to mitigate the impact of dumpsite pollution. Overall, the WQI assessment provides valuable evidence that can guide policy

decisions aimed at ensuring sustainable groundwater management and protecting public health in Port Harcourt metropolis.

**Table 1 Water Quality Index in relation to Seasons Across Sampled Dumpsites**

| <b>Sampled Dumpsites</b> | <b>Season</b> | <b>WQI Value</b> | <b>WQI Status</b>  |
|--------------------------|---------------|------------------|--------------------|
| <b>Agholu</b>            | Dry           | 124.855          | Unfit for drinking |
|                          | Rainy         | 165.791          | Unfit for drinking |
| <b>Choba</b>             | Dry           | 10.248           | Excellent          |
|                          | Rainy         | 10.655           | Excellent          |
| <b>Eleme</b>             | Dry           | 56.362           | Poor               |
|                          | Rainy         | 140.928          | Unfit for drinking |
| <b>Eliozu</b>            | Dry           | 73.502           | Very poor          |
|                          | Rainy         | 43.221           | Good               |
| <b>Eneka</b>             | Dry           | 41.466           | Good               |
|                          | Rainy         | 41.408           | Good               |
| <b>Epirikom</b>          | Dry           | 337.154          | Unfit for drinking |
|                          | Rainy         | 274.744          | Unfit for drinking |
| <b>Olumeni</b>           | Dry           | 372.698          | Unfit for drinking |
|                          | Rainy         | 433.028          | Unfit for drinking |
| <b>Oyigbo</b>            | Dry           | 172.53           | Unfit for drinking |
|                          | Rainy         | 178.979          | Unfit for drinking |
| <b>Rumuola</b>           | Dry           | 167.061          | Unfit for drinking |
|                          | Rainy         | 167.061          | Unfit for drinking |
| <b>Sasun</b>             | Dry           | 2553.23          | Unfit for drinking |
|                          | Rainy         | 2634.939         | Unfit for drinking |

**Source: Researcher Field work (2021)**

Table 2 shows the Water Quality Index (WQI) values across different sampling locations at varying distances (200m, 400m, and 600m) from dumpsites in Port Harcourt reveal significant spatial variations in groundwater contamination. Choba, Eneka, and Rumuolumeni exhibit the best water quality, with WQI values consistently falling within the “excellent” category at all sampling distances, indicating minimal

leachate infiltration from dumpsites. In contrast, Sasun recorded the highest contamination levels, with a WQI of 7222.46 at 200m, highlighting extreme groundwater pollution near the dumpsite. This severe contamination persists at farther distances (400m and 600m), suggesting extensive groundwater infiltration by pollutants. Similarly, Epirikom, Rumuola, and Oyigbo show consistently high WQI values across all sampling distances, classifying them as “unfit for drinking.” The trend in contamination levels suggests that proximity to dumpsites significantly affects groundwater quality, but other environmental and hydrogeological factors may also be at play. While some locations, like Agholu and Eleme, experience high WQI values at 200m and 400m but improved conditions at 600m, others, such as Rumuola and Rumuolumeni, remain highly contaminated even at greater distances. This indicates that groundwater pollution from dumpsites does not dissipate uniformly with distance, likely due to factors such as subsurface flow, soil composition, and the extent of leachate migration. The elevated WQI at deeper distances in Rumuola and Sasun suggests that contamination plumes may be spreading further through underground water channels, making distance alone an unreliable indicator of water safety.

The real-world implications of these findings are critical, particularly for communities relying on groundwater for drinking and domestic use. The widespread classification of groundwater as “unfit for drinking” in several locations highlights the urgent need for intervention. Prolonged consumption of contaminated groundwater can lead to severe health issues, including heavy metal toxicity, gastrointestinal infections, and

long-term organ damage. The presence of high WQI values even at 600m from certain dumpsites suggests that leachate pollution is pervasive and cannot be mitigated simply by increasing the distance of wells from dumpsites. This calls for stricter regulation of waste disposal practices, improved landfill engineering, and the implementation of leachate management systems to prevent further groundwater contamination. From a policy standpoint, these findings emphasize the need for comprehensive groundwater monitoring and remediation efforts in Port Harcourt. Authorities should prioritize the enforcement of environmental laws to ensure that dumpsites are properly managed, and communities should be provided with alternative potable water sources. Additionally, remediation strategies such as bioremediation and the use of constructed wetlands could be explored to reduce the impact of existing contamination. The extreme pollution levels in certain locations, particularly Sasun, indicate that immediate intervention is necessary to prevent long-term environmental and public health crises. These insights serve as a valuable basis for developing targeted policies that protect groundwater resources and ensure safe drinking water for affected populations.

**Table 2. Water Quality Index Values Across Sampling Points in the sampled Dumpsites.**

| Sample Dumpsite | Sample locations | WQI      | WQI Status     |     |
|-----------------|------------------|----------|----------------|-----|
| <b>Agholu</b>   | BH1(200m)        | 147.2567 | Unfit drinking | for |
|                 | BH2(400m)        | 175.0579 | Unfit drinking | for |
|                 | BH3(600m)        | 113.6544 | Unfit drinking | for |
| <b>Choba</b>    | BH1(200m)        | 10.23994 | Excellent      |     |



|                    |           |          |                |     |
|--------------------|-----------|----------|----------------|-----|
| <b>Eleme</b>       | BH2(400m) | 10.86233 | Excellent      |     |
|                    | BH3(600m) | 10.25229 | Excellent      |     |
|                    | BH1(200m) | 121.6781 | Unfit drinking | for |
| <b>Eliozu</b>      | BH2(400m) | 150.5528 | Unfit drinking | for |
|                    | BH3(600m) | 23.70329 | Excellent      |     |
|                    | BH1(200m) | 106.3705 | Unfit drinking | for |
| <b>Eneka</b>       | BH2(400m) | 11.6465  | Excellent      |     |
|                    | BH3(600m) | 57.06804 | Poor           |     |
|                    | BH1(200m) | 10.26892 | Excellent      |     |
| <b>Epirikom</b>    | BH2(400m) | 103.6225 | Unfit drinking | for |
|                    | BH3(600m) | 10.41918 | Excellent      |     |
|                    | BH1(200m) | 419.6739 | Unfit drinking | for |
| <b>Rumuolumeni</b> | BH2(400m) | 202.2795 | Unfit drinking | for |
|                    | BH3(600m) | 295.8937 | Unfit drinking | for |
|                    | BH1(200m) | 10.37225 | Excellent      |     |
| <b>Oyigbo</b>      | BH2(400m) | 644.3554 | Unfit drinking | for |
|                    | BH3(600m) | 553.8604 | Unfit drinking | for |
|                    | BH1(200m) | 162.5117 | Unfit drinking | for |
| <b>Rumuola</b>     | BH2(400m) | 187.2124 | Unfit drinking | for |
|                    | BH3(600m) | 177.539  | Unfit drinking | for |
|                    | BH1(200m) | 12.50133 | Excellent      |     |
| <b>Sasun</b>       | BH2(400m) | 198.2805 | Unfit drinking | for |
|                    | BH3(600m) | 290.4002 | Unfit drinking | for |
|                    | BH1(200m) | 7222.46  | Unfit drinking | for |
|                    | BH2(400m) | 341.1783 | Unfit drinking | for |
|                    | BH3(600m) | 218.6153 | Unfit drinking | for |

**Source: Researcher Field work (2021)**

#### **4. Discussion**

#### **4.1 Spatial Variations Groundwater Contamination Near Dumpsites: A WQI-Based Analysis in Port Harcourt**

The findings of this study reveal significant spatial and seasonal variations in groundwater quality around dumpsites in Port Harcourt. The Water Quality Index (WQI) values indicate that while Choba consistently exhibits excellent water quality, locations such as Sasun, Olumeni, and Epirikom have dangerously high WQI values, rendering their groundwater unfit for drinking. Seasonal fluctuations suggest that rainfall exacerbates contamination by increasing leachate infiltration, as observed in Eleme, where WQI rises from 56.362 in the dry season to 140.928 in the rainy season. These findings are consistent with Kumar *et al.* [84], who reported that WQI values in Anna Nagar, Chennai, fluctuated seasonally due to surface runoff and leachate percolation, making groundwater quality highly variable throughout the year. A similar trend was reported by Boateng *et al.* [85] in their study on groundwater contamination near the Oti landfill site in Kumasi, Ghana. Their findings highlight the severe impact of landfill leachates on groundwater quality, where heavy metal contamination significantly contributed to high WQI values. This aligns with the present study, particularly in locations like Sasun and Olumeni, where extreme contamination levels suggest extensive leachate migration. Additionally, Muhammad *et al.* [86] utilized the DRASTIC index method in Lahore, Pakistan, to evaluate groundwater vulnerability and found that areas close to waste disposal sites exhibited heightened contamination risks, similar to the patterns observed in Port Harcourt. Moreover, the study's findings resonate with the work of Li *et al.* [87], who

assessed groundwater quality in Pengyang County, Northwest China. Their research demonstrated that human activities, particularly waste disposal practices, significantly influence WQI levels. The consistently poor groundwater quality near dumpsites in Port Harcourt further corroborates Li *et al.*'s [87] assertion that uncontrolled waste disposal and inadequate landfill management exacerbate groundwater contamination. Similarly, Prasad & Sangita [88] examined groundwater pollution in an abandoned open-cast mine repurposed for municipal waste disposal and found excessive heavy metal accumulation, reinforcing the conclusion that dumpsites contribute significantly to groundwater degradation.

Furthermore, Han *et al.* [89] reviewed groundwater contamination near municipal solid waste dumpsites in China and reported that heavy metal leaching from waste was a dominant factor in determining water quality. This aligns with the present study, where high WQI values in certain areas indicate potential heavy metal contamination. Ojekunle *et al.* [90] also assessed water quality indices and ecological risk factors for heavy metals in scrap yard neighborhoods, emphasizing that waste accumulation sites pose severe threats to public health. Similarly, the research by Naveedullah *et al.* [91] on agricultural soil contamination due to heavy metals underscores the environmental risks associated with waste disposal, which likely contribute to the poor groundwater quality observed in Port Harcourt. Lastly, Adimalla *et al.* [92] studied groundwater suitability for domestic and agricultural purposes in Telangana, India, and found that proximity to industrial and waste disposal sites significantly affected water quality, mirroring the trends identified in this study. In

conclusion, the present study's findings align with a substantial body of existing research, reinforcing the critical link between dumpsite leachates and groundwater contamination. The correlation with previous studies underscores the urgent need for stringent waste management policies and groundwater remediation efforts in Port Harcourt. The widespread presence of high WQI values in multiple locations highlights the necessity for regulatory interventions to mitigate groundwater pollution and protect public health. Moving forward, implementing leachate containment strategies and promoting alternative potable water sources will be essential in safeguarding water quality in affected areas.

#### **4.2 Seasonal Variations in Groundwater Contamination Near Dumpsites in Port Harcourt: Insights from Water Quality Index Analysis**

The findings of this study reveal significant spatial and seasonal variations in groundwater quality around dumpsites in Port Harcourt. The Water Quality Index (WQI) values indicate that while Choba consistently exhibits excellent water quality, locations such as Sasun, Olumeni, and Epirikom have dangerously high WQI values, rendering their groundwater unfit for drinking. Seasonal fluctuations suggest that rainfall exacerbates contamination by increasing leachate infiltration, as observed in Eleme, where WQI rises from 56.362 in the dry season to 140.928 in the rainy season. These findings are consistent with Kumar *et al.* [84], who reported that WQI values in Anna Nagar, Chennai, fluctuated seasonally due to surface runoff and leachate percolation, making groundwater quality highly variable throughout the year. A similar trend was reported by

Boateng *et al.* [85] in their study on groundwater contamination near the Oti landfill site in Kumasi, Ghana. Their findings highlight the severe impact of landfill leachates on groundwater quality, where heavy metal contamination significantly contributed to high WQI values. This aligns with the present study, particularly in locations like Sasun and Olumeni, where extreme contamination levels suggest extensive leachate migration. Additionally, Muhammad *et al.* [86] utilized the DRASTIC index method in Lahore, Pakistan, to evaluate groundwater vulnerability and found that areas close to waste disposal sites exhibited heightened contamination risks, similar to the patterns observed in Port Harcourt.

Moreover, the study's findings resonate with the work of Li *et al.* [87], who assessed groundwater quality in Pengyang County, Northwest China. Their research demonstrated that human activities, particularly waste disposal practices, significantly influence WQI levels. The consistently poor groundwater quality near dumpsites in Port Harcourt further corroborates Li *et al.*'s [87] assertion that uncontrolled waste disposal and inadequate landfill management exacerbate groundwater contamination. Similarly, Prasad & Sangita [88] examined groundwater pollution in an abandoned open-cast mine repurposed for municipal waste disposal and found excessive heavy metal accumulation, reinforcing the conclusion that dumpsites contribute significantly to groundwater degradation.

The spatial variations in WQI values observed in this study align with the findings of Carpenter *et al.* [93], who used resistivity surveys to identify groundwater contamination near a dumpsite in Maoming, China.

Their study showed that contamination levels varied with distance from the dumpsite, similar to the findings in Port Harcourt, where locations such as Sasun and Epirikom remained highly contaminated even at greater distances. Additionally, Aizebeokhai & Olayinka [81] highlighted the role of subsurface geological structures in groundwater contamination, supporting the conclusion that factors such as soil composition and hydrogeology influence contamination spread in Port Harcourt. Han *et al.* [89] reviewed groundwater contamination near municipal solid waste dumpsites in China and reported that heavy metal leaching from waste was a dominant factor in determining water quality. This aligns with the present study, where high WQI values in certain areas indicate potential heavy metal contamination. Ojekunle *et al.* [90] also assessed water quality indices and ecological risk factors for heavy metals in scrap yard neighborhoods, emphasizing that waste accumulation sites pose severe threats to public health. Similarly, the research by Naveedullah *et al.* [91] on agricultural soil contamination due to heavy metals underscores the environmental risks associated with waste disposal, which likely contribute to the poor groundwater quality observed in Port Harcourt.

The findings of Batarseh *et al.* [94] further reinforce the present study by demonstrating the widespread impact of waste disposal on groundwater quality. Their study in Abu Dhabi showed that groundwater in proximity to waste disposal sites exhibited significantly higher levels of contamination, similar to the trends observed in Port Harcourt. Likewise, Armanuos *et al.* [95] used GIS-based WQI analysis to assess groundwater pollution in the Western Nile Delta, Egypt, and found that industrial and

municipal waste significantly deteriorated water quality, echoing the contamination patterns in this study. Furthermore, El Osta *et al.* [96] examined groundwater quality in Makkah Province, Saudi Arabia, and reported that unregulated waste disposal sites led to persistent contamination, reinforcing the necessity for improved landfill management and water quality monitoring in Port Harcourt. In conclusion, the present study's findings align with a substantial body of existing research, reinforcing the critical link between dumpsite leachates and groundwater contamination. The correlation with previous studies underscores the urgent need for stringent waste management policies and groundwater remediation efforts in Port Harcourt. The widespread presence of high WQI values in multiple locations highlights the necessity for regulatory interventions to mitigate groundwater pollution and protect public health. Moving forward, implementing leachate containment strategies and promoting alternative potable water sources will be essential in safeguarding water quality in affected areas.

## 5. Summary

This study assessed groundwater contamination near dumpsites in Port Harcourt using the Water Quality Index (WQI) as an indicator of water safety. The findings revealed significant spatial and seasonal variations in groundwater quality, with some locations, such as Choba, maintaining excellent water conditions, while others, including Sasun, Epirikom, and Rumuola, exhibited extreme contamination. The influence of rainfall was evident, as leachate infiltration intensified water pollution during the rainy season, leading to elevated WQI values in certain areas. These findings

align with global research, demonstrating that dumpsite leachates serve as a primary source of groundwater pollution, exacerbated by poor waste management and hydrogeological conditions. The study further highlights that contamination does not dissipate uniformly with distance from dumpsites, suggesting that subsurface flow dynamics, soil composition, and landfill conditions influence pollutant migration. The consistently poor water quality in several areas underscores the urgent need for intervention, as groundwater remains a vital source of drinking and domestic water for many communities. By correlating these findings with previous literature, this study reinforces the conclusion that unregulated waste disposal severely degrades groundwater quality, posing long-term environmental and public health risks.

## **6. Conclusion**

The study's results confirm that dumpsites in Port Harcourt significantly impact groundwater quality, with several locations consistently classified as “unfit for drinking” based on WQI values. High contamination levels, particularly in Sasun, Epirikom, and Rumuola, indicate the extensive migration of pollutants through groundwater channels. Seasonal trends suggest that rainfall enhances leachate percolation, further deteriorating water quality. The presence of high WQI values even at 600m from some dumpsites suggests that groundwater contamination extends far beyond immediate landfill boundaries. Comparing these findings with previous studies from China, Ghana, and Egypt highlights a global pattern of groundwater contamination due to improper waste disposal. The influence of local hydrogeological factors,



such as subsurface flow and soil composition, further complicates pollution dynamics, making distance alone an unreliable predictor of water safety. This study emphasizes the need for continuous groundwater monitoring and robust waste management strategies to mitigate environmental degradation and protect public health.

## **7. Policy Implications**

The widespread contamination of groundwater near dumpsites in Port Harcourt necessitates stricter environmental regulations to control waste disposal practices. The findings suggest that existing landfill management strategies are inadequate, allowing pollutants to infiltrate groundwater supplies. Authorities must enforce stringent regulations on landfill engineering, including the mandatory use of leachate containment systems, liners, and proper waste segregation to minimize contamination risks. Additionally, policies should mandate periodic groundwater quality assessments to identify pollution hotspots and ensure prompt remediation. Beyond landfill regulations, urban planning policies must prioritize the relocation of communities relying on highly contaminated water sources and provide alternative potable water solutions. Public health policies should also integrate groundwater safety into broader environmental health programs, ensuring that communities are educated on the risks associated with consuming untreated groundwater. By implementing these measures, policymakers can mitigate the long-term health and ecological impacts of dumpsite-induced groundwater contamination.


## **8. Recommendations**

To address the severe groundwater contamination observed in this study, immediate and long-term interventions are necessary. First, improved landfill engineering practices, such as the installation of impermeable liners and leachate collection systems, should be prioritized to reduce the migration of pollutants into groundwater. Additionally, the establishment of buffer zones around dumpsites, where groundwater extraction for drinking purposes is prohibited, could help minimize public exposure to contaminated water sources. Long-term solutions should focus on enhancing waste management infrastructure through recycling programs, waste-to-energy initiatives, and the development of modern sanitary landfills with proper environmental safeguards. Routine groundwater monitoring using WQI and other hydrogeochemical assessments should be institutionalized to detect contamination trends early. Furthermore, investment in alternative potable water sources, such as rainwater harvesting and treated surface water systems, is essential to reduce dependency on groundwater in high-risk areas.

## **9. Significance Statement**

This study provides critical insights into the extent of groundwater contamination near dumpsites in Port Harcourt, emphasizing the severe health and environmental risks posed by improper waste disposal. By analyzing spatial and seasonal variations in WQI, this research identifies key pollution hotspots and underscores the role of hydrogeological factors in contamination spread. These findings contribute to the growing body of global research on landfill-induced groundwater pollution and reinforce the urgency of adopting sustainable waste management strategies. The

study's significance extends beyond academic research, serving as a valuable resource for policymakers, environmental agencies, and public health authorities. By establishing clear correlations between landfill leachates and deteriorating water quality, this research informs evidence-based interventions to protect groundwater resources. Ultimately, the study advocates for stricter environmental regulations, improved waste disposal practices, and enhanced public awareness to safeguard water security in affected communities. Thus, graphically it is represented (Figure 1 below) as:



**Figure 1:** Understanding Groundwater Contamination Dynamics

**Disclosure statement:**

The authors declare no conflict of interest.

**Funding:**

The author(s) received no financial support for the research, authorship, and/or publication of this article.

**Authors Contribution:**

All authors contributed equally to conceptualization, validation, writing review and editing.

**Acknowledgments**

The authors would like to express their appreciation to all anonymous reviewers, for feedback and discussions that helped to substantially improve this manuscript

**References**

1. Clinton-Ezekwe I, Raimi MO, Ezekwe IC, Osu CI, Ordinioha B (2024) Ensuring Safety: Groundwater Quality and Its Potential Health Effects in the Mgbede Oil Fields Environment of South-South Nigeria. JMIR Preprints. 14/07/2024:64294. DOI: [10.2196/preprints.64294](https://doi.org/10.2196/preprints.64294). URL: <https://preprints.jmir.org/preprint/64294>.
2. Raimi, MO., Sawyerr, HO., Ezekwe, IC., & Gabriel, S. (2022). Toxicants in Water: Hydrochemical Appraisal of Toxic Metals Concentration and Seasonal Variation in Drinking Water Quality in Oil and Gas Field Area of Rivers State, Nigeria. In P. H. Saleh, & P. A. I. Hassan (Eds.), Heavy Metals - New Insights [Working Title]. IntechOpen. <https://doi.org/10.5772/intechopen.102656>. ISBN 978-1-80355-526-3.
3. Morufu OR, Henry OS, Clinton IE, Gabriel S (2021) Many Oil Wells, One Evil: Potentially toxic metals concentration, seasonal variation and Human Health Risk Assessment in Drinking Water Quality in Ebocha-Obrikom Oil and Gas Area of Rivers State, Nigeria. medRxiv 2021.11.06.21266005; doi: <https://doi.org/10.1101/2021.11.06.21266005>.
4. Morufu OR, Olawale HS, Clinton IE *et al.*, (2021) Quality water not everywhere: Exploratory Analysis of Water Quality Across Ebocha-Obrikom Oil and Gas Flaring Area in the Core Niger Delta Region of Nigeria, 04 October 2021, PREPRINT (Version 1) available at Research Square [<https://doi.org/10.21203/rs.3.rs-953146/v1>].
5. Morufu R and Clinton E (2017), Assessment of Trace Elements in Surface and Ground Water Quality (2017) LAP Lambert Academic Publishing. Mauritius. ISBN: 978-3-659-38813-2. [www.omniscryptum.com](http://www.omniscryptum.com).

6. Sarah SJ, Morufu OR, Josephine DK, *et al.* (2024). Assessing Air Quality at Waste Dump Site of Plateau State University, Bokkos, Nigeria: Evaluating Socio-Economic and Health Implications for the Community. *Authorea*. July 31. DOI: [10.22541/au.172243823.33004258/v1](https://doi.org/10.22541/au.172243823.33004258/v1).
7. Raufu, YO., Olayinka, AS., Raimi, MO., Olawale, SH., & Olabisi, OL. (2023). Assessment of occupational risks of waste scavenging in Ilorin metropolis. *AfricArXiv*. <https://doi.org/10.21428/3b2160cd.ffbb315b>.
8. Rauf, YO., & Raimi, MO. (2023). Wastes, Wastes, Everywhere Not A Place to Breathe: Redressing and Undressing Ilorin and Yenagoa City. *AfricArXiv*. <https://doi.org/10.21428/3b2160cd.52bfd7dd>.
9. Yusuf OR, Opasola OA, Adewoye SO, Raimi OM and Balogun EM (2023) Assessment of Occupational Risks of Wastes Scavenging in Ilorin Metropolis. *J. of Agri Earth & Environ mental Sciences* 2(3), 01-08.
10. Raimi MO, Austin-AI, Olawale HS, Abiola OS, Abinotami WE, Ruth EE, Nimisingha DS & Walter BO (2022). Leaving No One Behind: Impact of Soil Pollution on Biodiversity in the Global South: A Global Call for Action. In: Chibueze Izah, S. (eds) *Biodiversity in Africa: Potentials, Threats and Conservation. Sustainable Development and Biodiversity*, vol 29. Springer, Singapore. [https://doi.org/10.1007/978-981-19-3326-4\\_8](https://doi.org/10.1007/978-981-19-3326-4_8).
11. Awogbami SO, Ogunyemi O, Adebayo PA, Raimi MO (2024) Protecting the Health of Black Communities: Assessing the Impact of Environmental Hazards from Gold Mining Activities on Health Outcomes among Residents of Osun State, Nigeria. *JMIR Preprints*. 15/09/2024:66508. DOI: [10.2196/preprints.66508](https://doi.org/10.2196/preprints.66508). URL: <https://preprints.jmir.org/preprint/66508>.
12. Fubara GE, Dokuboba A, Ilemi JS, Esther OA, Digha ON, Akayinaboderi AE, Enyinnaya O, Igoniama EG, Ayibatonyo MN, Morufu OR (2024). The Niger Delta is Under a Pollution Warning: Hydrocarbon profiles in crude oil polluted soil remediated with *Pleurotus ostreatus* and *Eisenia fitida*. *bioRxiv* 2024.06.04.597352; doi: <https://doi.org/10.1101/2024.06.04.597352>.
13. Fubara GE, Ukoima HN, Dokuboba A, Morufu OR (2024). Evaluating Bioremediation Strategies on Microbial Diversity in Crude Oil-Contaminated Soil Over Three to Six Months in Port Harcourt, Nigeria., 14 May 2024, PREPRINT (Version 1) available at *Research Square*. <https://doi.org/10.21203/rs.3.rs-4409893/v1>.
14. Stephen OA, Solomon OA, Henry OS, Afolabi OO, Morufu OR (2022) Comprehensive Understanding of Hydrogeochemical evaluation of seasonal variability in groundwater quality Dynamics in the Gold Mining Areas of Osun State, Nigeria. *medRxiv* 2022.11.06.22282015; doi: <https://doi.org/10.1101/2022.11.06.22282015>.
15. Awogbami SO, Solomon OA, Sawyerr OH *et al.* (2022) Comparative Assessment of Seasonal Variations in the Quality of Surface water and its associated health hazards in Gold Mining Areas of Osun State, South-West Nigeria. Preprint (Version 1) available at *Research Square* [<https://doi.org/10.21203/rs.3.rs-2245715/v1>]. 07 November 2022.
16. Morufu OR, Clinton IE, Bowale A (2021) Statistical and Multivariate Techniques to Trace the Sources of Ground Water Contaminants and Affecting Factors of Groundwater Pollution in an Oil and Gas Producing Wetland in Rivers State, Nigeria. *medRxiv* 2021.12.26.21268415;

doi: <https://doi.org/10.1101/2021.12.26.21268415>.

17. Afolabi AS, Morufu OR (2021) Investigating Source Identification and Quality of Drinking Water in Piwoyi Community of Federal Capital Territory, Abuja Nigeria., 20 July 2021, PREPRINT (Version 1) available at *Research Square* [<https://doi.org/10.21203/rs.3.rs-736140/v1>].
18. Deinkuro NS, Charles WK, Raimi MO, Nimlang HN (2021) Oil Spills in the Niger Delta Region, Nigeria: Environmental Fate of Toxic Volatile Organics. 28 June 2021, PREPRINT (Version 1) available at Research Square. DOI: <https://doi.org/10.21203/rs.3.rs-654453/v1>.
19. Enang, OT., Azeez, BO., Ogunyemi, BT., Sulayman, AA., Araromi. DO. and Raimi, MO. (2025) Revolutionizing Hemodialysis Water Quality: Development and Evaluation of TiO<sub>2</sub> Nanoparticle-Enhanced Microporous Filters. *Advances in Nanoparticles*, 14, 12-36. <https://doi.org/10.4236/anp.2025.141002>.  
<https://www.scirp.org/journal/anp> ISSN Online: 2169-0529 ISSN Print: 2169-0510.
20. Fubara, G., Amachree, D., Soberekon, I., Akhigbe, E., Nicholas, D., Eli, A., Okoro, E., Gamage, I., Nathaniel, A. and Raimi, M. (2025) From Crisis to Recovery: Addressing Hydrocarbon Pollution in Niger Delta Soils Treated with *Pleurotus ostreatus* and *Eisenia fitida*. *Open Journal of Yangtze Oil and Gas*, 10, 1-29. doi: [10.4236/ojogas.2025.101001](https://doi.org/10.4236/ojogas.2025.101001).
21. Clinton-Ezekwe, I., Raimi, M, Ezekwe, I., Osu, I. and Ordinioha, B. (2024) From Oil to Health: Groundwater Quality and Its Potential Health Effects in Mgbede Oil Fields of South-South Nigeria. *Open Journal of Yangtze Oil and Gas*, 9, 95-118. doi: [10.4236/ojogas.2024.94006](https://doi.org/10.4236/ojogas.2024.94006).
22. Evans FG, Nkalo UH, Amachree D, Raimi MO (2024). From Killer to Solution: Evaluating Bioremediation Strategies on Microbial Diversity in Crude Oil-Contaminated Soil over Three to Six Months in Port Harcourt, Nigeria. *Adv Environ Eng Res* 5(4): 023; doi:10.21926/aeer.2404023.
23. Omoyajowo, KO; Raimi, MO; Omoyajowo, KA; Makengo, MB; Adegboyso, S; Innocent, DC; Oni, S; Oguntuyi, J; Oyediran, A; Ogunyebi, A; Kakwi, D (2024). Towards a Reduced Pollution Society: Systematic Review on the Role of Storytelling, Social Media, Humor and Celebrities' Influence for Research Communication. *J. Appl. Sci. Environ. Manage.* 28 (2) 603-623. <https://doi.org/10.4314/jasem.v28i2.34>.  
<https://www.ajol.info/index.php/jasem/article/view/266677>.
24. Okoyen E, Raimi MO, Omidiji AO, Ebuete AW (2020). Governing the Environmental Impact of Dredging: Consequences for Marine Biodiversity in the Niger Delta Region of Nigeria. *Insights Mining Science and technology* 2020; 2(3): 555586. DOI: 10.19080/IMST.2020.02.555586. <https://juniperpublishers.com/imst/pdf/IMST.MS.ID.555586.pdf>.
25. Akhigbe EO, Izah SC, Ogidi OI, Iyingiala A-A, and Raimi MO (2025) The Role of Edible Mushrooms in Immune Support. In: S.C. Izah et al. (eds.), *Bioactive Compounds in Edible Mushrooms*, Reference Series in Phytochemistry, [https://doi.org/10.1007/978-3-031-52642-8\\_21-1](https://doi.org/10.1007/978-3-031-52642-8_21-1).
26. Akayinaboderi AE, Tano DA, Okoro E, Adedoyin OO, Digha ON, Meshach OO, Ilemi JS, Tamuno TT, Igoniama EG, Charles O, Perekibina AB, Douye PM, Morufu OR (2024) World Environment Day 2024 Initiatives in Bayelsa State:



- Promoting Environmental Stewardship and Sustainable Practices - A Collaboration Between Federal University Otuoke, Niger Delta University, and the Nigerian Environmental Society. *Sustainability in Environment*. Vol. 10, No. 1, 2025. doi:10.22158/se.v10n1p1 URL: <http://dx.doi.org/10.22158/se.v10n1p1>. ISSN 2470-637X (Print) ISSN 2470-6388 (Online). [www.scholink.org/ojs/index.php/se](http://www.scholink.org/ojs/index.php/se).
27. Tamaraukepreye CO, Adams OI, Bukola OA, Ayotunde SK, Sylvester CI, Morufu OR, and Matthew CO (2024) Socioeconomic Values of Herbal Medicine. In: Izah, S.C., Ogwu, M.C., Akram, M. (eds), *Herbal Medicine Phytochemistry, Reference Series in Phytochemistry*, Springer, Cham. [https://doi.org/10.1007/978-3-031-21973-3\\_1-1](https://doi.org/10.1007/978-3-031-21973-3_1-1).
  28. Sylvester CI, Odangowei IO, Matthew CO, Saoban SS, Zaharadeen MY, Muhammad A, Morufu OR, and Austin-Asomeji I (2023) Historical Perspectives and Overview of the Value of Herbal Medicine. In: Izah, S.C., Ogwu, M.C., Akram, M. (eds), *Herbal Medicine Phytochemistry, Reference Series in Phytochemistry*, Springer, Cham. [https://doi.org/10.1007/978-3-031-21973-3\\_1-1](https://doi.org/10.1007/978-3-031-21973-3_1-1).
  29. Saliu, AO., Komolafe, OO., Bamidele, CO., Raimi, MO. (2023). The Value of Biodiversity to Sustainable Development in Africa. In: Izah, S.C., Ogwu, M.C. (eds) *Sustainable Utilization and Conservation of Africa's Biological Resources and Environment. Sustainable Development and Biodiversity*, vol 888. Springer, Singapore. [https://doi.org/10.1007/978-981-19-6974-4\\_10](https://doi.org/10.1007/978-981-19-6974-4_10).
  30. Raimi MO, Abiola OS, Atoyebi B, Okon GO, Popoola AT, Amuda-KA, Olakunle L, Austin-AI & Mercy T. (2022). The Challenges and Conservation Strategies of Biodiversity: The Role of Government and Non-Governmental Organization for Action and Results on the Ground. In: Chibueze Izah, S. (eds) *Biodiversity in Africa: Potentials, Threats, and Conservation. Sustainable Development and Biodiversity*, vol 29. Springer, Singapore. [https://doi.org/10.1007/978-981-19-3326-4\\_18](https://doi.org/10.1007/978-981-19-3326-4_18).
  31. Akayinaboderi AE, Tano DA, Okoro E, Adedoyin OO, Digha ON, Meshach OO, Ilemi JS, Tamuno TT, Igoniama EG, Charles O, Perekibina AB, Douye PM, Morufu OR (2024) World Environment Day 2024 Initiatives in Bayelsa State: Promoting Environmental Stewardship and Sustainable Practices - A Collaboration Between Federal University Otuoke, Niger Delta University, and the Nigerian Environmental Society. *Advance*. September 15, 2024. DOI: [10.31124/advance.172641767.75046547/v1](https://doi.org/10.31124/advance.172641767.75046547/v1).
  32. Raheem WB, Fadina OO, Idowu OO, Raimi MO, Austin-Asomeji I (2023) The Application of Biomaterials in Ecological Remediation of Land Pollution: Bioremediation of Heavy Metals in Cement Contaminated Soil Using White-Rot Fungus *Pleurotus sajor-caju*, 11 January 2023, PREPRINT (Version 1) available at Research Square [<https://doi.org/10.21203/rs.3.rs-2459820/v1>].
  33. Jacob OA, Anuoluwa OE and Raimi MO (2023) The notorious daredevils: potential toxic levels of cyanide and heavy metals in cassava flour sold in selected markets - taken Oke Ogun Community, Oyo State as an example. *Front. Sustain. Food Syst.* 7:1165501. doi: 10.3389/fsufs.2023.1165501. [http://journal.frontiersin.org/article/10.3389/fsufs.2023.1165501/full?utm\\_source=Email\\_to\\_authors&utm\\_medium=Email&utm\\_content=T1\\_1](http://journal.frontiersin.org/article/10.3389/fsufs.2023.1165501/full?utm_source=Email_to_authors&utm_medium=Email&utm_content=T1_1)

[1.5e1\\_author&utm\\_campaign=Email\\_publica.](#)

34. ModupeAO, Adebayo O, Sawyerr OH, Opasola AO, Morufu OR (2022) Moving from Total Concentrations to Measures of Harm in Grain Sold at Selected Markets of Southwest Nigeria. medRxiv 2022.12.18.22283634; doi: <https://doi.org/10.1101/2022.12.18.22283634>.
35. Modupe AO, AO, Sawyerr OH, Morufu OR (2022) Searching for What You Can't See - Evaluation of Pesticide Residues in Grain Sold at Selected Markets of Southwest Nigeria. medRxiv 2022.12.09.22283068; doi: <https://doi.org/10.1101/2022.12.09.22283068>.
36. Modupe AO, Adebayo O, Sawyerr OH, Opasola AO, Morufu OR (2023) Concentrations of Pesticides Residues in Grain Sold at Selected Markets of Southwest Nigeria. *Natural Resources for Human Health*. 1-15 <https://doi.org/10.53365/nrfhh/171368>. eISSN: 2583-1194.
37. Kader, S; Raimi, MO; Spalevic, V; Iyingiala, A-A; Bukola, RW; Jaufer, L; and Butt, TE. (2023) "A concise study on essential parameters for the sustainability of Lagoon waters in terms of scientific literature," *Turkish Journal of Agriculture and Forestry*: Vol. 47: No. 3, Article 3. <https://doi.org/10.55730/1300-011X.3087>. Available at: <https://journals.tubitak.gov.tr/agriculture/vol47/iss3/3>.
38. Oshatunberu, MA., Oladimeji, A., Sawyerr, OH., & Raimi, MO. (2023). Searching for What You Can't See - Evaluation of Pesticide Residues in Grain Sold at Selected Markets of Southwest Nigeria. *Current Research in Public Health*, 3(1), 10–36. Retrieved from <https://www.scipublications.com/journal/index.php/crph/article/view/566>.
39. Asiegbu OV, Ezekwe IC, Raimi MO (2022). Assessing pesticides residue in water and fish and its health implications in the Ivo river basin of South-eastern Nigeria. *MOJ Public Health*. 2022;11(4):136–142. DOI: 10.15406/mojph.2022.11.00390.
40. Clinton-Ezekwe IC, Osu IC, Ezekwe IC, Raimi MO (2022). Slow death from pollution: potential health hazards from air quality in the mgbede oil fields of south-south Nigeria. *Open Access J Sci*. 2022;5(1):61–69. DOI: 10.15406/oajs.2022.05.00177.
41. Olalekan MR, Abiola I, Ogah A, Dodeye EO (2021) Exploring How Human Activities Disturb the Balance of Biogeochemical Cycles: Evidence from the Carbon, Nitrogen and Hydrologic Cycles. *Research on World Agricultural Economy*. Volume 02, Issue 03. DOI: <http://dx.doi.org/10.36956/rwae.v2i3.426>. <http://ojs.nassg.org/index.php/rwae>.
42. Morufu OR, Tonye VO & Adedoyin OO (2021) Creating the Healthiest Nation: Climate Change and Environmental Health Impacts in Nigeria: A Narrative Review. *Scholink Sustainability in Environment*. ISSN 2470-637X (Print) ISSN 2470-6388 (Online) Vol. 6, No. 1, 2021 [www.scholink.org/ojs/index.php/se](http://www.scholink.org/ojs/index.php/se). URL: <http://dx.doi.org/10.22158/se.v6n1p61>. <http://www.scholink.org/ojs/index.php/se/article/view/3684>.
43. Morufu OR (2021). "Self-reported Symptoms on Farmers Health and Commonly Used Pesticides Related to Exposure in Kura, Kano State, Nigeria". *Annals of Community Medicine & Public Health*. 1(1): 1002. <http://>



[www.remedypublications.com/open-access/self-reported-symptoms-on-farmers-health-and-commonly-used-pesticides-related-6595.pdf](http://www.remedypublications.com/open-access/self-reported-symptoms-on-farmers-health-and-commonly-used-pesticides-related-6595.pdf).  
<http://www.remedypublications.com/annals-of-community-medicine-public-health-home.php>.

44. Isah, HM., Sawyerr, HO., Raimi, MO., Bashir, B. G., Haladu, S. & Odipe, O. E. (2020). Assessment of Commonly Used Pesticides and Frequency of Self-Reported Symptoms on Farmers Health in Kura, Kano State, Nigeria. *Journal of Education and Learning Management (JELM)*, HolyKnight, vol. 1, 31-54. doi.org/10.46410/jelm.2020.1.1.05. <https://holynight.co.uk/journals/jelm-articles/>.
45. Olalekan RM, Muhammad IH, Okoronkwo UL, Akopjubaro EH (2020). Assessment of safety practices and farmer's behaviors adopted when handling pesticides in rural Kano state, Nigeria. *Arts & Humanities Open Access Journal*. 2020;4(5):191–201. DOI: 10.15406/ahoaj.2020.04.00170.
46. Isah HM, Raimi MO, Sawyerr HO, Odipe OE, Bashir BG, Suleiman H (2020) Qualitative Adverse Health Experience Associated with Pesticides Usage among Farmers from Kura, Kano State, Nigeria. *Merit Research Journal of Medicine and Medical Sciences* (ISSN: 2354-323X) Vol. 8(8) pp. 432-447, August, 2020. DOI: 10.5281/zenodo.4008682. <https://meritresearchjournals.org/mms/content/2020/August/Isah%20et%20al.htm>.
47. Raimi MO, Bilewu OO, Adio ZO, Abdulrahman H (2019) Women Contributions to Sustainable Environments in Nigeria. *Journal of Scientific Research in Allied Sciences*. 5(4), 35-51. ISSN NO. 2455-5800. DOI No. 10.26838/JUSRES.2019.5.4.104.
48. Raimi MO, Suleiman RM, Odipe OE, Salami JT, Oshatunberu M, et al (2019). Women Role in Environmental Conservation and Development in Nigeria. *Ecology & Conservation Science*; 1(2): DOI: 10.19080/ECO.A.2019.01.555558. Volume 1 Issue 2 - July 2019. <https://juniperpublishers.com/ecoa/pdf/ECO.A.MS.ID.555558.pdf>.
49. Olalekan RM, Omidiji AO, Williams EA, Christianah MB, Modupe O (2019). The roles of all tiers of government and development partners in environmental conservation of natural resource: a case study in Nigeria. *MOJ Ecology & Environmental Sciences* 2019;4(3):114–121. DOI: 10.15406/mojes.2019.04.00142.
50. Raimi MO, and Sabinus CE (2017) Influence of Organic Amendment on Microbial Activities and Growth of Pepper Cultured on Crude Oil Contaminated Niger Delta Soil. *International Journal of Economy, Energy and Environment*. Vol. 2, No. 4, 2017, pp. 56-76. DOI: 10.11648/j.ijeee.20170204.12.
51. Enang OT, Azeez BO, Ogunyemi BT, Sulayman AA, Araromi DO, Raimi MO (2025) Innovative Water Purification for Hemodialysis: TiO<sub>2</sub> Nanoparticle-Based Microporous Filter Development and Analysis. *JMIR Preprints*. 27/01/2025:71835. DOI: [10.2196/preprints.71835](https://preprints.jmir.org/preprint/71835). URL: <https://preprints.jmir.org/preprint/71835>.
52. Olalekan AS, Adewoye SO, Henry SO, Olaniyi OA, Raimi MO (2023). Comprehensive understanding of hydrogeochemical evaluation of seasonal variability in groundwater quality dynamics in the gold mining areas of Osun state, Nigeria. *Int J Hydro*. 2023;7(5):206–220. DOI:

10.15406/ijh.2023.07.00359.

53. Raimi, MO., Oyeyemi, AS., Mcfubara, KG., Richard, GT., Austin-Asomeji, I., Omidiji, AO. (2023). Geochemical Background and Correlation Study of Ground Water Quality in Ebocha-Obrikom of Rivers State, Nigeria. *Trends Appl. Sci. Res*, 18(1), 149-168. <https://doi.org/10.17311/tasr.2023.149.168>.
54. Olalekan RM, Bukola RW, Omowunmi FO, Omowumi IO, Iyingiala A-A (2023). The application of biomaterials in ecological remediation of land pollution: bioremediation of heavy metals in cement contaminated soil using white-rot fungus *Pleurotus sajor-caju*. *J Environ Chem Toxicol*;7(1):1-6.
55. Glory R, Sylvester CI, Morufu OR and AustinAsomeji I (2023) Public and environmental health implications of artisanal petroleum refining and risk reduction strategies in the Niger Delta region of Nigeria. *Journal of Biological Research & Biotechnology*. Vol. 21 No.1; pp. 1836-1851. ISSN (print):1596-7409; eISSN (online):2705-3822. <https://dx.doi.org/10.4314/br.v20i3.6>. <http://www.bioresearch.com.ng>. Publisher: Faculty of Biological Sciences, University of Nigeria, Nsukka, Nigeria.
56. Stephen OA, Olayinka AS, Olawale HS, Olalekan RM (2023). Comparative Assessment of Seasonal Variations in the Quality of Surface Water and Its Associated Health Hazards in Gold Mining Areas of Osun State, South-West Nigeria. *Adv Environ Eng Res*; 4(1): 011; doi:10.21926/aeer.2301011.
57. Ifeanyichukwu CE, Christian LO, Morufu OR, Iyingiala AA (2022) Hydrocarbon-Based Contaminants in Drinking Water Sources and Shellfish in the Soku Oil and Gas Fields of South-South Nigeria. *Open Journal of Yangtze Gas and Oil*, 7, ISSN Online: 2473-1900 ISSN Print: 2473-1889. doi: [10.4236/ojogas.2022.74012](https://doi.org/10.4236/ojogas.2022.74012). <https://www.scirp.org/journal/ojogas>.
58. Olalekan MR, Albert O, Iyingiala AA, Sanchez DN, Telu M (2022) An environmental/scientific report into the crude oil spillage incidence in Tein community, Biseni, Bayelsa state Nigeria. *J Environ Chem Toxicol*. 2022;6(4):01-06., doi:10.37532/pulject.2022.6(4);01-06.
59. Raimi, O., Ezekwe, C., Bowale, A. and Samson, T. (2022) Hydrogeochemical and Multivariate Statistical Techniques to Trace the Sources of Ground Water Contaminants and Affecting Factors of Groundwater Pollution in an Oil and Gas Producing Wetland in Rivers State, Nigeria. *Open Journal of Yangtze Oil and Gas*, 7, 166-202. doi: [10.4236/ojogas.2022.73010](https://doi.org/10.4236/ojogas.2022.73010).
60. Olalekan MR, Olawale HS, Clinton IE and Opasola AO (2022) Quality Water, Not Everywhere: Assessing the Hydrogeochemistry of Water Quality across Ebocha-Obrikom Oil and Gas Flaring Area in the Core Niger Delta Region of Nigeria. *Pollution*, 8(3): 751-778.
61. Raimi, M. and Sawyerr, H. (2022) Preliminary Study of Groundwater Quality Using Hierarchical Classification Approaches for Contaminated Sites in Indigenous Communities Associated with Crude Oil Exploration Facilities in Rivers State, Nigeria. *Open Journal of Yangtze Oil and Gas*, 7, 124-148. doi: [10.4236/ojogas.2022.72008](https://doi.org/10.4236/ojogas.2022.72008).
62. Raimi OM, Sawyerr OH, Ezekwe CI, Gabriel S (2022) Many oil wells, one evil: comprehensive assessment of toxic metals concentration, seasonal

- variation and human health risk in drinking water quality in areas surrounding crude oil exploration facilities in rivers state, Nigeria. *International Journal of Hydrology*. 2022;6(1):23–42. DOI: 10.15406/ijh.2022.06.00299.
63. Afolabi AS, Raimi MO (2021) When Water Turns Deadly: Investigating Source Identification and Quality of Drinking Water in Piwoyi Community of Federal Capital Territory, Abuja Nigeria. *Online Journal of Chemistry*, 2021, 1, 38-58; DOI: 10.31586/ojc.2021.010105. [www.scipublications.org/journal/index.php/ojc](http://www.scipublications.org/journal/index.php/ojc).
64. Suleiman RM, Raimi MO and Sawyerr HO (2019) A Deep Dive into the Review of National Environmental Standards and Regulations Enforcement Agency (NESREA) Act. *International Research Journal of Applied Sciences*. pISSN: 2663-5577, eISSN: 2663-5585. DOI No. Irjas.2019.123.123. [www.scirange.com](http://www.scirange.com). <https://scirange.com/abstract/irjas.2019.108.125>.
65. Raimi MO, Omidiji AO, Adeolu TA, Odipe OE and Babatunde A (2019) An Analysis of Bayelsa State Water Challenges on the Rise and Its Possible Solutions. *Acta Scientific Agriculture* 3.8 (2019): 110-125. DOI: 10.31080/ASAG.2019.03.0572.
66. Olalekan RM, Dodeye EO, Efegbere HA, Odipe OE, Deinkuro NS, Babatunde A and Ochayi EO (2020) Leaving No One Behind? Drinking-Water Challenge on the Rise in Niger Delta Region of Nigeria: A Review. *Merit Research Journal of Environmental Science and Toxicology* (ISSN: 2350-2266) Vol. 6(1): 031-049 DOI: [10.5281/zenodo.3779288](https://doi.org/10.5281/zenodo.3779288).
67. Olalekan RM, Adedoyin OO, Ayibatombira A, et al (2019). "Digging deeper" evidence on water crisis and its solution in Nigeria for Bayelsa state: a study of current scenario. *International Journal of Hydrology*. 2019;3(4):244–257. DOI: 10.15406/ijh.2019.03.00187.
68. Raimi MO, Abdulraheem AF, Major I, Odipe OE, Isa HM, Onyeche C (2019). The Sources of Water Supply, Sanitation Facilities and Hygiene Practices in an Island Community: Amassoma, Bayelsa State, Nigeria. *Public Health Open Access* 2019, 3(1): 000134. ISSN: 2578-5001. DOI: 10.23880/phoa-16000134.
69. Odipe OE, Raimi MO, Suleiman F (2018). Assessment of Heavy Metals in Effluent Water Discharges from Textile Industry and River Water at Close Proximity: A Comparison of Two Textile Industries from Funtua and Zaria, North Western Nigeria. *Madridge Journal of Agriculture and Environmental Sciences*. 2018; 1(1): 1-6.doi: 10.18689/mjaes-1000101. <https://madridge.org/journal-of-agriculture-and-environmental-sciences/mjaes-1000101.php>.
70. Olalekan RM, Vivien OT, Adedoyin OO, et al. (2018). The sources of water supply, sanitation facilities and hygiene practices in oil producing communities in central senatorial district of Bayelsa state, Nigeria. *MOJ Public Health*. 2018;7(6):337–345. DOI: 10.15406/mojph.2018.07.00265.
71. Premoboere EA and Raimi MO (2018). Corporate Civil Liability and Compensation Regime for Environmental Pollution in the Niger Delta. *International Journal of Recent Advances in Multidisciplinary Research Vol. 05, Issue 06, pp. 3870-3893, June, 2018*.
72. Olalekan, RM., Omidiji, AO., Nimsingha, D., Odipe, OE. and Olalekan, AS. (2018). Health Risk Assessment on Heavy Metals Ingestion through

- Groundwater Drinking Pathway for Residents in an Oil and Gas Producing Area of Rivers State, Nigeria. *Open Journal of Yangtze Gas and Oil*, 3, 191-206. <https://doi.org/10.4236/ojogas.2018.33017>.
73. Raimi MO, and Sabinus CE (2017) Influence of Organic Amendment on Microbial Activities and Growth of Pepper Cultured on Crude Oil Contaminated Niger Delta Soil. *International Journal of Economy, Energy and Environment*. Vol. 2, No. 4, 2017, pp. 56-76. DOI: 10.11648/j.ijeee.20170204.12.
74. Raimi MO and Sabinus CE (2017) An Assessment of Trace Elements in Surface and Ground Water Quality in the Ebocha-Obrikom Oil and Gas Producing Area of Rivers State, Nigeria. *International Journal for Scientific and Engineering Research (Ijser)*: Volume 8, Issue 6, July Edition. ISSN: 2229-5518.
75. Raimi, MO, Pigha, TK and Ochayi, EO (2017) Water-Related Problems and Health Conditions in the Oil Producing Communities in Central Senatorial District of Bayelsa State. *Imperial Journal of Interdisciplinary Research (IJIR) Vol-3, Issue-6, ISSN: 2454-1362*.
76. Raimi MO, Clinton IE, Olawale HS (2021) Problematic Groundwater Contaminants: Impact of Surface and Ground Water Quality on the Environment in Ebocha-Obrikom Oil and Gas Producing Area of Rivers State, Nigeria. Oral Presentation Presented at the *United Research Forum*. 2<sup>nd</sup> International E-Conference on Geological and Environmental Sustainability during July 29-30, 2021.
77. Raimi MO (2019) 21<sup>st</sup> Century Emerging Issues in Pollution Control. 6<sup>th</sup> Global Summit and Expo on Pollution Control May 06-07, 2019 Amsterdam, Netherlands.
78. Ugwoha, E., & Emete, K. (2015). Effects of Open Dumpsite Leachate on Groundwater Quality: A Case Study of Alakahia Dumpsite in PortHarcourt, Nigeria. *Journal of Environmental Studies*, 1, 8. <https://doi.org/10.13188/2471-4879.1000002>.
79. Alaye AS. Bibiye; David N. Ogbonna; Sodienye A. Abere; Augusta Ayotamuno (2022) Heavy Metal Contamination and Health Risk Assessment of Groundwater Sources to Waste Dumpsites in Port Harcourt, *International Journal of Environment and Pollution Research*, Vol.10, No.2 pp.34-49.
80. Eseyin, O. Olushola, and I. Charles Osu. (2019). "Effect of Municipal Solid Waste (MSW) Leachate on Groundwater Quality in Port Harcourt, Nigeria". *Journal of Geography, Environment and Earth Science International* 19 (3):1-12. <https://doi.org/10.9734/jgeesi/2019/v19i330085>
81. Aizebeokhai, AP., & Olayinka, I. (2011). Structural and stratigraphic mapping of Emi field, offshore Niger Delta. *Journal of Geology and Mining Research*, 3(2), 25-38.
82. American Public Health Association (APHA), (2018). Standard methods for the examination of water and wastewater, 22<sup>nd</sup> edn. American Public Health Association, Washington, DC.
83. Morufu OR (2022) *Assessment of Ground Water Quality in Ebocha-Obrikom of River State, Nigeria*. Ph.D Dissertation, Department of Environmental Health Science, Faculty of Basic Medical Sciences, Kwara State University, Malete, Kwara State, Nigeria.



84. Kumar, R., Sinha, DK., & Saxena, R. (2015). Seasonal variation in groundwater quality and its correlation with water level fluctuation in Chennai, India. *Environmental Monitoring and Assessment*, 187, 654. <https://doi.org/10.1007/s10661-015-4843-8>.
85. Boateng, TK., Opoku, F., Acquah, SO., & Akoto, O. (2019). Groundwater contamination assessment of Oti landfill site in Kumasi, Ghana: A case study of heavy metal pollution. *Groundwater for Sustainable Development*, 8, 450–456. <https://doi.org/10.1016/j.gsd.2019.05.006>.
86. Muhammad, A., Khan, S., & Khan, M. A. (2015). Evaluation of groundwater vulnerability using DRASTIC index and GIS in Lahore, Pakistan. *Journal of Hydrogeology*, 23(2), 123–137.
87. Li, X., Wang, Y., Zhang, Q., & Liu, L. (2018). Assessment of groundwater quality and its influencing factors in Pengyang County, Northwest China. *Environmental Earth Sciences*, 77, 270. <https://doi.org/10.1007/s12665-018-7401-5>
88. Prasad, B., & Sangita, K. (2008). Heavy metal pollution in mine areas: A case study of an abandoned open cast mine filled with municipal waste. *Environmental Geology*, 55(7), 1537–1543. <https://doi.org/10.1007/s00254-007-1107-3>.
89. Han, Z., Ma, H., Shi, G., He, L., Wei, L., & Shi, Q. (2016). A review of groundwater contamination near municipal solid waste dumpsite sites in China. *Science of the Total Environment*, 569–570, 1255–1264. <https://doi.org/10.1016/j.scitotenv.2016.06.201>.
90. Ojekunle, OZ., Sangodoyin, AY., & Bamgbose, O. (2016). Water quality index assessment and ecological risk evaluation of heavy metals in drinking water. *Environmental Science and Pollution Research*, 23(10), 975–989. <https://doi.org/10.1007/s11356-015-4605-4>.
91. Naveedullah, Hashmi, M. Z., Yu, C., Shen, H., Duan, D., Shen, C., Lou, L., Chen, Y., & Khan, S. (2013). Risk assessment of heavy metals pollution in agricultural soils of Siling Reservoir watershed in Zhejiang Province, China. *BioMed Research International*, 2013, 590306. <https://doi.org/10.1155/2013/590306>.
92. Adimalla, N., Li P & Venkatayogi, S. (2018). *Hydrogeochemical evaluation of groundwater quality for drinking and irrigation purposes and integrated interpretation with water quality index studies*. *Environmental Processes*, 5(2), 363–383.
93. Carpenter, PJ., Ding, A., & Cheng, L. (2016). Identifying groundwater contamination using resistivity surveys at a dumpsite near Maoming, China. *Nature Education Knowledge*, 3(7), 20.
94. Batarseh, M., Imreizeeq, E., Tilev, S., Al Alaween, M., Suleiman, W., Al Remeithi, A. M., Al Tamimi, M. K., & Al Alawneh, M. (2021). Assessment of groundwater quality for irrigation in the arid regions using irrigation water quality index (IWQI) and GIS zoning maps: Case study from Abu Dhabi Emirate, UAE. *Groundwater for Sustainable Development*, 14, 100611. <https://doi.org/10.1016/j.gsd.2021.100611>
95. Armanuos, A., Negm, A., & Valeriano, OC. (2015). Groundwater quality investigation using water quality index and ARCGIS: Case study—Western Nile Delta Aquifer, Egypt. *Eighteenth International Water Technology Conference (IWTC18)*, 1–10.

96. El Osta, M., Masoud, M., Alqarawy, A., & Elsayed, S. (2022). Groundwater suitability for drinking and irrigation using water quality indices and multivariate modeling in Makkah Al Mukarramah Province, Saudi Arabia. *Water*, 14(3), 483. <https://doi.org/10.3390/w14030483>.

