

Eczema and Transepidermal Moisture Loss: A Systematic Review and Meta-Analysis

Torgyn Ibrayeva, Togzhan Algazina, Gulnar Batpenova, Dinara Azanbayeva,
Gulnaz Tourir, Issa Ardakuly Emir, Assel Iskakova, Aizhan Shakhanova

Submitted to: JMIR Dermatology
on: August 02, 2025

Disclaimer: © The authors. All rights reserved. This is a privileged document currently under peer-review/community review. Authors have provided JMIR Publications with an exclusive license to publish this preprint on its website for review purposes only. While the final peer-reviewed paper may be licensed under a CC BY license on publication, at this stage authors and publisher expressly prohibit redistribution of this draft paper other than for review purposes.

Table of Contents

Original Manuscript.....	4
---------------------------------	----------

Preprint
JMIR Publications

Eczema and Transepidermal Moisture Loss: A Systematic Review and Meta-Analysis

Torgyn Ibrayeva¹; Togzhan Algazina¹; Gulnar Batpenova¹; Dinara Azanbayeva¹; Gulnaz Touri¹; Issa Ardakuly Emir^{2,3}; Assel Iskakova³; Aizhan Shakhanova⁴

¹Department of Dermatology and Cosmetology Astana Medical University Astana KZ

²Shakarim University Semey KZ

³Department of Public Health Semey Medical University Semey KZ

⁴Department of Internal diseases and Rheumatology Semey Medical University Semey KZ

Abstract

Background: Eczema is a chronic, recurrent, inflammatory skin disease. Its main symptom is intermittent pruritus, as well as xerosis of the skin and eczematous lesions caused by the interaction of genetic and environmental factors. Risk factors for development are also associated with skin barrier defects that reduce skin hydration (capacity) and increase transepidermal water loss (TEWL). There are many studies and, in our analysis, we aimed to determine how TEWL scores differed significantly between patients with skin diseases, including eczema.

Objective: The specific aims of our systematic review and meta-analysis were:

- To determine, by analyzing studies, the change in the level of the epidermal barrier in skin diseases, namely eczema;
- To study the effect of changes in transepidermal moisture loss in patients with eczema, taking into account gender characteristics.

Methods: By conducting a systematic review and meta-analysis, we synthesized data from various studies that evaluated the association between TEWL scores and clinical severity of disease. Fifteen articles describing 1355 cases of eczema were included in our study.

Results: About one third of patients with eczema had elevated TEWL levels. We found that the difference in TEWL levels between men and women was statistically significant. In the studies included in this meta-analysis, TEWL elevation in eczema patients ranged from 13.3% to 76.7%. Among the age group 19 to 44 years - 59.8% of eczema cases had TEWL elevation in contrast to other age groups.

Conclusions: Disruption of epidermal barrier permeability in eczematous conditions may also occur in other skin diseases. We have found that improvement in TEWL levels can prevent and alleviate the clinical severity of inflammatory dermatoses.

(JMIR Preprints 02/08/2025:81724)

DOI: <https://doi.org/10.2196/preprints.81724>

Preprint Settings

1) Would you like to publish your submitted manuscript as preprint?

✓ **Please make my preprint PDF available to anyone at any time (recommended).**

Please make my preprint PDF available only to logged-in users; I understand that my title and abstract will remain visible to all users.

Only make the preprint title and abstract visible.

No, I do not wish to publish my submitted manuscript as a preprint.

2) If accepted for publication in a JMIR journal, would you like the PDF to be visible to the public?

✓ **Yes, please make my accepted manuscript PDF available to anyone at any time (Recommended).**

Yes, but please make my accepted manuscript PDF available only to logged-in users; I understand that the title and abstract will remain visible to all users.

Yes, but only make the title and abstract visible (see Important note, above). I understand that if I later pay to participate in <https://preprints.jmir.org/preprint/81724>

No. Please do not make my accepted manuscript PDF available to anyone. I understand that if I later pay to participate in <https://preprints.jmir.org/preprint/81724>

Original Manuscript

Review

Eczema and Transepidermal Moisture Loss: A Systematic Review and Meta-Analysis

Torgyn Ibrayeva¹, Togzhan Algazina¹, Gulnar Batpenova¹, Dinara Azanbayeva¹, Gulnaz Tourir¹, Issa Emir Ardakuly², Assel Iskakova³ and Aizhan Shakhanova⁴

¹ Department of Dermatology and Cosmetology, Astana Medical University; Astana, Kazakhstan

² Department of Foreign Philology, Shakarim University; 20A Glinka st., 071410, Semey, Kazakhstan

³ Department of Public Health, Semey Medical University; 103 Abai st., 18000, Semey, Kazakhstan

⁴ Department of Internal diseases and Rheumatology, Semey Medical University; 103 Abai st., 18000, Semey, Kazakhstan

*these authors contributed equally

Corresponding Author:

Torgyn Ibrayeva, MD

Department of Dermatology and Cosmetology

Astana Medical University

49A, Beybitshilik st.,

Astana, 01000

Kazakhstan,

E-mail: ibraeva.t@amu.kz

Abstract

Background: Eczema is a chronic, recurrent, inflammatory skin disease. Its main symptom is intermittent pruritus, as well as xerosis of the skin and eczematous lesions caused by the interaction of genetic and environmental factors. Risk factors for development are also associated with skin barrier defects that reduce skin hydration (capacity) and increase transepidermal water loss (TEWL). There are many studies and, in our analysis, we aimed to determine how TEWL scores differed significantly between patients with skin diseases, including eczema.

In the course of our study, we were exposed to a variety of data that supported our hypotheses.

The specific aims of our systematic review and meta-analysis were:

- To determine, by analyzing studies, the change in the level of the epidermal barrier in skin diseases, namely eczema;
- To study the effect of changes in transepidermal moisture loss in patients with eczema, taking into account gender characteristics.

Methodology: By conducting a systematic review and meta-analysis, we synthesized data from various studies that evaluated the association between TEWL scores and clinical severity of disease. Fifteen articles describing 1355 cases of eczema were included in our study.

Results: About one third of patients with eczema had elevated TEWL levels. We found that the difference in TEWL levels between men and women was statistically significant. In the studies included in this meta-analysis, TEWL elevation in eczema patients ranged from 13.3% to 76.7%. Among the age group 19 to 44 years - 59.8% of eczema cases had TEWL elevation in contrast to other age groups.

Conclusions: Disruption of epidermal barrier permeability in eczematous conditions may also occur in other skin diseases. We have found that improvement in TEWL levels can prevent and alleviate the clinical severity of inflammatory dermatoses.

Keywords: Skin; Eczema; Epidermal barrier; Transepidermal moisture loss; Systematic review; Meta-analysis.



INTRODUCTION

Eczema is an inflammatory skin disease caused by the interaction of both exogenous and endogenous risk factors, with a significant burden on the patient in terms of psychosocial and economic consequences. The prevalence of eczema differs between developing and industrialized countries [1]. Over the past three decades, prevalence rates in industrialized countries have increased to 15-30% in children and 2-10% in adults [2]. For example, among the statistical studies conducted in developing countries such as Mexico, Chile, Kenya and South-East Asia [2], there has been an increase in the prevalence of eczema in adults. However, rates are also increasing in children (6-7 years) with 84% of participating centers, especially in Western Europe, Canada, South America, Australasia and the Far East [2], older adults (13-14 years) with 58% of participating centers reporting eczema prevalence [3, 4]. According to the sources studied, authors state that in 60% of cases eczema manifests itself in a child from birth in the first months of life [2]. Therefore, the development of eczema is divided into 3 periods (infancy, childhood and adolescence/adulthood) [2].

Eczema has many causes, ranging from psychological disorders to permanent mechanical damage. Eczema is one of the most common skin diseases, but despite this, the exact etiology is not known. Although treatment is available, it is not always successful, and well-being and activities are limited due to the severe pain and irritation of the skin. Considering the pathomechanism of inflammation of eczema disease it is known that having a history of atopic dermatitis is an endogenous risk factor, whereas working in a humid environment, exposure to irritants and contact sensitization are exogenous risk factors.

Eczema does not have precise criteria for diagnosis, hence the indicators that change in eczema have been studied. The upper epidermal layer of the human skin functions as a physical and chemical barrier consisting of a brick and mortar-like structure called the stratum corneum [2, 5]. There is abundant evidence in the literature supporting the role of skin barrier dysfunction in eczema [2, 6]. TEWL assessment is one of the methods used to quantify the barrier function of the skin. Changes in epidermal lipids caused by water loss allow cracks to form in the stratum corneum, allowing the penetration of external antigens, irritants and microbial pathogens, which can provoke further inflammation [2, 7, 8].

Non-invasive tools that assess the level of hydration and the integrity of the skin barrier function have been found to be most informative for determining the severity and monitoring disease progression using various criteria, for diagnosis and follow-up of patients with eczema as medical examinations to obtain objective and reproducible results of skin barrier measurements. Thus, increased TEWL and decreased skin elasticity are characteristic features of epidermal barrier disruption in atopic eczema [9]. In the course of our study, we were exposed to a variety of data that supported our hypotheses.

The specific aims of our systematic review and meta-analysis were:

- To determine, by analyzing studies, the change in the level of the epidermal barrier in skin diseases, namely eczema;
- To study the effect of changes in transepidermal moisture loss in patients with eczema, taking into account gender characteristics.
- Patients with eczema, taking into account gender characteristics.

MATERIAL AND METGODS

We conducted this systematic review and meta-analysis in accordance with the statement of preferred reporting elements for systematic reviews and meta-analyses (PRISMA) (Figure 1). Before proceeding with the quantitative and systematic synthesis, we analyzed all studies that aimed to assess the association between eczema and measures of transepidermal moisture loss.

Search Strategy

A comprehensive search was conducted using Pubmed and Cochrane Library Pubmed, Cochrane Library, MEDLINE, Embase, Up ToDate, Research Gate, Google Scholar, and eLibrary databases. The following search criteria were applied: ["Eczema" (MeSH)] AND ["TEWL" (MeSH) OR "Transepidermal moisture loss" (MeSH)]. This search was restricted to English language studies published from inception to 15 April 2024. Subsequently, we analysed the abstracts of all identified papers to determine whether they met the inclusion criteria. Finally, we reviewed the reference lists of all eligible papers to find additional relevant articles

Inclusion Criteria

Our inclusion criteria were (i) studies that included patients with eczema, (ii) studies that assessed transepidermal moisture loss, (iii) studies that assessed transepidermal moisture loss based on standardised and validated instruments/questionnaires, and (iv) studies published in English.

Exclusion Criteria

Our exclusion criteria were (i) studies that did not report the rate of transepidermal moisture loss among patients with eczema or provided insufficient data to calculate the rate of transepidermal moisture loss, (ii) lack of full text for the full review, (iii) studies with poor methodological quality, i.e. case reports, case series and commentaries, and (iv) studies published in languages other than English.

Article selection

The initial search and selection of articles was conducted independently by the author (T.I.), who checked titles and abstracts and excluded all articles that did not fulfil the inclusion criteria. As a next step, we collected the full text of articles that were deemed suitable for publication and assessed all studies based on their design. All disagreements about the eligibility of studies were resolved through discussions. The selection process according to PRISMA guidelines is summarized in **Error: Reference source not found**Error: Reference source not found.

Studies should have provided sufficient data, such as odds ratio (OR) and 95% confidence interval (CI), so that the corresponding standard errors (SES) could be calculated. If such information was not available, we looked for crude data with the number of cases.

Data Extraction and Study Evaluation

The reviewer (T.I.) extracted data from the screened articles. All selected data were presented in a standardized form that included (i) first author's surname and year of publication, (ii) country of study origin, (iii) sample size in groups with and without eczema, if any, (iv) prevalence of TEWL scores in patients with eczema and controls, if any, (v) TEWL assessment tool, (vi) proportion of female and male patients with eczema, if any, (vii) age category. The quality of included articles was assessed using the Newcastle-Ottawa scale for non-randomized studies [10]. Studies scoring ≥ 7 on the Newcastle-Ottawa scale were considered high quality, 4-6 met the criteria of medium quality studies, and less than 4 were considered low quality studies. Each article was evaluated independently by a reviewer (T.I.). All differences in scores were confirmed in the AI discussion.

Statistical Analysis

We performed statistical analyses using Review Manager (RevMan) software version 5.3 (Copenhagen: Nordic Cochrane Centre, Cochrane Collaboration, 2014). Then, using a random-effects model, we calculated the prevalence and frequency of changes in TEWL levels in eczema patients and in the general population of study participants. An i^2 value was processed to assess study heterogeneity. We used a random-effects model when the heterogeneity of studies was large ($i^2 > 60\%$, $p < 0.05$) and a fixed-effects meta-analysis when heterogeneity was small ($i^2 < 60\%$, $p < 0.05$).

Statement of Ethics

As we did not perform any patient interventions and all information was recycled from previous scientific studies, ethical committee approval was not required.

Subgroup Analysis

Subgroup analysis was performed to examine the influence of sex, age, localization, severity

of skin manifestations, continence and skin disease on the prevalence of changes in TEWL levels among eczema patients. Data from the World Health Organization were used to classify all included studies by age category [11]. The countries considered for incidence information by publication were China, Germany, Malaysia, UK, Belgium, Denmark, USA, Sweden, South Korea, Brazil [11]. By continentally, European and American countries were covered, China and South Korea were allocated to the East Asia continent subgroup, and Malaysia was allocated to South-east Asia. Controlled studies have compared the prevalence of TEWL changes in groups of patients with and without eczema (or in remission). Regarding the type of skin disease, some studies contained data on the prevalence of TEWL changes among patients with eczema and atopic dermatitis, while other studies contained data on different skin diseases. The control group was most often comprised of patients with eczema.

RESULTS

3.1. Identification, Screening, Eligibility

A search using Pubmed, Cochrane Library, Research Gate, Google Scholar, eLibrary identified 348 studies (Error: Reference source not found). There were no studies that met the inclusion criteria in the Cochrane Library. One hundred and eighty-three studies were removed as duplicate studies. The titles and abstracts of the remaining 165 articles were selected. The full texts of 165 studies were analyzed and 90 of them were excluded for the following reasons: irrelevant results; review articles; not in English/non-Russian language. Finally, 15 articles describing 1355 cases of eczema were included in our study.

3.2. Demographic Data of the Studies

All articles presented data on TEWL levels in patients with eczema. When analyzing the reviewed publications, the authors assessed the prevalence of changes in TEWL levels, where scientific studies using modern non-invasive ways of studying skin structures were examined. Thus, our meta-analysis included 15 studies involving 1355 patients with eczema, with sample sizes ranging from 18 to 650 participants. The vast majority of studies included participants whose age was >18 years, some articles were older than 60 years, and one study did not include the age of the participants. Participants aged 0 to 70 years, whose mean age was available in the articles, were included. We categorized by age group: 0-18 "Children"; 19-44 "Adults"; 45-59 "Middle age"; 60-74 "Older age". The number of patients with eczema ranged from 13 to 103, in one study the sex ratio was not specified.

Almost all studies were designed as cohort studies. Assessing the quality of included studies using the Newcastle-Ottawa scale, we found that eight studies met the criteria for acceptable quality, the same numbers of studies were considered of good quality, and two studies were of poor quality. Study countries, geographical regions, sample size, TEWL rates among eczema patients, screening tools, study design and quality of included publications are summarized in Error: Reference source not found.

3.3 Analysis of random effects models

Error: Reference source not found summarizes the incidence of elevation in the total sample and in patients with eczema. The sample size in the fifteen studies ranged from 18 to 650 participants. The number of patients with eczema in the included studies ranged from 18 to 60. TEWL elevation in patients with eczema ranged from 9.8 % to 67.0 %, thus the minimum and maximum case rate values were 0.09 and 0.67 respectively.

Analysis of TEWL levels among eczema patients using random effects models showed that the prevalence was 77 %. The pooled OR for change in TEWL levels among eczema patients was 22.89 [19.62, 26.16] (Figure 2). There was high heterogeneity between studies ($I^2 = 99\%$ ($P < 0.00001$)). The increase in TEWL was 39.54 % in patients with eczema versus 31.74 % in patients with atopic dermatitis and this result was statistically significant ($P=0.04$). The increase in TEWL in

patients with atopic dermatitis was 24.37 % compared to 29.27 % in patients with eczema. A statistically significant detail was also observed: The age group "adults" (19-44 years) had more cases of eczema with TEWL elevation and the percentage was 59.8% in favor of this category out of 15 articles as compared to other age groups.

We demonstrated that patients with eczema had higher levels of TEWL scores. About one third of patients with eczema had elevated TEWL levels. We found that the difference in TEWL scores between men and women was statistically significant. In the studies included in this meta-analysis, TEWL elevation in eczema patients ranged from 13.3% to 76.7%.

DISCUSSION

This meta-analysis aimed to assess changes in TEWL levels among eczema patients from different countries and to elucidate associated risk factors. In addition, we searched for publications on the incidence from early infancy and childhood, where a chronic recurrent skin disease like atopic dermatitis was most commonly encountered. There was no single distinguishing sign or diagnostic laboratory test; therefore, the diagnosis was based on the totality of clinical findings. As a result of the different clinical features, the term "atopic eczema/dermatitis syndrome" was used, which was consistent with the study of our problem.

In the studies included in this systematic review, a comparison of TEWL levels in eczema patients according to age category showed that people in the "adult" category had higher TEWL levels. We hypothesize that this finding may be explained by the greater work capacity of this category. TEWL scores were positively correlated with disease severity regardless of age category, both adult and infant [12,13,14]. In addition, infants with higher TEWL scores during the first week of life predicted a higher risk of atopic dermatitis later in life [15]. Similarly, TEWL scores predicted psoriasis recurrence, i.e., psoriasis may recur earlier in subjects with higher TEWL scores [12,16]. We also surmise from most of the publications studied that the role of epidermal permeability dysfunction could serve as an indirect indicator in predicting the risk of developing eczema, as TEWL scores were higher in measurements performed on affected or adjacent areas and on unaffected skin areas in subjects with eczema. Studies demonstrating a significant increase in TEWL on intact skin of eczema patients have been attributed to impaired maturation of lamellar cells associated with a genetic predisposition to high protease production, leading to early destruction of stratum corneum desmosomes and subsequent disruption of the epidermal barrier [17]. Disruption of barrier function promotes exposure to antigens and other environmental stimuli, so that interleukins and cytokines are activated, leading to increased. In addition, studies have identified the importance of measuring the barrier function of epidermal permeability in the effectiveness of therapeutic action not only for eczema but also for some skin diseases such as atopic dermatitis, psoriasis [20, 21,22].

In atopic dermatitis (AD), TEWL was elevated in both normal-looking skin and in affected or dry skin. In dermatitis in the active phase, TEWL values increased and gradually decreased during clinical improvement [23]. Assessing skin barrier function in longitudinal dimension during therapy, measuring TEWL and percutaneous absorption of hydrocortisone, found that there was rapid improvement during treatment. The water content of the stratum corneum is also important for maintaining skin barrier function. The minimum water content of the stratum corneum to maintain intact skin is about 10%. The water content of normal skin is 15-40% [24, 25]. Dry skin in AD showed reduced water content in the stratum corneum, in contrast to normal skin or normal looking skin in AD [26]. Barrier restoration strategies delay relapses or improve eczema and other skin diseases, such as atopic dermatitis, with defective permeability barrier being risk factors for eczema [12, 27]. The difference in eczema prevalence between male and female patients was statistically significant as females had higher TEWL [12].

Analyses of the studies included in our meta-analysis showed that changes in TEWL levels are most often work-related. In some observational studies on gender differences in TEWL rates, differences in eczema prevalence between men and women are likely because they tend to have

different occupations that may influence the development of eczema [12]. Neither prior personal nor family history of allergy was associated with the prevalence of eczema, but certain occupations and frequent contact with disinfectants were independently associated with the incidence of hand eczema. Both skin surface pH and TEWL scores differed significantly between normal control groups and subjects with active hand eczema or a prior history of hand eczema [12]. In addition, when comparing the prevalence of eczema in both sexes, female patients had a statistically higher risk of developing eczema. We hypothesize that the higher TEWL observed in patients with eczema may be explained by hormonal imbalances and the fact that eczema affects adolescents and young adults at a time when they are more likely to be concerned about their health and body and more susceptible to public opinion.

When comparing patients between male and female sexes, TEWL levels and pH showed a small correlation of eczema prevalence per level higher in females than in males [28, 29, 30]. The higher prevalence of elevated TEWL levels in patients with eczema may be explained by the fact that most of the studies included in this meta-analysis were conducted in Asian and European countries. Comparison of the incidence in patients according to their continent of residence showed that hand eczema is a common dermatosis in the United States and Europe, information on the prevalence of hand eczema in China is limited. The lifetime prevalence of hand eczema was 9.7%, while the point prevalence was 5.4% [12, 31].

Also, aqueous capacitance is a widely used method that assesses stratum corneum and skin barrier function, and skin pH is a useful method for assessing skin barrier function [32, 33]. Skin has an acidic pH, which is important for maintaining skin barrier homeostasis, and an alkaline pH causes or exacerbates disease. Recovery of skin barrier function after disruption is normal at acidic pH, but recovery is difficult at neutral or alkaline pH. Skin pH is known to be elevated in AD, eczema, although no statistical correlation with clinical severity has been described, but skin pH is above normal, and alkaline pH may be an exacerbating factor [34]. Since skin pH can be influenced by many endogenous factors such as skin moisture, age and genetic predisposition, as well as exogenous factors such as detergents and cosmetics, the pH value can be both a cause and a result of epidermal abnormalities [35, 36]. Given the important role of pH in the regulation of skin barrier function [36, 37] and the likely involvement of skin barrier dysfunction in the pathogenesis of CE, we hypothesize that elevated skin pH contributes to skin barrier dysfunction in patients with CE.

This meta-analysis included 15 publications involving 1441 patients, and the authors found that approximately one in three people with eczema had elevated TEWL levels. In the pathophysiology of this disease, a defect in the skin barrier is associated with decreased ceramide levels and profilaggrin production, leading to greater TEWL and increased susceptibility to aggression, causing inflammation.

Among the publications studied, biophysical properties of non-eczematous skin in atopic and non-atopic people were characterized by localization in specific areas using non-invasive physical methods with different instruments [38, 39].

The areas studied included the widespread localization of the back of the hand, the palm surface of the forearm and the lower back, TEWL was significantly increased in all areas in atopic patients. Higher TEWL was found in the hands than in the other areas studied. This could be due to frequent exposure to water and detergents, which may have disrupted the skin barrier. In addition, in some areas of normal skin, TEWL tended to increase with increasing capacitance. A possible reason for this could be that an increase in skin moisture caused, for example, by occlusion reduced the diffusion resistance of the stratum corneum [40, 41].

In the reviewed studies, the authors repeatedly emphasized the heterogeneity of the study groups and instruments used and reported that the overall prevalence of TEWL ranged from 23% to 56%. There are several differences between these studies and our work, which are mainly related to the main objective. While we focused exclusively on the level of TEWL, Second, we included more studies in our meta-analysis due to the two-year interval between the two studies, during which a

number of additional publications appeared.

Our study has several strengths and limitations. First, we tried to minimize publication bias by strictly adhering to the rules for selecting studies for systematic reviews. Second, our meta-analysis includes studies using different non-invasive physical methods instruments from different countries and regions of the world. Third, our study focuses on assessing the prevalence of TEWL levels among eczema patients. However, there are limitations and the main one is moderate heterogeneity as a result of broad inclusion criteria. Therefore, the results of our study should be generalized with caution. Secondly, the wide range of measuring instruments may have affected the results of the study. Finally, we took into account age and sex characteristics, localization of the skin process, course or severity of the disease, occupational gender differences and sociocultural characteristics, which may also have contributed to the heterogeneity of our meta-analysis.

CONCLUSIONS

Defects in the epidermal permeable barrier can cause cutaneous inflammation. Conversely, improvement of the epidermal permeable barrier may prevent and alleviate some inflammatory dermatoses. We found that the prevalence of TEWL elevation among eczema patients was 67.7%. The lack of statistical significance may be due to the wide range of screening tools, suggesting the need for their validation in the field of dermatology. In addition, more qualitative observational studies would be desirable to establish clearer associations between different aspects of TEWL levels in patients with eczema. In addition, dermatologists working with eczema patients should be aware of their TEWL changes and be able to envisage corrective interventions to alleviate the burden of their disease.

Author Contributions: Conceptualization, T.I., T.A. and G.B.; methodology, D.A. and G.T.; investigation, T.I. and A.I.; writing—original draft preparation, I.E.A and A.S.; writing—review and editing, T.I., T.A, G.B., D.A., G.T. and I.E.A.; visualization, A.I., and A.S.; supervision, T.I. and A.S. All authors have read and agreed to the published version of the manuscript.

Acknowledgments

Not applicable.

Funding

This research has not received any specific grant from funding agencies in the public, for-profit or non-profit sectors.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

Search Strategy

Standardized search strategy for Review of the Literature on Anxiety in patients with vitiligo. Search fields: Title and Abstract only Search terms: ["Vitiligo"(MeSH)] AND ["Anxiety"(MeSH) OR "Anxiety Disorder" (MeSH) OR Anx* (title/abstract;TIAB)].

Abbreviations

The following abbreviations are used in this manuscript:

TEWL - Transepidermal water loss

PRISMA - Reporting Elements For Systematic Reviews And Meta-Analyses

OR - Odds Ratio

CI - Confidence Interval

SES - Corresponding Standard Errors

Preprint
JMIR Publications

Table 1. This is a table. Tables should be placed in the main text near to the first time they are cited.

First autor, year	Country	Continent	Instrument Used	Study Design
1. Xiaohua Wang, 2020	China	East Asia	1. GPskin Barrier® (GPower Inc., Seoul, South Korea), portable skin pH meter	cross-sectional
2. Ben Wang, 2018	China	East Asia	2. pH value, decreased water content, elevated TEWL, and impaired integrity of the stratum corneum	cohort
3. R. Voegeli, 2009	Germany	Europe	3. CR 300, PH 905, CM825, Standard D-Squame, SquameScanTM 850A	cohort
4. Kam-lun Ellis Hon , 2008	China	East Asia	4. CM 825, TM 210, NESS,SCORAD index, CDLQI	cohort
5. Hazleen Zainal, 2019	Malaysia	SouthEast Asia	5. Hanna Instruments H199181 pH meter, TEWL was measured using Tewameter TM300, and hydration measured using DermaLab Combo	cross-sectional
6. L. Ye, 2018	China	East Asia	6. Corneometer® CM 825 and Tewameter® TM300, VapoMeter SWL4001TJ	cross-sectional
A. Bornkessel, 2005	Germany	Europe	Corneometer CM 825, skin-pH-meter PH 900, e Tewameter TM 300, thermometer, D-Squames	cohort
7. Paul V. Andrew, 2024	UK	Europe	7. Corneometer CM825, AquaFlux AF200, Skin-pH-Meter 905, D-Squame scan 850A, Mexameter MK18	cohort
8. Teresa Jaeger, 2015	Belgium	Europe	8. Corneometer (Courage and Khazaka electronic GmbH, Cologne, Germany), vapometer (Delfin Technologies, Kuopio, Finland), pH meter (Courage and Khazaka electronic GmbH, Cologne, Germany)	cohort
9. Frederik Plum, 2020	Denmark	Europe	9. H-18049625, VD-2019-15, DermaLab Combo	cross-sectional
10. Erin E. Grinich, 2021	USA	America	10. Corneometer CM 825, F 200 Evaporimeter	cohort
11. Marie Loden, 1992	Sweden	Europe	11. Corneometer SM H20, Evaporimeter EP1	cohort
12. Suk-Jin Choi, 2003	South Korea	East Asia	12. Tewameter TM210, CM 820	cohort
13. Do-Won	South	East Asia	13. SM810,CM825,pH900	cohort

Kim, 2006	Korea			
14. Flavia A.S, 2012	Brazil	South America	14. Corneometer MPA 580, Tewameter TM300	cohort

Preprint
JMIR Publications

Table 2. Information about the studies included in the meta-analysis.

Study year	Number of patients	Female patients with eczema	Male patients with eczema	Age categories	Average age	Average TEWL g/(m ² h)	Average Ph
Xiaohua Wang, 2020	650	462	188	Adult	36.7 ± 0.46	12.4±1.3	5.0±0.5
Ben Wang, 2018	30	19	11	Middle age	50	28.82±7.52	5.80±0.41
R.Voegeli, 2009	18	-	-	Adult	23-31.2	12.48 ± 3.74	5.32 ± 0.68
Kam-lun Ellis Hon, 2008	44	21	23	Children	11.5 ± 4.6	7.4 ± 2.9	5.1 ± 2.4
Hazleen Zainal, 2019	48	33	15	Adult	28.46 ± 12.07	16.34	5.33
L.Ye, 2018	200	126	74	Middle age	45.24	16.35	9.5
A.Bornkessel, 2005	30	19	11	Adult	29.13	8.11±1.2	4.82
Paul.V.Andrew, 2024	23	14	9	Adult	38 ± 15	10.44 ± 3.18	4.49 ± 0.38
Teresa Jaeger, 2015	20	14	6	Adult	36.85	12.23	5.42
Frederik Plum, 2020	20	16	4	Adult	25.9	4.90	5.25
Erin E.Grinch, 2021	50	30	20	Adult	29.1	12	5.25
Marie Loden, 1992	26	9	2	Adult	29 ± 8	16.3±0.8	5.1±1.3
Suk-Jin Choi, 2003	51	22	29	Children	8.877	26.7	5.2
Do-Won Kim, 2006	25	8	17	Adult	20.5	25.14	5.568
Flavia A.S, 2012	120	66	54	Adult	22.1	10.2	7.0±0.6




Figure 1. Flow diagram PRISMA.




Figure 2. Analysis of TEWL levels.

References

1. Loman L., Brands M.J., Massella Patsea A.A.L., Politiek K., Arents B.W.M., Schuttelaar M.L.A. Lifestyle factors and hand eczema: A systematic review and meta-analysis of observational studies. *Contact Dermatitis*. **2022**, 87(3), 211-232.
2. Biagini Myers J.M., Khurana Hershey G.K. Eczema in early life: genetics, the skin barrier, and lessons learned from birth cohort studies. *J Pediatr*. **2010**, 157(5), 704-14.
3. Williams H., Stewart A., von Mutius E., Cookson W., Anderson H.R. Is eczema really on the increase worldwide? *J Allergy Clin Immunol*. **2008**, 121(4), 947–54.e15.
4. Strachan D., Sibbald B., Weiland S., Ait-Khaled N., Anabwani G., Anderson H.R. Worldwide variations in prevalence of symptoms of allergic rhinoconjunctivitis in children: the International Study of Asthma and Allergies in Childhood (ISAAC) *Pediatr Allergy Immunol*. **1997**, 8(4), 161–76.
5. Bieber T. Atopic dermatitis. *N Engl J Med*. **2008**, 358(14), 1483–94.
6. Sugarman J.L. The epidermal barrier in atopic dermatitis. *Semin Cutan Med Surg*. **2008**, 27(2), 108–14.
7. Cork M. The importance of skin barrier function. *Journal of Dermatological Treatment*. **1997**, 8(S1), S7–S13.
8. Ong P.Y., Leung D.Y. Immune dysregulation in atopic dermatitis. *Curr Allergy Asthma Rep*. **2006**, 6(5), 384–9.

9. Choi S.J., Song M.G., Sung W.T., Lee D.Y., Lee J.H., Lee E.S., Yang J.M. Comparison of Transepidermal Water Loss, Capacitance and pH Values in the Skin between Intrinsic and Extrinsic Atopic Dermatitis Patients. *J Korean Med Sci.* **2003**, *18*(1), 93-96.
10. Moher D., Liberati A., Tetzlaff J., Altman D.G., The PRISMA Group Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 2009, *6*(7), e1000097.
11. Wells G.A., Shea B., O'Connell D., Peterson J., Welch V., Losos M., Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses https://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
12. Voegeli R., Rawlings A.V., Breternitz M., Doppler S., Schreier T., Fluhr J.W. Increased stratum corneum serine protease activity in acute eczematous atopic skin. *Br J Dermatol.* **2009**, *161*(1), 70-7.
13. Wang X., Ye L., Lai Q., Wen S., Long Z., Qiu X., Elias P.M., Yang B., Man M.Q. Altered Epidermal Permeability Barrier Function in the Uninvolved Skin Supports a Role of Epidermal Dysfunction in the Pathogenesis of Occupational Hand Eczema. *Skin Pharmacol Physiol.* **2020**, *33*(2), 94-101.
14. Gupta J., Grube E., Ericksen M.B., Stevenson M.D., Lucky A.W., Sheth A.P., et al. Intrinsically defective skin barrier function in children with atopic dermatitis correlates with disease severity. *Journal of Allergy and Clinical Immunology* **2008**, *121*(3), 725–730.e2.
15. Kim D.W., Park J.Y., Na G.Y., Lee S.J., Lee W.J. Correlation of clinical features and skin barrier function in adolescent and adult patients with atopic dermatitis. *Int J Dermatol.* **2006**, *45*(6), 698–701.
16. Horimukai K., Morita K., Narita M., Kondo M., Kabashima S., Inoue E. Transepidermal water loss measurement during infancy can predict the subsequent development of atopic dermatitis regardless of filaggrin mutations. *Allergol Int.* **2016**, *65*(1), 103–8.
17. Man M.Q., Ye L., Hu L., Jeong S., Elias P.M., Lv C. Improvements in epidermal function prevent relapse of psoriasis: a self-controlled study. *Clin Exp Dermatol.* **2019**, *44*(6), 654–7.
18. Grinich E.E., Topham C., Haynes D., Chung J., Latour E., Simpson E.L. Validation of a novel patient-operated device for measuring skin barrier function in atopic dermatitis. *Skin Res Technol.* **2021**, *27*(5), 824-830.

19. Addor F.A., Takaoka R., Rivitti E.A., Aoki V. Atopic dermatitis: correlation between non-damaged skin barrier function and disease activity. *Int J Dermatol.* **2012**, 51(6), 672-6.
20. Morar N., Willis-Owen S.A.G., Moffatt M.F., Cookson W.O.C.M. The genetics of atopic dermatitis. *Journal of Allergy and Clinical Immunology.* **2006**, 118(1), 24-34.
21. Ye L., Wang Z., Li Z., Lv C., Man M.Q. Validation of GP Skin Barrier for assessing epidermal permeability barrier function and stratum corneum hydration in humans. *Skin Res Technol.* **2019**, 25(1), 25-29.
22. Andrew P.V., Pinnock A., Poyner A., Brown K., Chittock J., Kay L.J., Cork M.J., Danby S.G. Maintenance of an Acidic Skin Surface with a Novel Zinc Lactobionate Emollient Preparation Improves Skin Barrier Function in Patients with Atopic Dermatitis. *Dermatol Ther (Heidelb).* **2024**, 14(2), 391-408.
23. Montero-Vilchez T/, Sanabria-de-la-Torre R/, Sanchez-Diaz M/, Ureña-Paniego C/, Molina-Leyva A/, Arias-Santiago S. The impact of dupilumab on skin barrier function: A systematic review. *J Eur Acad Dermatol Venereol.* **2023**, 37(7), 1284-1292.
24. Aalto-Korte K. Improvement of skin barrier function during treatment of atopic dermatitis. *J Am Acad Dermatol.* **1995**, 33(6), 969-72.
25. Warner R.R., Myers M.C., Taylor D.A. Electron probe analysis of human skin: determination of the water concentration profile. *J Invest Dermatol.* **1988**, 90(2), 218-24.
26. Addor F.A., Takaoka R., Rivitti E.A., Aoki V. Atopic dermatitis: correlation between non-damaged skin barrier function and disease activity. *Int J Dermatol.* **2012**, 51(6), 672-6.
27. Lodén M., Wirén K., Smerud K., Meland N., Hønnås H., Mørk G. Treatment with a barrier-strengthening moisturizer prevents relapse of hand-eczema. An open, randomized, prospective, parallel group study. *Acta Derm Venereol.* **2010**, 90(6), 602–6.
28. Jordan L. Efficacy of a Hand Regimen in Skin Barrier Protection in Individuals With Occupational Irritant Contact Dermatitis. *J Drugs Dermatol.* **2016**, 15(11), s81-s85.
29. Jaeger T., Rothmaier M., Zander H., Ring J., Gutermuth J., Anliker M.D. Acid-coated Textiles (pH 5.5-6.5)--a New Therapeutic Strategy for Atopic Eczema? *Acta Derm Venereol.* **2015**, 95(6), 659-63.
30. Eberlein-König B., Schäfer T., Huss-Marp J., Darsow U., Möhrenschrager M., Herbert O., Abeck D., Krämer U., Behrendt H., Ring J. Skin surface pH, stratum corneum hydration,

- trans-epidermal water loss and skin roughness related to atopic eczema and skin dryness in a population of primary school children. *Acta DermVenereol.* **2000**, 80(3), 188-91.
31. Zainal H., Jamil A., Md Nor N., Tang M.M. Skin pH mapping and its relationship with transepidermal water loss, hydration and disease severity in adult patients with atopic dermatitis. *Skin Res Technol.* **2020**, 26(1), 91-98.
 32. Hon K.L., Wong K.Y., Leung T.F., Chow C.M., Ng P.C. Comparison of skin hydration evaluation sites and correlations among skin hydration, transepidermal water loss, SCORAD index, Nottingham Eczema Severity Score, and quality of life in patients with atopic dermatitis. *Am J Clin Dermatol.* **2008**, 9(1), 45-50.
 33. Kim D.W., Park J.Y., Na G.Y., Lee S.J., Lee W.J. Correlation of clinical features and skin barrier function in adolescent and adult patients with atopic dermatitis. *Int J Dermatol.* **2006**, 45(6), 698-701.
 34. Seidenari S., Giusti G. Objective assessment of the skin of children affected by atopic dermatitis: a study of pH, capacitance and TEWL in eczematous and clinically uninvolved skin. *Acta Derm Venereol.* **1995**, 75(6), 429-33.
 35. Grayson S., Johnson-Winegar A.G., Wintroub B.U., Isseroff R.R., Epstein E.H.Jr., Elias P.M. Lamellar body-enriched fractions from neonatal mice: preparative techniques and partial characterization. *J Invest Dermatol.* **1985**, 85(4), 289-94.
 36. Schmid-Wendtner M.H., Korting H.C. The pH of the skin surface and its impact on the barrier function. *Skin Pharmacol Physiol.* **2006**, 19(6), 296-302.
 37. Bornkessel A., Flach M., Arens-Corell M., Elsner P., Fluhr J.W. Functional assessment of a washing emulsion for sensitive skin: mild impairment of stratum corneum hydration, pH, barrier function, lipid content, integrity and cohesion in a controlled washing test. *Skin Res Technol.* **2005**, 11(1), 53-60.
 38. Wang B., Liu L.L., Zhao Z.T., Tu P. Impaired Skin Barrier Function and Downregulated Expression of Caspase-14 in Moderate to Severe Chronic Hand Eczema. *Dermatology.* **2018**, 234(5-6), 180-185.
 39. Lodén M., Olsson H., Axéll T., Linde Y.W. Friction, capacitance and transepidermal water loss (TEWL) in dry atopic and normal skin. *Br J Dermatol.* **1992**, 126(2), 137-41.

40. Plum F., Yüksel Y.T., Agner T., Nørreslet L.B. Skin barrier function after repeated short-term application of alcohol-based hand rub following intervention with water immersion or occlusion. *Contact Dermatitis*. **2020**, 83(3), 215-219.
41. Hachem J.P., Crumrine D., Fluhr J., Brown B.E., Feingold K.R., Elias P.M. pH directly regulates epidermal permeability barrier homeostasis, and stratum corneum integrity/cohesion. *J Invest Dermatol*. **2003**, 121(2), 345-53.